

Supplementary File 13 Seizure scenario sets and presentation order participants were randomised to and response rates

SF 13 Table 1 *Seizure scenario sets participants were randomised to complete DCEs on*

Seizure scenarios	n (%)	Presentation order	n (%)
<i>People with epilepsy (n=427)</i>			
Atypical seizure / Public typical seizure	145 (34.0%)	Standard	68 (15.9%)
		Reverse	77 (18.0%)
Home typical seizure / Atypical seizure	146 (34.2%)	Standard	71 (16.6%)
		Reverse	75 (17.6%)
Public typical seizure / Home typical seizure	136 (31.8%)	Standard	71 (16.7%)
		Reverse	65 (15.2%)
<i>Significant others (n=167)</i>			
Atypical seizure / Public typical seizure	54 (32.3%)	Standard	25 (14.9%)
		Reverse	29 (17.4%)
Home typical seizure / Atypical seizure	61 (36.5%)	Standard	31 (18.6%)
		Reverse	30 (17.9%)
Public typical seizure / Home typical seizure	52 (31.2%)	Standard	29 (17.4%)
		Reverse	23 (13.8%)


Notes: DCE, Discrete choice experiment; n, number.

SF 13 Table 2

DCE response rate by participant type and hypothetical seizure scenario


Seizure scenario	Participant type		
	<i>PWE</i> (<i>n=427</i>)	<i>Significant other</i> (<i>n=167</i>)	<i>Pooled</i> (<i>n=594</i>)
Home typical seizure	239/282 [85%]	97/113 [86%]	336/395 [85%]
Public typical seizure	252/281 [90%]	96/106 [91%]	348/387 [90%]
Atypical seizure	258/291 [89%]	105/115 [91%]	363/406 [89%]
	749/854 [88%]	298/334 [89%]	1047/1188 [88%]

Notes: DCE, Discrete choice experiment; n, number; PWE, people with epilepsy; Each participant was randomised to complete DCEs on 2/3 seizure scenarios. Each scenario involved 12 forced, pairwise choice questions.



SF 13 Figure 1. The randomisation of people with epilepsy to the different possible DCE sets and extent of fully complete responses

Notes: DCE, Discrete choice experiment; n, number; Each participant was randomised to complete DCEs on 2/3 seizure scenarios. Each scenario involved 12 forced, pairwise choice questions. Thus, a complete DCE set for a participant equated to 24 DCE choice questions being answered.



SF 13 Figure 2. The randomisation of 'significant other' participants to the different possible DCE sets and extent of fully complete responses

Notes: DCE, Discrete choice experiment; n, number; Each participant was randomised to complete DCEs on 2/3 seizure scenarios. Each scenario involved 12 forced, pairwise choice questions. Thus, a complete DCE set for a participant equated to 24 DCE choice questions being answered.