

S3. Review 1 list of excluded full texts

Exclude on language	
Author (Year)	Title
Afnan (2015)	[the Added Value of Art Therapy for Mothers with Post-Partum Depression in Arabic Society in Israel]/td>
Andaroon (2018)	Effect of individual counseling program by a midwife on anxiety during pregnancy in nulliparous women. [Persian]/td>
Azadi (2016)	Effectiveness of stress management training on stress reduction in pregnant women. [Persian]/td>
Benhajjoub (2008)	Maternal and perinatal depression and prevention: A clinical and projective approach. [French]/td>
Bowen (2008)	Optimizing maternal mental health within a primary health care centre: a model program/td>
Castanon (2008)	[Use of the Edinburgh Postnatal Depression Scale to detect postpartum depression]/td>
Chabrol (2001)	[Acceptability of antidepressants and psychotherapies in the depressions of the post-partum: inquiry to 198 delivered]/td>
Chabrol (2002)	Detection, prevention and treatment of postpartum depression: A randomized, controlled study on a sample of 859 women/td>
Chang (2008)	[Evidence-based management of perinatal depression]/td>
Cheng (2016)	[The Effects of a Mobile Application Social Support Program on Postpartum Perceived Stress and Depression]/td>
Chrzan-Detkos (2019)	The role of early psychological intervention in the prevention and treatment of postpartum depression. [Polish]/td>
Collado (2015)	Can we explain the depression of mothers who don't have a psychiatric history by the response to inhibition of the action?/td>
de Noose (2011)	Prenatal depression and therapeutic engagement: A clinical and projective approach/td>
de Sousa (2003)	Interactional orientation as an alternative of precocious parents-baby intervention in the context of maternal depression: Some initial reflections/td>
Des Rivieres-Pigeon (2000)	[The Edinburgh Postnatal Depression Scale: the validity of its Quebec version for a population low socioeconomic status mothers]/td>
Fakhri (2017)	Effect of education based on health belief model with relaxation on anxiety of nulliparous women. [Persian]/td>
Fang (2011)	Uncertainty, stress, and birth outcomes in non-hospitalized, high-risk pregnancy women: the effectiveness of health consultation/td>
Glasser (1999)	[Depression scale for research in and identification of postpartum depression]/td>
Hofecker-Fallahpour (2003)	[Group therapy for depression during early motherhood: first results of a pilot study]/td>
Hohm (2008)	In-patient mother baby therapy for postpartum mental disorders - A contribution to prevention in infancy. [German]/td>
Hok (2011)	Psychologists at home: Practices, models and issues of a preventive intervention/td>
Hornstein (2009)	Prevention of postnatal disorders starts during pregnancy! multi-professional network of expertise to encourage mentally burdened and teenage mothers and their children after birth: "perinatal prevention network in the rhine-neckar region". [German]/td>
Hribersek (2014)	Choice of psychopharmaceutical among pregnant women and nursing mothers. [Slovene]/td>
Huber (2009)	Postpartum depression should obstetric hospitals perform routine screening?. [German]/td>
Jalenques (2009)	[Post-partum blues and depression]/td>
Kapfhammer (2009)	Bipolar affective disorders in pregnancy and puerperium. New developments in diagnostics and treatment. [German]/td>
Karamoozian (2015)	The impact of prenatal group stress management therapy on anxiety in mothers and newborns' physiological parameters/td>
Kim (2016)	[Effects of a Hospital Based Follow-Up Program for Mothers with Very Low Birth Weight Infants]/td>
Kim (2018)	[Impact of Uncertainty on the Anxiety of Hospitalized Pregnant Women Diagnosed with Preterm Labor: Focusing on Mediating Effect of Uncertainty Appraisal and Coping Style]/td>
Komminaho (2003)	Postpartum depression screening at Rauma Health District mother and baby clinics/td>
Lara (2006)	Depressive symptoms in pregnancy and associated factors in patients of three health institutions in Mexico City. [Spanish]/td>
Lefebvre-Chanson (2017)	Treating postnatal depression with group therapy/td>

Luis (1998)	[Mental disorders during pregnancy, childbirth and puerperium in the region of Ribeiro Preto-SP, Brazil]/td>
Martinez (2015)	Ethnographic analysis at general practice waiting rooms: Implications for postpartum depression detection and treatment. [Spanish]/td>
Mirabella (2016)	[Effectiveness of a postnatal psychological treatment for women who had screened positive for depression]/td>
Moschner (2018)	[Treatment Satisfaction of Mothers with Postpartum Depression Concerning Circle of Security Intervention]/td>
Piacentini (2009)	Results of a manualised cognitive behavioural treatment in postnatal depression/td>
Piacentini (2011)	[Effectiveness of a manualized cognitive-behavioural intervention for postnatal depression]/td>
Pukas (2010)	Level of anxiety in parents of neonates treated in the NICU and support from family and medical staff. [Polish]/td>
Radonic (2018)	Improvement of non-institutional treatment and care for women with postpartum depression. [Croatian]/td>
Reyhani (2014)	Investigating the effects of spiritual self-care training on psychological stress of mothers with preterm infants admitted in neonatal intensive care unit. [Persian]/td>
Rincon-Pabon (2014)	[Postnatal depression in Colombian women: secondary analysis of the 2010 Colombian Demographic and Health Survey]/td>
Rojas (2015)	[Barriers restricting postpartum depression treatment in Chile]/td>
Rojas (2018)	[Screening and barriers for treatment of postpartum depression in Chilean public primary health care centers]/td>
Saias (2016)	Prevention of perinatal depression in high-risk families through home visitation: Clinical issues/td>
Tang (2019)	[Comparative study of minimally invasive technique and traditional decaying method for dental caries and the impact of dental treatment on pain and anxiety during pregnancy]/td>
Varo (2012)	Impact of a nursing intervention on pregnant women as a preventive tool for postpartum depression/td>
Yamashita (2003)	[Screening and intervention for depressive mothers of new-born infants]/td>
Yin (2017)	Impact of nucleos(t)ide analogue antiviral therapy on anxiety and depression in pregnant women with chronic hepatitis B. [Chinese]/td>
Zelkowitz (1995)	Screening for post-partum depression in a community sample/td>
Can't find full text	
Aigbe (2015)	Providers' perspectives on barriers to care, system constraints and piloting the HIV-modified centering pregnancy model among HIV-positive pregnant women/td>
Bhati (2015)	The effect of the Sleep Support for Moms Intervention on postpartum sleep and depressive symptoms. a pilot Randomized Controlled Trial/td>
Brown (2006)	Health visitors and postnatal depression: identification and practice/td>
Brown (2018)	The use of mobile technology in managing mild depressive symptomology in the Primipara of low socioeconomic status: A feasibility study/td>
Cubison (2005)	Acceptability of Using the EPDS as a Screening Tool for Postnatal Depression/td>
Davies (2004)	A first-stage evaluation of a group programme for PND/td>
Fritz (2013)	Postpartum depression screening in yemeni and punjabi immigrant women/td>
Gordon (2006)	Universal perinatal depression screening in an Academic Medical Center/td>
Gutteridge (2001)	The Tamworth Postnatal Depression Support Group/td>
Harmon (2018)	Barriers to diagnosis of peripartum depression in Latino women/td>
Harris (2017)	Barriers to and Facilitators of Perinatal Depression Screening/td>
Hayes (2010)	From 'postnatal depression' to 'perinatal anxiety and depression': key points of the National Perinatal Depression Plan for nurses and midwives in Australian primary health care settings/td>
Horowitz (1996)	Identification of symptoms of postpartum depression: linking research to practice/td>
Howland (2017)	Feasibility of a Relaxation Guided Imagery Intervention to Reduce Maternal Stress in the NICU/td>
Jallo (2009)	The biobehavioral effects of relaxation guided imagery on maternal stress/td>
Kye (2014)	Facilitating Mental Health Intervention in Home Visiting/td>
Lehman (2012)	Centering pregnancy: A combined quantitative and qualitative appraisal of women's experiences of depression and anxiety during group prenatal care/td>
Lewis (2011)	Improving the quality of perinatal mental health: a health visitor-led protocol/td>

Liebert (2018)	Postpartum depression: Screening and treatment of low income women in a primary care setting/td>
Lindsey (2012)	MotherFirst: Developing a Maternal Mental Health Strategy in Saskatchewan/td>
Marshall (2019)	Mindfulness Training among Parents with Preterm Neonates in the Neonatal Intensive Care Unit: A Pilot Study/td>
McGregor (2009)	A physician-based Cognitive Behavioral Intervention for depressed pregnant women in primary care: A pilot study/td>
Miller (2019)	Screening and Treatment After Implementation of a Universal Perinatal Depression Screening Program/td>
Mounts (2009)	Screening for maternal depression in the neonatal ICU/td>
Owen (2008)	Improving perinatal emotional well-being and mental health/td>
Pearson (2019)	Reducing Barriers That Hinder Obstetric Providers From Addressing Perinatal Depression: A Provider Education Module/td>
Poustie (1997)	Co-ordinated care for women with postnatal depression/td>
Rastle (2013)	Screening for depression at first obstetrical visits at Naval Hospital Camp Pendleton in 2008: A comparison study of active duty members and military dependent wives with spouse's deployment status included/td>
Robison (1999)	Effects of a stress management intervention on levels of stress and coping among pregnant and parenting adolescents: An investigation.(girls)/td>
Romain (1995)	Postnatal depression. Facilitating peer group support/td>
Rowe (2014)	Self-management of mild to moderate anxiety in women who have recently given birth: Development and acceptability of a theoretically sound complex intervention/td>
Rowe (2015)	Prevention of postnatal mental health problems in women: knowledge exchange in primary care in Victoria, Australia/td>
Stevens (2019)	A feasibility study of trauma-sensitive obstetric care for low-income, ethno-racial minority pregnant abuse survivors/td>
Tyldesley (2009)	Beat the Blues -- a self-help group for parents experiencing postnatal depression/td>
Webster (1997)	The impact of screening for risk factors associated with postnatal depression at the first prenatal visit/td>
Wesley (2006)	Reduce stress: a stress reduction project for pregnant black women/td>
Wilkinson (2003)	"Mad for Fitness": an exercise group to combat a high incidence of postnatal depression/td>
Perinatal loss/child outcomes	
Bagnall (2014)	Long-term follow-up of NetmumsHWD: A feasibility randomised controlled trial of telephone supported online behavioural activation for postnatal depression at 16 months post-randomisation/td>
Barlow (2015)	Paraprofessional-Delivered Home-Visiting Intervention for American Indian Teen Mothers and Children: 3-Year Outcomes From a Randomized Controlled Trial/td>
Beeber (2007)	Flying under the radar: engagement and retention of depressed low-income mothers in a mental health intervention/td>
Cohen (2019)	The process of change in psychotherapy with a pregnant patient following perinatal losses: An analysis of a case study/td>
Cote-Arsenault (2014)	Evidence-based intervention with women pregnant after perinatal loss/td>
Fletcher (2018)	Sms4 perinatal parents: Designing parenting support via text messages for mothers with severe mental illness (smi) and their partners/td>
Volpe (2017)	Assessing the feasibility and acceptability of Narrative Exposure Therapy to address IPV-related mental health in parenting and pregnant adolescents/td>
Not mental health	
Golfenshtein (2017)	Coping with the Stress in the Cardiac Intensive Care Unit: Can Mindfulness Be the Answer?/td>
Keller (2014)	Social marketing: approach to cultural and contextual relevance in a community-based physical activity intervention/td>
Mason (2010)	The role of antenatal pelvic floor muscle exercises in prevention of postpartum stress incontinence: a randomised controlled trial/td>
Melnyk (2014)	Validation and psychometric properties of the neonatal intensive care unit parental beliefs scale/td>
Ngai (2016)	The effect of telephone-based cognitive-behavioural therapy on parenting stress: A randomised controlled trial/td>
Ngai (2017)	The effect of a telephone-based cognitive behavioral therapy on quality of life: a randomized controlled trial/td>

Nolan (2009)	A pilot study of a nursing intervention protocol to minimize maternal-infant separation after Cesarean birth/td>
O'Brien (2013)	A pilot cohort analytic study of Family Integrated Care in a Canadian neonatal intensive care unit/td>
Ong (2019)	The effectiveness of a structured nursing intervention program on maternal stress and ability among mothers of premature infants in a neonatal intensive care unit/td>
Osman (2014)	Interventions to reduce postpartum stress in first-time mothers: a randomized-controlled trial/td>
Petteys (2018)	Mindfulness-Based Neurodevelopmental Care: Impact on NICU Parent Stress and Infant Length of Stay; A Randomized Controlled Pilot Study/td>
Pouyan (2019)	Effect of interpersonal psychotherapy oriented child birth education on stress and role adaptation in mothers with premature birth: A randomized clinical trial/td>
Pulliainen (2019)	Experiences of interactive ultrasound examination among women at risk of preterm birth: a qualitative study/td>
Reza (2007)	The impact of music on postoperative pain and anxiety following cesarean section/td>
Samra (2015)	Effect of Skin-to-Skin Holding on Stress in Mothers of Late-Preterm Infants: A Randomized Controlled Trial/td>
Simavli (2014)	Effect of music on labor pain relief, anxiety level and postpartum analgesic requirement: A randomized controlled clinical trial/td>
Snyder (2019)	Mental and physical support (MAPS) for moms: preliminary findings from a prenatal health support program/td>
Urizar (2019)	The SMART Moms Program: A Randomized Trial of the Impact of Stress Management on Perceived Stress and Cortisol in Low-Income Pregnant Women/td>
Urizar (2019)	Process Evaluation of a Stress Management Program for Low-Income Pregnant Women: The SMART Moms/Mamas LISTAS Project/td>
Villamizar-Carvajal (2018)	Decrease of premature mothers' stress levels in the intensive care unit/td>
Substance misuse	
Kern (2004)	Reductions in stress and depressive symptoms in mothers of substance-exposed infants, participating in a psychosocial program/td>
Not about implementing services/treatment/assessment	
Abrams (2009)	Barriers to service use for postpartum depression symptoms among low-income ethnic minority mothers in the United States/td>
Abrams (2016)	Perceptions and experiences of perinatal mental disorders in rural, predominantly ethnic minority communities in northern Vietnam/td>
Aftyka (2017)	Risk factors for the development of post-traumatic stress disorder and coping strategies in mothers and fathers following infant hospitalisation in the neonatal intensive care unit/td>
Alfayumi-Zeadna (2019)	Barriers to postpartum depression treatment among Indigenous Bedouin women in Israel: A focus group study/td>
Alhasanat (2017)	Risk For Postpartum Depression Among Immigrant Arabic Women in the United States: A Feasibility Study/td>
Allbaugh (2015)	Development of a screening and recruitment registry to facilitate perinatal depression research in obstetrics settings in the USA/td>
Armstrong (2010)	The paradox of screening: rural women's views on screening for postnatal depression/td>
Austin (2013)	The antenatal risk questionnaire (ANRQ): acceptability and use for psychosocial risk assessment in the maternity setting/td>
Battle (2013)	Perinatal antidepressant use: Understanding women's preferences and concerns/td>
Bayrampour (2017)	A qualitative inquiry on pregnant women's preferences for mental health screening/td>
Bentley (2007)	Implementing a clinical and research registry in obstetrics: overcoming the barriers/td>
Bray (2019)	The psychosocial impact of nausea and vomiting during pregnancy as a predictor of postpartum depression/td>
Byatt (2018)	How obstetric settings can help address gaps in psychiatric care for pregnant and postpartum women with bipolar disorder/td>
Chabrol (2004)	Acceptability of psychotherapy and antidepressants for postnatal depression among newly delivered mothers/td>
Da Costa (2018)	Mental health help-seeking patterns and perceived barriers for care among nulliparous pregnant women/td>
Dietsch (2011)	Australian Aboriginal kinship: a means to enhance maternal well-being/td>

El-Den (2019)	Development and Psychometric Evaluation of a Questionnaire to Measure Attitudes Toward Perinatal Depression and Acceptability of Screening: The PND Attitudes and Screening Acceptability Questionnaire (PASAQ)/td>
Elkhodr (2018)	The role of community pharmacists in the identification and ongoing management of women at risk for perinatal depression: A qualitative study/td>
Evans (2017)	Women's views on anxiety in pregnancy and the use of anxiety instruments: a qualitative study/td>
Fonseca (2015)	Women's help-seeking behaviours for depressive symptoms during the perinatal period: Socio-demographic and clinical correlates and perceived barriers to seeking professional help/td>
Foulkes (2011)	Enablers and Barriers to Seeking Help for a Postpartum Mood Disorder/td>
Gemmill (2006)	A survey of the clinical acceptability of screening for postnatal depression in depressed and non-depressed women/td>
Godderis (2009)	Applying new techniques to an old ally: a qualitative validation study of the Edinburgh Postnatal Depression Scale/td>
Goodman (2009)	Women's attitudes, preferences, and perceived barriers to treatment for perinatal depression/td>
Goodman (2013)	Pregnant African American women's attitudes toward perinatal depression prevention/td>
Gordon (2016)	Participatory design of ehealth solutions for women from vulnerable populations with perinatal depression/td>
Goyal (2015)	Postpartum Depression/td>
Grissette (2018)	Barriers to Help-Seeking Behavior Among Women With Postpartum Depression/td>
Guintivano (2018)	PPD ACT: an app-based genetic study of postpartum depression/td>
Henshaw (2013)	Treatment-seeking behaviors and attitudes survey among women at risk for perinatal depression or anxiety/td>
Holden (2019)	Brief Report. A qualitative study of maternal mental health services in New Zealand: Perspectives of Maori and Pacific mothers and midwives/td>
Jarrett (2016)	Pregnant women's experience of depression care/td>
Jesse (2008)	Barriers to seeking help and treatment suggestions for prenatal depressive symptoms: focus groups with rural low-income women/td>
Jomeen (2013)	Assessing women's perinatal psychological health: Exploring the experiences of health visitors/td>
Khajehpour (2013)	Health status of women with intended and unintended pregnancies/td>
Kingston (2014)	Public views of acceptability of perinatal mental health screening and treatment preference: a population based survey/td>
Kingston (2015)	Preferences for Mental Health Screening Among Pregnant Women: A Cross-Sectional Study/td>
Kinser (2015)	"I just start crying for no reason": the experience of stress and depression in pregnant, urban, African-American adolescents and their perception of yoga as a management strategy/td>
Lara-Cinisomo (2014)	Perinatal depression treatment preferences among Latina mothers/td>
Lasater (2019)	Integrating mental health into community-based maternal health services in Selingue, Mali/td>
Leung (2005)	Stress in women with postpartum depression: a phenomenological study/td>
Logsdon (2009)	Barriers to depression treatment in low-income, unmarried, adolescent mothers in a southern, urban area of the United States/td>
Lowenhoff (2019)	Using the TIDieR checklist to describe health visitor support for mothers with mental health problems: Analysis of a cross-sectional survey/td>
Massoudi (2007)	Screening for postnatal depression in Swedish child health care/td>
McGookin (2017)	Student midwives' awareness, knowledge, and experiences of antenatal anxiety within clinical practice/td>
Mohamed (2014)	A social and Biological Approach for Postpartum Depression in Egypt/td>
Moore (2018)	Advice for Health Care Professionals and Users: An Evaluation of Websites for Perinatal Anxiety/td>
Musyimi (2019)	Integration of Traditional Birth Attendants into Mental Healthcare: A Multistakeholder Qualitative Study Exploration/td>
Muzik (2016)	Teenage mothers of black and minority ethnic origin want access to a range of mental and physical health support: a participatory research approach/td>
Mwape (2012)	Socio-cultural factors surrounding mental distress during the perinatal period in Zambia: a qualitative investigation/td>
Negron (2013)	Social support during the postpartum period: mothers' views on needs, expectations, and mobilization of support/td>
Ng'oma (2019)	"Passing through difficult times": Perceptions of perinatal depression and treatment needs in Malawi -

	A qualitative study to inform the development of a culturally sensitive intervention/td>
Nyberg (2010)	Midwives' experience of encountering women with posttraumatic stress symptoms after childbirth/td>
Nygaard (2015)	Balancing Risk: A Grounded Theory Study of Pregnant Women's Decisions to (Dis)Continue Antidepressant Therapy/td>
Oh (2018)	Trends in depression among low-income mothers in the United States, 2005-2015/td>
Olin (2017)	Beyond Screening: A Stepped Care Pathway for Managing Postpartum Depression in Pediatric Settings/td>
Olson (2002)	Primary care pediatricians' roles and perceived responsibilities in the identification and management of maternal depression/td>
O'Mahen (2012)	Modifying CBT for Perinatal Depression: What Do Women Want?. A Qualitative Study/td>
O'Mahony (2012)	Using Critical Ethnography to Explore Issues among Immigrant and Refugee Women Seeking Help for Postpartum Depression/td>
O'Mahony (2012)	Barriers and facilitators of social supports for immigrant and refugee women coping with postpartum depression/td>
Osok (2018)	Adversities and mental health needs of pregnant adolescents in Kenya: identifying interpersonal, practical, and cultural barriers to care/td>
Pawils (2016)	Patients with Postpartum Depression in Gynaecological Practices in Germany - Results of a Representative Survey of Local Gynaecologists about Diagnosis and Management/td>
Pilkington (2016)	Enhancing reciprocal partner support to prevent perinatal depression and anxiety: a Delphi consensus study/td>
Place (2017)	Detection and care practices for postpartum depressive symptoms in public-sector obstetric units in Mexico: Qualitative results from a resource-constrained setting/td>
Ransing (2019)	Perinatal depression-knowledge gap among service providers and service utilizers in India/td>
Raymond (2014)	Addressing perinatal depression in a group of underserved urban women: a focus group study/td>
Recto (2018)	"We Don't Want to be Judged": Perceptions about Professional Help and Attitudes Towards Help-Seeking among Pregnant and Postpartum Mexican-American Adolescents/td>
Reilly (2013)	Disparities in reported psychosocial assessment across public and private maternity settings: a national survey of women in Australia/td>
Rompala (2016)	Prenatal Depression Screening by Certified Nurse-Midwives, Oregon/td>
Sanders (2006)	Attitudes, perceived ability, and knowledge about depression screening: a survey of certified nurse-midwives/certified midwives/td>
Segre (2010)	Nursing care for postpartum depression, part 1: do nurses think they should offer both screening and counseling?/td>
Segre (2010)	Screening and counseling for postpartum depression by nurses: the women's views/td>
Silverwood (2019)	Healthcare professionals' perspectives on identifying and managing perinatal anxiety: a qualitative study/td>
Skocir (2006)	Are Slovenian midwives and nurses ready to take on a greater role in caring for women with postnatal depression?/td>
Skoog (2017)	'There's something in their eyes' - Child Health Services nurses' experiences of identifying signs of postpartum depression in non-Swedish-speaking immigrant mothers/td>
Sleath (2005)	Ethnicity and depression treatment preferences of pregnant women/td>
Surjaningrum (2018)	The feasibility of a role for community health workers in integrated mental health care for perinatal depression: a qualitative study from Surabaya, Indonesia/td>
Tabb (2015)	Perinatal depression screening in a Women, Infants, and Children (WIC) program: perception of feasibility and acceptability among a multidisciplinary staff/td>
Teng (2007)	Healthcare worker's perceptions of barriers to care by immigrant women with postpartum depression: an exploratory qualitative study/td>
Thomas (2014)	Pilot early intervention antenatal group program for pregnant women with anxiety and depression/td>
Thomas (2014)	Stories of postpartum depression: exploring health constructs and help-seeking in mothers' talk/td>
Tol (2018)	Maternal mental health priorities, help-seeking behaviors, and resources in post-conflict settings: a qualitative study in eastern Uganda/td>
Vanderkruik (2019)	Perspectives on task-shifting depression care to peers for depressed Latina mothers/td>
Vigod (2016)	A Population-Based Study of Postpartum Mental Health Service Use by Immigrant Women in Ontario, Canada/td>
Walker (2013)	Maternal health needs and interest in screening for depression and health behaviors during pediatric visits/td>

Wenze (2018)	Perinatal Mental Health Treatment Needs, Preferences, and Barriers in Parents of Multiples/td>
Zittel-Palamara (2008)	Desired assistance versus care received for postpartum depression: access to care differences by race/td>
Effectiveness studies	
Abdollahpour (2016)	The effect of the magical hour on post-traumatic stress disorder (PTSD) in traumatic childbirth: a clinical trial/td>
Abdollahpour (2019)	Effect of the Sacred Hour on Postnatal Depression in Traumatic Childbirth: a Randomized Controlled Trial/td>
Abdollahpour (2019)	Effect of Brief Cognitive Behavioral Counseling and Debriefing on the Prevention of Post-traumatic Stress Disorder in Traumatic Birth: A Randomized Clinical Trial/td>
Abiodun (2006)	Postnatal depression in primary care populations in Nigeria/td>
Abrahams (2019)	Validation of a brief mental health screening tool for pregnant women in a low socio-economic setting/td>
Adams (2011)	Starting a perinatal and infant mental health service at Winnunga Nimmityjah/td>
Ahlqvist-Bjorkroth (2019)	An educational intervention for NICU staff decreased maternal postpartum depression/td>
Ahn (2004)	The effects of a home-visiting discharge education on maternal self-esteem, maternal attachment, postpartum depression and family function in the mothers of NICU infants/td>
Aite (2002)	Multidisciplinary management of fetal surgical anomalies: the impact on maternal anxiety/td>
Aite (2003)	A challenging intervention with maternal anxiety: Babies requiring surgical correction of a congenital anomaly after missed prenatal diagnosis/td>
Akbarzadeh (2018)	Investigation of breastfeeding training based on BASNEF model on the intensity of postpartum blues/td>
Alder (2002)	Counselling for postnatal depression in the voluntary sector/td>
Alder (2008)	Policy and practice in the management of postnatal depression in Scotland/td>
Alexander (2013)	Innovation:CBT-based support groups for postnatal depression/td>
Ali (2009)	Post partum anxiety and depression in peri-urban communities of Karachi, Pakistan: a quasi-experimental study/td>
Ammerman (2012)	Predictors of treatment response in depressed mothers receiving in-home cognitive-behavioral therapy and concurrent home visiting/td>
Ammerman (2013)	Treatment of depressed mothers in home visiting: impact on psychological distress and social functioning/td>
Ammerman (2013)	A clinical trial of in-home CBT for depressed mothers in home visitation/td>
Ammerman (2015)	Depression improvement and parenting in low-income mothers in home visiting/td>
An exploratory parallel-group... (Trevillion)	An exploratory parallel-group randomised controlled trial of antenatal Guided Self-Help (plus usual care) versus usual care alone for pregnant women with depression: DAWN trial/td>
Anagnostaki (2019)	Implementation of a brief early intervention in times of socio-economic crisis: effects on parental stress/td>
Ancill (1986)	Screening for antenatal and postnatal depressive symptoms in general practice using a microcomputer-delivered questionnaire/td>
Anderson (2018)	Pregnancy-related anxiety: Re-examining its distinctiveness+/td>
Anderson (2018)	Predicting posttraumatic stress and depression symptoms among adolescents in the extended postpartum period/td>
Appleby (2003)	The treatment of postnatal depression by health visitors: impact of brief training on skills and clinical practice/td>
Armstrong (1999)	A randomized, controlled trial of nurse home visiting to vulnerable families with newborns/td>
Armstrong (2004)	The effectiveness of a pram-walking exercise programme in reducing depressive symptomatology for postnatal women/td>
Armstrong (2007)	Screening for postnatal depression: not a simple task/td>
Arroyo (2013)	Before and after: A mother and infant painting group/td>
Ashby (2016)	Implications of Comprehensive Mental Health Services Embedded in an Adolescent Obstetric Medical Home/td>
Ashby (2019)	Trauma-informed care in a patient-centered medical home for adolescent mothers and their children/td>
Ashford (2018)	Feasibility and Acceptability of a Web-Based Treatment with Telephone Support for Postpartum

	Women With Anxiety: Randomized Controlled Trial/td>
Atif (2017)	Mother-to-mother therapy in India and Pakistan: adaptation and feasibility evaluation of the peer-delivered Thinking Healthy Programme/td>
Atif (2019)	Delivering maternal mental health through peer volunteers: a 5-year report/td>
Atif (2019)	Scaling-up psychological interventions in resource-poor settings: training and supervising peer volunteers to deliver the 'Thinking Healthy Programme' for perinatal depression in rural Pakistan/td>
Avalos (2016)	Improved Perinatal Depression Screening, Treatment, and Outcomes With a Universal Obstetric Program/td>
Ayers (2006)	Birth trauma. What happens after a difficult birth? Postnatal debriefing services/td>
Ayers (2015)	Brief Online Self-help Exercises for Postnatal Women to Improve Mood: A Pilot Study/td>
Babu (2018)	Sociodemographic and Medical Risk Factors Associated With Antepartum Depression/td>
Bakare (2017)	Improving access to interventions among mothers screened positive for post-partum depression (PPD) at National Programme on Immunization (NPI) clinics in south-western and south-eastern Nigeria - A service development report/td>
Baker (2005)	Prevalence of postpartum depression in a native American population/td>
Baker (2009)	Web-based training for implementing evidence-based management of postpartum depression/td>
Baker-Ericzen (2008)	Partnership for women's health: A new-age collaborative program for addressing maternal depression in the postpartum period/td>
Baker-Ericzen (2012)	A collaborative care telemedicine intervention to overcome treatment barriers for Latina women with depression during the perinatal period/td>
Balbino (2016)	Measurement of Family-centered care perception and parental stress in a neonatal unit/td>
Baldisserotto (2019)	Barriers to Seeking and Accepting Treatment for Perinatal Depression: A Qualitative Study in Rio de Janeiro, Brazil/td>
Bansal (2018)	Screening for postpartum depression in a neonatal intensive care unit and postnatal ward with its impact on newborn care practices: A hospital based survey/td>
Barnes (2009)	The utility of volunteer home-visiting support to prevent maternal depression in the first year of life/td>
Barnet (2007)	Home visiting for adolescent mothers: effects on parenting, maternal life course, and primary care linkage/td>
Barnett (1999)	Screening for postnatal depression in women of non-English speaking background/td>
Barrera (2015)	Online prevention of postpartum depression for Spanish- and English-speaking pregnant women: A pilot randomized controlled trial/td>
Barrera (2019)	A preliminary study on the acceptability of a brief SMS program for perinatal women/td>
Barry (2001)	Reducing maternal psychological distress after the NICU experience through journal writing/td>
Battle (2014)	A mother-baby psychiatric day hospital: History, rationale, and why perinatal mental health is important for obstetric medicine/td>
Battle (2015)	Potential for prenatal yoga to serve as an intervention to treat depression during pregnancy/td>
Bauer (2010)	Alleviating distress during antepartum hospitalization: a randomized controlled trial of music and recreation therapy/td>
Baumel (2018)	Digital Peer-Support Platform (7Cups) as an Adjunct Treatment for Women With Postpartum Depression: Feasibility, Acceptability, and Preliminary Efficacy Study/td>
Beattie (2017)	Effects of mindfulness on maternal stress, depressive symptoms and awareness of present moment experience: A pilot randomised trial/td>
Beckjord (2011)	Potential benefits of health information technology for integrating physical and behavioral health care: perinatal depression as a case-in-point/td>
Beddoe (2008)	Mindfulness-based yoga during pregnancy: A pilot study examining relationships between stress, anxiety, sleep, and pain/td>
Beddoe (2009)	The effects of mindfulness-based yoga during pregnancy on maternal psychological and physical distress/td>
Beeber (2010)	Short-term in-home intervention reduces depressive symptoms in Early Head Start Latina mothers of infants and toddlers/td>
Beevi (2016)	Impact of Hypnosis Intervention in Alleviating Psychological and Physical Symptoms During Pregnancy/td>
Bell (2016)	Barriers and Facilitators to the Use of Mental Health Services by Women With Elevated Symptoms of Depression and Their Partners/td>
Benediktsson (2013)	Comparing CenteringPregnancy to standard prenatal care plus prenatal education/td>

Berard (2019)	Impact of antidepressant use, discontinuation, and dosage modification on maternal depression during pregnancy/td>
Bershadsky (2014)	The effect of prenatal Hatha yoga on affect, cortisol and depressive symptoms/td>
Bhat (2018)	Text messaging to support a perinatal collaborative care model for depression: A multi-methods inquiry/td>
Bhat (2018)	Delivering perinatal depression care in a rural obstetric setting: a mixed methods study of feasibility, acceptability and effectiveness/td>
Bhusal (2016)	Validating the Edinburgh Postnatal Depression Scale as a screening tool for postpartum depression in Kathmandu, Nepal/td>
Biggs (2015)	Exploring the views and experiences of callers to the PANDA Post and Antenatal Depression Association Australian National Perinatal Depression Helpline: a cross-sectional survey/td>
Biggs (2019)	Peer supporters' experiences on an Australian perinatal mental health helpline/td>
Biggs (2019)	'I need help': Reasons new and re-engaging callers contact the PANDA-Perinatal Anxiety and Depression Australia National Helpline/td>
Bina (2018)	Nurses perceived preparedness to screen, intervene, and refer women with suspected postpartum depression/td>
Bitew (2017)	Antenatal depressive symptoms and utilisation of delivery and postnatal care: a prospective study in rural Ethiopia/td>
Bittner (2014)	Early intervention in pregnant women with elevated anxiety and depressive symptoms: efficacy of a cognitive-behavioral group program/td>
Bledsoe (2018)	Feasibility of treating depression in pregnant adolescents using brief interpersonal psychotherapy/td>
Boath (2004)	Users' views of two alternative approaches to the treatment of postnatal depression/td>
Borghini (2014)	Effects of an early intervention on maternal post-traumatic stress symptoms and the quality of mother-infant interaction: the case of preterm birth/td>
Bose (2015)	Changes in depression status in low socioeconomic perinatal subjects in rural India after supervised physical exercise: A randomized controlled study/td>
Bowen (2012)	Patterns of depression and treatment in pregnant and postpartum women/td>
Bowman (2015)	Sources Impacting Pharmacological Treatment for Anxiety and/or Depression During Pregnancy/td>
Boyd (2015)	A Pilot Investigation to Enhance Behavioral Health Referral for Perinatal, Low-Income Women With Mood Disorders/td>
Boyd (2019)	Pilot RCT of a social media parenting intervention for postpartum mothers with depression symptoms/td>
Brandon (2012)	Proof of concept: Partner-Assisted Interpersonal Psychotherapy for perinatal depression/td>
Brecht (2014)	Prevention of postpartum traumatic stress in mothers with preterm infants: Manual development and evaluation/td>
Brock (2017)	Depression Treatment by Non-Mental-Health Providers: Incremental Evidence for the Effectiveness of Listening Visits/td>
Broom (2015)	Feasibility and Perception of Using Text Messages as an Adjunct Therapy for Low-Income, Minority Mothers With Postpartum Depression/td>
Brugha (2000)	Pragmatic randomized trial of antenatal intervention to prevent post-natal depression by reducing psychosocial risk factors/td>
Brugha (2016)	Can community midwives prevent antenatal depression? An external pilot study to test the feasibility of a cluster randomized controlled universal prevention trial/td>
Buil (2019)	Kangaroo supported diagonal flexion positioning: Positive impact on maternal stress and postpartum depression risk and on skin-to-skin practice with very preterm infants/td>
Buist (2006)	Acceptability of routine screening for perinatal depression/td>
Buist (2007)	National program for depression associated with childbirth: the Australian experience/td>
Buist (2007)	Impact of education on women with perinatal depression/td>
Burns (2013)	A pilot randomised controlled trial of cognitive behavioural therapy for antenatal depression/td>
Butler (2014)	The acceptability and feasibility of the Baby Triple P Positive Parenting Programme on a mother and baby unit: Q-methodology with mothers with severe mental illness/td>
Buttner (2014)	Efficacy of yoga for depressed postpartum women: A randomized controlled trial/td>
Buultjens (2008)	A holistic programme for mothers with postnatal depression: pilot study/td>
Buzi (2015)	A Socioecological Framework to Assessing Depression Among Pregnant Teens/td>
Byatt (2016)	Improving perinatal depression care: the Massachusetts Child Psychiatry Access Project for

	Moms/td>
Byatt (2016)	PRogram in Support of Moms (PRISM): Development and beta testing/td>
Byatt (2018)	PRogram In Support of Moms (PRISM): a pilot group randomized controlled trial of two approaches to improving depression among perinatal women/td>
Byatt (2018)	Massachusetts Child Psychiatry Access Program for Moms: Utilization and Quality Assessment/td>
Callahan (2002)	Identifying mothers at risk for postnatal emotional distress: further evidence for the validity of the Perinatal Posttraumatic Stress Disorder Questionnaire/td>
Calvert (2009)	Structural family therapy as a treatment modality to decrease depressive symptoms for women suffering from postpartum depression and improve family functioning/td>
Cardone (2006)	Psychosocial assessment by phone for high-scoring patients taking the Edinburgh Postnatal Depression Scale: communication pathways and strategies/td>
Carlin (2019)	'Having a Quiet Word': Yarning with Aboriginal Women in the Pilbara Region of Western Australia about Mental Health and Mental Health Screening during the Perinatal Period/td>
Carolan (2012)	The Limerick Lullaby project: an intervention to relieve prenatal stress/td>
Carolan (2012)	Experiences of pregnant women attending a lullaby programme in Limerick, Ireland: A qualitative study/td>
Carter (2019)	Effectiveness of a peer support intervention for antenatal depression: a feasibility study/td>
Chae (2012)	Can we effectively use the two-item PHQ-2 to screen for postpartum depression?/td>
Chambers (2009)	Relaxation during pregnancy to reduce stress and anxiety and their associated complications/td>
Chang (2008)	Effects of music therapy on psychological health of women during pregnancy/td>
Chang (2015)	The effects of music listening on psychosocial stress and maternal-fetal attachment during pregnancy/td>
Chaudhry (2016)	The Cornell Peripartum Psychosis Management Tool: A case series and template/td>
Chaudron (2004)	Detection of postpartum depressive symptoms by screening at well-child visits/td>
Chen (2000)	Effects of support group intervention in postnatally distressed women. A controlled study in Taiwan/td>
Chen (2011)	Addressing maternal mental health needs in Singapore/td>
Chen (2011)	Identifying mothers with postpartum depression early: integrating perinatal mental health care into the obstetric setting/td>
Cheng (2016)	The Influence of Antenatal Partner Support on Pregnancy Outcomes/td>
Cherry (2016)	Postpartum depression screening in the Neonatal Intensive Care Unit: program development, implementation, and lessons learned/td>
Chibanda (2014)	Group problem-solving therapy for postnatal depression among HIV-positive and HIV-negative mothers in Zimbabwe/td>
Christie (2011)	The effect of health visitors' postpartum home visit frequency on first-time mothers: cluster randomised trial/td>
Chuffo (2015)	Implementation of an innovative nurse-delivered depression intervention for mothers of NICU infants/td>
Chung (2012)	Randomized non-invasive sham-controlled pilot trial of electroacupuncture for postpartum depression/td>
Clarici (2015)	Intranasal administration of oxytocin in postnatal depression: implications for psychodynamic psychotherapy from a randomized double-blind pilot study/td>
Clark (2000)	Discussing emotional health in pregnancy: the Edinburgh Postnatal Depression Scale/td>
Clarke (2014)	Impact of a participatory intervention with women's groups on psychological distress among mothers in rural Bangladesh: secondary analysis of a cluster-randomised controlled trial/td>
Clevesy (2019)	A Project to Improve Postpartum Depression Screening Practices Among Providers in a Community Women's Health Care Clinic/td>
Cluxton-Keller (2019)	Engaging rural young mothers in a technology-based intervention for depression/td>
Coates (2017)	The experiences of women who have accessed a perinatal and infant mental health service: A qualitative investigation/td>
Cobiella (1990)	A comparison of two stress-reduction treatments for mothers of neonates hospitalized in a neonatal intensive care unit/td>
Coe (2013)	Supporting women with perinatal mental health problems: the role of the voluntary sector/td>
Coffman (2019)	Postpartum Depression Screening and Referrals in Special Supplemental Nutrition Program for Women, Infant, and Children Clinics/td>

Cohen (2002)	Effects of an intervention program to decrease distress in pregnant Latina women at risk for preterm birth/td>
Cole (2018)	Universal Postpartum Mental Health Screening for Parents of Newborns With Prenatally Diagnosed Birth Defects/td>
Coll (2019)	Efficacy of Regular Exercise During Pregnancy on the Prevention of Postpartum Depression: The PAMELA Randomized Clinical Trial/td>
Cooper (2015)	Attempting to prevent postnatal depression by targeting the mother-infant relationship: a randomised controlled trial/td>
Corey (2019)	Bedside Music Therapy for Women during Antepartum and Postpartum Hospitalization/td>
Corna (2019)	Supporting maternal mental health of Rohingya refugee women during the perinatal period to promote child health and wellbeing: a field study in Cox's Bazar/td>
Corno (2018)	Effect of a web-based positive psychology intervention on prenatal well-being: A case series study/td>
Craig (2005)	Therapeutic group programme for women with postnatal depression in rural Victoria: a pilot study/td>
Crockett (2008)	A depression preventive intervention for rural low-income African-American pregnant women at risk for postpartum depression/td>
Curry (1998)	The Prenatal Psychosocial Profile: a research and clinical tool/td>
Cussino (2013)	Postnatal depression and prevention: Effects of a psychoeducational intervention. [Italian, English]/td>
Cwikel (2018)	Addressing the needs of new mothers in a multi-cultural setting: an evaluation of home visiting support for new mothers - Mom to Mom (Negev)/td>
Cynthia (2018)	Efficacy of an Internet-based depression intervention to improve rates of treatment in adolescent mothers/td>
Da Costa (2009)	A randomized clinical trial of exercise to alleviate postpartum depressed mood/td>
Dabas (2019)	Impact of audio assisted relaxation technique on stress, anxiety and milk output among postpartum mothers of hospitalized neonates: A randomized controlled trial/td>
Dagher (2009)	The psychosocial work environment and maternal postpartum depression/td>
Dagklis (2016)	Prevalence of antenatal depression and associated factors among pregnant women hospitalized in a high-risk pregnancy unit in Greece/td>
Daire (2018)	Positively impacting maternal stress and parental adjustment through community-based relationship education (re)/td>
Daley (2008)	Feasibility of an exercise intervention for women with postnatal depression: a pilot randomised controlled trial/td>
Daley (2015)	A pragmatic randomized controlled trial to evaluate the effectiveness of a facilitated exercise intervention as a treatment for postnatal depression: the PAM-PeRS trial/td>
Danaher (2012)	Web-Based Intervention for Postpartum Depression: Formative Research and Design of the MomMoodBooster Program/td>
Danaher (2013)	MomMoodBooster web-based intervention for postpartum depression: feasibility trial results/td>
Darius (2018)	Adaptation of an evidence-based postpartum depression intervention: feasibility and acceptability of mothers and babies 1-on-1/td>
Darwin (2013)	Assessment acting as intervention: Findings from a study of perinatal psychosocial assessment/td>
Davies (2018)	Psychological Interventions in Perinatal Community Mental Health Teams in the United Kingdom/td>
Davis (2014)	The feasibility of yoga in the treatment of antenatal depression and anxiety: A pilot study/td>
Davis (2015)	A randomized controlled trial of yoga for pregnant women with symptoms of depression and anxiety/td>
de Alencar (2009)	Effect of kangaroo mother care on postpartum depression/td>
de la Fe (2017)	Feasibility of screening and prevalence of prenatal depression in an obstetric setting in Spain/td>
De Schepper (2016)	Post-Traumatic Stress Disorder after childbirth and the influence of maternity team care during labour and birth: A cohort study/td>
Dennis (2003)	The effect of peer support on postpartum depression: a pilot randomized controlled trial/td>
Dennis (2004)	Can we identify mothers at risk for postpartum depression in the immediate postpartum period using the Edinburgh Postnatal Depression Scale?/td>
Dennis (2010)	Postpartum depression peer support: maternal perceptions from a randomized controlled trial/td>
Dennis (2013)	Peer support for postpartum depression: volunteers' perceptions, recruitment strategies and training from a randomized controlled trial/td>

Dennis (2014)	The process of developing and implementing a telephone-based peer support program for postpartum depression: evidence from two randomized controlled trials/td>
Dennis-Tiwary (2017)	Salutary effects of an attention bias modification mobile application on biobehavioral measures of stress and anxiety during pregnancy/td>
DeOliveira (2018)	A screening tool for perinatal mood and anxiety disorders: Culturally sensitive prenatal care for Latina mothers/td>
Deshpande (2013)	Yoga for high-risk pregnancy: a randomized controlled trial/td>
Dikmen-Yildiz (2017)	Screening for birth-related PTSD: psychometric properties of the Turkish version of the Posttraumatic Diagnostic Scale in postpartum women in Turkey/td>
Dimidjian (2015)	An open trial of mindfulness-based cognitive therapy for the prevention of perinatal depressive relapse/recurrence/td>
Dimidjian (2016)	Staying well during pregnancy and the postpartum: A pilot randomized trial of mindfulness-based cognitive therapy for the prevention of depressive relapse/recurrence/td>
Ding (2014)	Epidural labor analgesia is associated with a decreased risk of postpartum depression: a prospective cohort study/td>
Docherty (2019)	Postpartum depression screening in the first year: A cross-sectional provider analysis in Oregon/td>
Dodge (2014)	Implementation and randomized controlled trial evaluation of universal postnatal nurse home visiting/td>
Doherty (2018)	A Mobile App for the Self-Report of Psychological Well-Being During Pregnancy (BrightSelf): Qualitative Design Study/td>
Dois (2013)	The effect of Mother/Infant skin-to-skin contact on postpartum depressive symptoms in women with low obstetric risk/td>
Dolan (2019)	Pregnancy in prison, mental health and admission to prison mother and baby units/td>
Dombrowska-Pali (2018)	Music therapy - An alternative form supporting the quality of mother and baby care/td>
Dornelas (2010)	A Pilot Study of Affect-Focused Psychotherapy for Antepartum Depression/td>
Downie (2003)	Using the Edinburgh Postnatal Depression Scale to achieve best practice standards/td>
Drake (2014)	Online screening and referral for postpartum depression: an exploratory study/td>
Drozd (2015)	An Internet-Based Intervention (Mamma Mia) for Postpartum Depression: Mapping the Development from Theory to Practice/td>
Dubus (2014)	Permission to be authentic: An intervention for postpartum women/td>
Duffecy (2019)	A Group-Based Online Intervention to Prevent Postpartum Depression (Sunnyside): Feasibility Randomized Controlled Trial/td>
Dugravier (2013)	Impact of a manualized multifocal perinatal home-visiting program using psychologists on postnatal depression: the CAPEDP randomized controlled trial/td>
Edge (2011)	Managed care networks improve perinatal mental healthcare for Black and minority ethnic (BME) women?/td>
Edward (2019)	Self- screening using the Edinburgh post natal depression scale for mothers and fathers to initiate early help seeking behaviours/td>
Edwards (2008)	Antenatal psychosocial risk factors and depression among women living in socioeconomically disadvantaged suburbs in Adelaide, South Australia/td>
Effects of a mindfulness... (Lonnberg)	Effects of a mindfulness based childbirth and parenting program on pregnant women's perceived stress and risk of perinatal depression-Results from a randomized controlled trial/td>
El-Khorazaty (2007)	Recruitment and retention of low-income minority women in a behavioral intervention to reduce smoking, depression, and intimate partner violence during pregnancy/td>
Elliott (2001)	Training health visitors to reduce levels of depression after childbirth: an evaluation/td>
Emerson (2018)	Postpartum Depression Screening for New Mothers at Well Child Visits/td>
Engqvist (2013)	Involving the family in the care and treatment of women with postpartum psychosis: Swedish psychiatrists' experiences/td>
Ettenberger (2018)	Music therapy song writing with mothers of preterm babies in the Neonatal Intensive Care Unit (NICU) - A mixed-methods pilot study/td>
Evans (2019)	The role of engagement in mindfulness-based cognitive therapy for the prevention of depressive relapse/recurrence in perinatal women/td>
Falcone (2005)	[Multiprofessional care and mental health in pregnant women]/td>
Fancourt (2019)	Creative interventions for symptoms of postnatal depression: A process evaluation of

	implementation/td>
Feeley (2008)	Assessing the feasibility and acceptability of an intervention to reduce anxiety and enhance sensitivity among mothers of very low birth-weight infants/td>
Feinberg (2012)	Adaptation of problem-solving treatment for prevention of depression among low-income, culturally diverse mothers/td>
Felder (2016)	Targeting barriers to care for pregnant women at risk for depression: Examining the role of stigma and the feasibility of a web-based depression prevention program/td>
Felder (2017)	Depressive symptoms and gestational length among pregnant adolescents: Cluster randomized control trial of CenteringPregnancy plus group prenatal care/td>
Felder (2017)	An Open Trial of Web-Based Mindfulness-Based Cognitive Therapy for Perinatal Women at Risk for Depressive Relapse/td>
Felder (2018)	Poor Sleep Quality, Psychological Distress, and the Buffering Effect of Mindfulness Training During Pregnancy/td>
Figueira (2009)	Edinburgh Postnatal Depression Scale for screening in the public health system/td>
Figueroa-Leigh (2015)	Screening for postpartum depression in a private health care network in Chile/td>
Filoché (2016)	Access to Secondary Mental Health Services in a Cohort of New Zealand Mothers/td>
Fisher (1997)	Adverse psychological impact of operative obstetric interventions: a prospective longitudinal study/td>
Fisher (2010)	Innovative psycho-educational program to prevent common postpartum mental disorders in primiparous women: a before and after controlled study/td>
Fisher (2012)	Uneven implementation of the National Perinatal Depression Initiative: findings from a survey of Australian women's hospitals/td>
Fisher (2016)	Gender-informed, psychoeducational programme for couples to prevent postnatal common mental disorders among primiparous women: cluster randomised controlled trial/td>
Flanagan (2016)	Perinatal Obstetric Office Depression Screening and Treatment: Implementation in a Health Care System/td>
Flynn (2006)	The impact of a brief obstetrics clinic-based intervention on treatment use for perinatal depression/td>
Flynn (2011)	Comparative performance of the Edinburgh Postnatal Depression Scale and the Patient Health Questionnaire-9 in pregnant and postpartum women seeking psychiatric services/td>
Flynn (2016)	The Effects of Guided Imagery on Stress in Pregnant Adolescents/td>
Fonseca (2016)	Women's use of online resources and acceptance of e-mental health tools during the perinatal period/td>
Fonseca (2018)	Be a Mom: Formative Evaluation of a Web-Based Psychological Intervention to Prevent Postpartum Depression/td>
Forsell (2017)	Internet delivered cognitive behavior therapy for antenatal depression: A randomised controlled trial/td>
Forsyth (2017)	Exercise as an adjunct treatment for postpartum depression for women living in an inner city-A pilot study/td>
Fossey (1997)	Postpartum blues: a clinical syndrome and predictor of postnatal depression?/td>
Freeman (2005)	Postpartum depression assessments at well-baby visits: screening feasibility, prevalence, and risk factors/td>
Freeman (2008)	Omega-3 fatty acids and supportive psychotherapy for perinatal depression: a randomized placebo-controlled study/td>
Fuggle (2000)	Estimating resources for setting up an HV service for postnatal depression/td>
Fuhr (2019)	Delivering the Thinking Healthy Programme for perinatal depression through peers: an individually randomised controlled trial in India/td>
Gamble (2004)	Counselling processes to address psychological distress following childbirth: perceptions of midwives/td>
Gamble (2004)	Counselling processes to address psychological distress following childbirth: perceptions of women/td>
Gao (2012)	Effects of an interpersonal-psychotherapy-oriented childbirth education programme for Chinese first-time childbearing women at 3-month follow up: randomised controlled trial/td>
Gao (2012)	Interpersonal psychotherapy-oriented program for Chinese pregnant women: delivery, content, and personal impact/td>
Gao (2015)	Effects of an interpersonal-psychotherapy-oriented postnatal programme for Chinese first-time mothers: a randomized controlled trial/td>

Garcia (2010)	Repetitive transcranial magnetic stimulation treats postpartum depression/td>
García (2018)	Effects of prenatal music stimulation on state/trait anxiety in full-term pregnancy and its influence on childbirth: a randomized controlled trial/td>
Garnier (2008)	Perinatal depression, prevention strategies, personality, and the importance of therapy acceptance/td>
George (2019)	Effectiveness of a group intervention led by lay health workers in reducing the incidence of postpartum depression in South India/td>
Georgiopoulos (1999)	Population-based screening for postpartum depression/td>
Georgiopoulos (2001)	Routine screening for postpartum depression/td>
Gerrard (1993)	A trainer's perspective of an innovative programme teaching health visitors about the detection, treatment and prevention of postnatal depression/td>
Gibbon (2004)	Developments in perinatal mental health assessments/td>
Gillis (2019)	Implementation of a Perinatal Depression Care Bundle in a Nurse-Managed Midwifery Practice/td>
Ginsburg (2012)	Postpartum Depression Prevention for Reservation-Based American Indians: Results from a Pilot Randomized Controlled Trial/td>
Gjerdingen (2009)	Postpartum depression screening at well-child visits: validity of a 2-question screen and the PHQ-9/td>
Gjerdingen (2009)	Stepped care treatment of postpartum depression: impact on treatment, health, and work outcomes/td>
Gjerdingen (2011)	Problems with a diagnostic depression interview in a postpartum depression trial/td>
Gjerdingen (2013)	Postpartum doula and peer telephone support for postpartum depression: a pilot randomized controlled trial/td>
Glavin (2010)	Redesigned community postpartum care to prevent and treat postpartum depression in women--a one-year follow-up study/td>
Goodman (2014)	CALM Pregnancy: results of a pilot study of mindfulness-based cognitive therapy for perinatal anxiety/td>
Goudarzi (2016)	The effect of educational program on stress, anxiety and depression of the mothers of neonates having colostomy/td>
Gray (2014)	Partnering for mental health promotion: Implementing evidence based mental health services within a maternal and child home health visiting program/td>
Green (2015)	Cognitive-behavioral group treatment for perinatal anxiety: a pilot study/td>
Grote (2004)	Culturally Relevant Psychotherapy for Perinatal Depression in Low-Income Ob/Gyn Patients/td>
Grote (2004)	Feasibility of Providing Culturally Relevant, Brief Interpersonal Psychotherapy for Antenatal Depression in an Obstetrics Clinic: A Pilot Study/td>
Grote (2009)	A randomized controlled trial of culturally relevant, brief interpersonal psychotherapy for perinatal depression/td>
Grote (2014)	Culturally relevant treatment services for perinatal depression in socio-economically disadvantaged women: the design of the MOMCare study/td>
Gunning (2010)	Assessing maternal anxiety in pregnancy with the State-Trait Anxiety Inventory (STAI): Issues of validity, location and participation/td>
Haakstad (2016)	What is the effect of regular group exercise on maternal psychological outcomes and common pregnancy complaints? An assessor blinded RCT/td>
Haga (2013)	Mamma mia: a feasibility study of a web-based intervention to reduce the risk of postpartum depression and enhance subjective well-being/td>
Haga (2019)	Mamma Mia - A randomized controlled trial of an internet-based intervention for perinatal depression/td>
Hajmohamadi (2018)	The "Cooperative-Supportive" Intervention for Improving Mental Health Status among Pregnant Women/td>
Hall (2014)	An analysis of Time 4U, a therapeutic group for women with postnatal depression/td>
Hanbury (2013)	Immediate versus sustained effects: interrupted time series analysis of a tailored intervention/td>
Hancock (2018)	A randomised trial of early palliative care for maternal stress in infants prenatally diagnosed with single-ventricle heart disease/td>
Hanna (2004)	The clinical application of three screening tools for recognizing post-partum depression/td>
Hantsoo (2014)	A randomized, placebo-controlled, double-blind trial of sertraline for postpartum depression/td>
Hantsoo (2017)	Pregnancy and the Acceptability of Computer-Based Versus Traditional Mental Health Treatments/td>

Hantsoo (2018)	A mobile application for monitoring and management of depressed mood in a vulnerable pregnant population/td>
Harrington (2007)	Healthy Start screens for depression among urban pregnant, postpartum and interconceptional women/td>
Harrison (2008)	Systematic prenatal screening for psychosocial risks/td>
Harrison-Hohner (2001)	Prenatal calcium supplementation and postpartum depression: An ancillary study to a randomized trial of calcium for prevention of preeclampsia/td>
Harvey (2012)	Evaluating the clinical efficacy of a primary care-focused, nurse-led, consultation liaison model for perinatal mental health/td>
Harvey (2018)	Evaluating a nurse-led community model of service for perinatal mental health/td>
Hauck (2008)	Healthy babies for mothers with serious mental illness: a case management framework for mental health clinicians/td>
Hayes (2001)	Perinatal depression: a randomized controlled trial of an antenatal education intervention for primiparas/td>
Heberlein (2015)	The comparative effectiveness of group prenatal care on women's psychosocial health/td>
Heberlein (2016)	The comparative effects of group prenatal care on psychosocial outcomes/td>
Hegarty (2007)	Women's views and outcomes of an educational intervention designed to enhance psychosocial support for women during pregnancy/td>
Hepp (2018)	Effects of music intervention during caesarean delivery on anxiety and stress of the mother a controlled, randomised study/td>
Herizchi (2017)	The impact of Kangaroo-Mother care on postpartum depression in mothers of premature infants/td>
Highet (2004)	A comparative evaluation of community treatments for post-partum depression: implications for treatment and management practices/td>
Highet (2018)	Perinatal mental health and psychosocial risk screening in a community maternal and child health setting: Evaluation of a digital platform/td>
Hillier (1989)	The impact of antenatal classes on knowledge, anxiety and confidence in primiparous women/td>
Hiscock (2007)	Improving infant sleep and maternal mental health: a cluster randomised trial/td>
Ho (2009)	Effectiveness of a discharge education program in reducing the severity of postpartum depression: a randomized controlled evaluation study/td>
Hoffman (2019)	A screening protocol for perinatal mood and anxiety disorders in expectant mothers/td>
Holt (2017)	Improving help-seeking for postnatal depression and anxiety: a cluster randomised controlled trial of motivational interviewing/td>
Honey (2002)	A brief psycho-educational group intervention for postnatal depression/td>
Horowitz (2011)	A community-based screening initiative to identify mothers at risk for postpartum depression/td>
Horsch (2016)	Improving Maternal Mental Health Following Preterm Birth Using an Expressive Writing Intervention: A Randomized Controlled Trial/td>
Hosseinian (2016)	Effectiveness of mindfulness-based cognitive therapy and metacognition therapy on reduction of symptoms of depression, core self-evaluation and emotion regulation difficulties in pregnant women with depression/td>
Howard (2006)	A psychiatric mother-baby day hospital for pregnant and postpartum women/td>
Huang (2007)	Depressive symptomatology and mental health help-seeking patterns of U.S.- and foreign-born mothers/td>
Hubley (2014)	Training obstetrics nurses and behavioral health care providers in behavioral activation for antenatal depression/td>
Hung (2014)	Community-based prenatal screening for postpartum depression in a South African township/td>
Ickovics (2011)	Effects of group prenatal care on psychosocial risk in pregnancy: results from a randomised controlled trial/td>
Iheanacho (2015)	Integrating mental health screening into routine community maternal and child health activity: experience from Prevention of Mother-to-child HIV transmission (PMTCT) trial in Nigeria/td>
Imran (2010)	Screening of antenatal depression in Pakistan: Risk factors and effects on obstetric and neonatal outcomes/td>
Imura (2006)	The psychological effects of aromatherapy-massage in healthy postpartum mothers/td>
Ingadottir (2006)	Evaluation of a web-based course for community nurses on postpartum emotional distress/td>
Ingram (2007)	Predictors of postnatal depression: Using an antenatal needs assessment discussion tool/td>
Ip (2015)	An evaluation of a nurse-led Comprehensive Child Development Service in Hong Kong/td>

Jackson (2018)	Maternal depression in Barbados: Exploring how black women experience, understand, manage and cope with self-reported 'baby blues' and postpartum depression/td>
Jahdi (2016)	Evaluating the effectiveness of empowerment-based education on empowerment and anxiety in nulliparous women/td>
Jallo (2013)	Effects of a guided imagery intervention on stress in hospitalized pregnant women: a pilot study/td>
Jardri (2010)	Impact of midwives' training on postnatal depression screening in the first week post delivery: a quality improvement report/td>
Jesse (2010)	A pilot study to reduce risk for antepartum depression among women in a public health prenatal clinic/td>
Jesse (2015)	Performance of a Culturally Tailored Cognitive-Behavioral Intervention Integrated in a Public Health Setting to Reduce Risk of Antepartum Depression: A Randomized Controlled Trial/td>
Jevitt (2005)	Screening for perinatal depression with limited psychiatric resources/td>
Jiang (2014)	Psychological intervention for postpartum depression/td>
Jiao (2019)	Web-based versus home-based postnatal psychoeducational interventions for first-time mothers: A randomised controlled trial/td>
John (2018)	Activity based group therapy reduces maternal anxiety in the Neonatal Intensive Care Unit - a prospective cohort study/td>
Johnson (2017)	Parents' adverse childhood experiences and mental health screening using home visiting programs: A pilot study/td>
Johnson-Agbakwu (2014)	Mental health screening among newly arrived refugees seeking routine obstetric and gynecologic care/td>
Jomeen (2008)	The impact of choice of maternity care on psychological health outcomes for women during pregnancy and the postnatal period/td>
Jones (2012)	Australian midwives' awareness and management of antenatal and postpartum depression/td>
Jones (2015)	A Home-Start peer support scheme for women with low mood following childbirth/td>
Kaaya (2013)	Randomized controlled trial evaluating the effect of an interactive group counseling intervention for HIV-positive women on prenatal depression and disclosure of HIV status/td>
Kadivar (2015)	The effect of narrative writing on maternal stress in neonatal intensive care settings/td>
Kalina (2016)	Standardized postpartum depression screening and treatment/td>
Kalra (2018)	An evaluation of routine antenatal depression screening and psychosocial assessment in a regional private maternity setting in Australia/td>
Kamibepu (2009)	Training health professionals to detect and support mothers at risk of postpartum depression or infant abuse in the community: a cross-sectional and a before and after study/td>
Karamoozian (2015)	Impact of prenatal cognitive-behavioral stress management intervention on maternal anxiety and depression and newborns' Apgar scores/td>
Karatas (2009)	Antenatal psychosocial assessment: how accurate are we in determining 'low-risk' status? A pilot study/td>
Kelman (2018)	A proof-of-concept pilot randomized comparative trial of brief Internet-based compassionate mind training and cognitive-behavioral therapy for perinatal and intending to become pregnant women/td>
Kelman (2018)	A proof-of-concept pilot randomized comparative trial of brief Internet-based compassionate mind training and cognitive-behavioral therapy for perinatal and intending to become pregnant women/td>
Khamseh (2019)	Effectiveness of applying problem-solving training on depression in Iranian pregnant women: Randomized clinical trial/td>
Khan (2017)	Evaluating feasibility and acceptability of a local psycho-educational intervention for pregnant women with common mental problems affected by armed conflict in Swat, Pakistan: A parallel randomized controlled feasibility trial/td>
Khan (2018)	Assessing the impact of mindfulness and life stress on maternal well-being/td>
Khan (2019)	Culturally-adapted cognitive behavioural therapy based intervention for maternal depression: a mixed-methods feasibility study/td>
Khodakarami (2017)	Impact of a Counseling Program on Depression, Anxiety, Stress, and Spiritual Intelligence in Pregnant Women/td>
Khojasteh (2016)	Comparison of the effects of massage therapy and guided imagery on anxiety of nulliparous women during pregnancy/td>
Kildea (2018)	Continuity of midwifery carer moderates the effects of prenatal maternal stress on postnatal maternal wellbeing: the Queensland flood study/td>
Kim (2007)	Automated depression screening in disadvantaged pregnant women in an urban obstetric clinic/td>

Kim (2010)	Barriers to mental health treatment among obstetric patients at risk for depression/td>
Kim (2011)	A survey of patient acceptability of repetitive transcranial magnetic stimulation (TMS) during pregnancy/td>
Kim (2012)	Screening for postpartum depression among low-income mothers using an interactive voice response system/td>
Kim (2014)	Computer-assisted cognitive behavioral therapy for pregnant women with major depressive disorder/td>
Kimmel (2017)	Maternal Experiences in a Parenting Group Delivered in an Urban General Pediatric Clinic/td>
Kingston (2015)	Disclosure during prenatal mental health screening/td>
Kingston (2015)	Barriers and facilitators of mental health screening in pregnancy/td>
Kingston (2017)	Pregnant Women's Perceptions of the Risks and Benefits of Disclosure During Web-Based Mental Health E-Screening Versus Paper-Based Screening: Randomized Controlled Trial/td>
Kingston (2017)	Pregnant Women's Views on the Feasibility and Acceptability of Web-Based Mental Health E-Screening Versus Paper-Based Screening: A Randomized Controlled Trial/td>
Kleiber (2017)	Treating Depression Among Adolescent Perinatal Women With a Dialectical Behavior Therapy-Informed Skills Group/td>
Ko (2008)	Effects of postpartum exercise program on fatigue and depression during 'doing-the-month' period/td>
Ko (2013)	Community-based postpartum exercise program/td>
Kohlhoff (2016)	Antenatal psychosocial assessment and depression screening in a private hospital/td>
Kopelman (2008)	Barriers to care for antenatal depression/td>
Kozinszky (2012)	Can a brief antepartum preventive group intervention help reduce postpartum depressive symptomatology?/td>
Kralj (2016)	Creating pathways of care: A multidisciplinary approach to increasing screening and treatment of postpartum depression within a rural health clinic/td>
Krusche (2018)	Mindfulness for pregnancy: A randomised controlled study of online mindfulness during pregnancy/td>
Kurzweil (2012)	Psychodynamic therapy for depression in women with infants and young children/td>
La Porte (2019)	Feasibility of perinatal mood screening and text messaging on patients' personal smartphones/td>
Laios (2010)	What are we screening for? Development of a psychiatric referral tool for use with adolescent pregnant women/td>
Langer (1998)	Effects of psychosocial support during labour and childbirth on breastfeeding, medical interventions, and mothers' wellbeing in a Mexican public hospital: a randomised clinical trial/td>
Lara (2010)	Outcome results of a psycho-educational intervention in pregnancy to prevent PPD: a randomized control trial/td>
Lara (2010)	Retention rates and potential predictors in a longitudinal randomized control trial to prevent postpartum depression/td>
Lara (2014)	Acceptability and barriers to treatment for perinatal depression. An exploratory study in Mexican women/td>
LaRocco-Cockburn (2013)	Improving depression treatment for women: integrating a collaborative care depression intervention into OB-GYN care/td>
Lawson (2019)	Use of Text Messaging for Postpartum Depression Screening and Information Provision/td>
Le (2009)	Using the internet to screen for postpartum depression/td>
Le (2010)	Transforming an evidence-based intervention to prevent perinatal depression for low-income Latina immigrants/td>
Le (2013)	In their own voices: Latinas' experiences with a randomized controlled trial/td>
Le (2019)	Preventing perinatal depression in Spain: a pilot evaluation of Mamas y Bebés/td>
Leigh (2007)	Acceptability of antenatal screening for depression in routine antenatal care/td>
Leis (2011)	Perceptions of mental health services among low-income, perinatal African-American women/td>
Lenze (2015)	A pilot, exploratory report on dyadic interpersonal psychotherapy for perinatal depression/td>
Lenze (2017)	Brief Interpersonal Psychotherapy for depression during pregnancy in a low-income population: A randomized controlled trial/td>
Letourneau (2007)	Canadian mothers' perceived support needs during postpartum depression/td>
Leung (2002)	Postpartum depression: Perceived social support and stress among Hong Kong Chinese women/td>

Leung (2013)	Culturally sensitive, preventive antenatal group cognitive-behavioural therapy for Chinese women with depression/td>
Lewis (2019)	Developing a Hospital-Based Postpartum Depression Education Intervention for Perinatal Nurses/td>
Lia-Hoagberg (1990)	Barriers and motivators to prenatal care among low-income women/td>
Liberto (2011)	Attitudes Toward Help Seeking Among Depressed and Non-depressed Postpartum Women/td>
Liu (2010)	Effects of music therapy on labour pain and anxiety in Taiwanese first-time mothers/td>
Liu (2014)	Effect of postpartum practices of doing the month on Chinese women's physical and psychological health/td>
Logsdon (2004)	Prediction of postpartum social support and symptoms of depression in pregnant adolescents: a pilot study/td>
Logsdon (2010)	Adapting and testing telephone-based depression care management intervention for adolescent mothers/td>
Logsdon (2013)	Testing of a prototype Web based intervention for adolescent mothers on postpartum depression/td>
Logsdon (2015)	Acceptability and initial efficacy of education for teen mothers/td>
Logsdon (2018)	Screening for Postpartum Depression by Hospital-Based Perinatal Nurses/td>
Lopez (2015)	Financial incentives for smoking cessation among depression-prone pregnant and newly postpartum women: effects on smoking abstinence and depression ratings/td>
Loudon (2016)	Using clinical decision support as a means of implementing a universal postpartum depression screening program/td>
Loughnan (2019)	A randomized controlled trial of 'MUMentum Pregnancy': Internet-delivered cognitive behavioral therapy program for antenatal anxiety and depression/td>
Lovero (2019)	Mixed-methods evaluation of mental healthcare integration into tuberculosis and maternal-child healthcare services of four South African districts/td>
Ludwick (2017)	Postpartum Depression: Implementing an Evidence-Based Social Support Network in North Carolina/td>
MacArthur (2003)	Redesigning postnatal care: a randomised controlled trial of protocol-based midwifery-led care focused on individual women's physical and psychological health needs/td>
Madden (2018)	Using action research to develop midwives' skills to support women with perinatal mental health needs/td>
Madhavanprabhakaran (2015)	Prevalence of pregnancy anxiety and associated factors/td>
Magalhaes (2007)	Impact of defense style on brief psychotherapy of postpartum depression/td>
Mahato (2018)	Qualitative evaluation of mental health training of auxiliary nurse midwives in rural Nepal/td>
Mak (2019)	Physical activity during early pregnancy and antenatal depression: A prospective cohort study/td>
Malis (2017)	Effects of an antenatal mindfulness-based childbirth and parenting programme on the postpartum experiences of mothers: a qualitative interview study/td>
Maloni (2013)	Web recruitment and internet use and preferences reported by women with postpartum depression after pregnancy complications/td>
Malqvist (2016)	Screening for Antepartum Depression Through Community Health Outreach in Swaziland/td>
Mane (2013)	A study on mental health status among pregnant women and the social factors influencing/td>
Mann (2015)	The acceptability of case-finding questions to identify perinatal depression/td>
Mao (2012)	Effectiveness of Antenatal Emotional Self-Management Training Program in Prevention of Postnatal Depression in Chinese Women/td>
Marcus (2003)	Depressive symptoms among pregnant women screened in obstetrics settings/td>
Marrs (2013)	Evaluation of the impact of a CBT-based group on maternal postnatal mental health difficulties/td>
Martin (2009)	Psychopharmacologic management of opioid-dependent women during pregnancy/td>
Martinez (2016)	Barriers to access to treatment for mothers with postpartum depression in primary health care centers: a predictive model/td>
Mathew (2017)	Perceived postpartum stress and coping strategies among postnatal mothers at aims, Kochi/td>
Matsumoto (2019)	Cognitive behavioral therapy for postpartum panic disorder: a case series/td>
Matthey (2018)	A comparison of two measures to screen for emotional health difficulties during pregnancy/td>
Mazzoni (2018)	The effect of group prenatal care for women with diabetes on social support and depressive symptoms: a pilot randomized trial/td>
McCabe-Beane (2018)	Anxiety Screening During Assessment of Emotional Distress in Mothers of Hospitalized

	Newborns/td>
McCarter (2018)	Measuring outcomes of digital technology-assisted nursing postpartum: A randomized controlled trial/td>
McCarter-Spaulding (2016)	Effectiveness of Discharge Education on Postpartum Depression/td>
McCarthy (2008)	Acceptance and experience of treatment for postnatal depression in a community mental health setting/td>
McFarlane (2017)	Outcomes of a Randomized Trial of a Cognitive Behavioral Enhancement to Address Maternal Distress in Home Visited Mothers/td>
McGregor (2014)	The effect of physician-based cognitive behavioural therapy among pregnant women with depressive symptomatology: a pilot quasi-experimental trial/td>
McKean (2018)	Postpartum Depression: When Should Health Care Providers Identify Those at Risk?/td>
McKee (2006)	Results of an Intervention to Reduce Perinatal Depression Among Low-Income Minority Women in Community Primary Care/td>
McKellar (2017)	Capture my mood: a feasibility study to develop a visual scale for women to self-monitor their mental wellbeing following birth/td>
McLeish (2017)	Mothers' accounts of the impact on emotional wellbeing of organised peer support in pregnancy and early parenthood: a qualitative study/td>
McLeish (2019)	"Being the best person that they can be and the best mum": a qualitative study of community volunteer doula support for disadvantaged mothers before and after birth in England/td>
Mehta (2012)	The 5-HTTLPR polymorphism modulates the influence on environmental stressors on peripartum depression symptoms/td>
Meijssen (2011)	Maternal psychological distress in the first two years after very preterm birth and early intervention/td>
Meine (2018)	Pregnancy unshackled: Increasing equity through implementation of perinatal depression screening, shared decision making, and treatment for incarcerated women/td>
Melnyk (2008)	Maternal anxiety and depression after a premature infant's discharge from the neonatal intensive care unit: explanatory effects of the creating opportunities for parent empowerment program/td>
Melnyk (2009)	Reducing hospital expenditures with the COPE (Creating Opportunities for Parent Empowerment) program for parents and premature infants: an analysis of direct healthcare neonatal intensive care unit costs and savings/td>
Mendelson (2013)	Impact of a preventive intervention for perinatal depression on mood regulation, social support, and coping/td>
Mendelson (2018)	A mindfulness intervention to reduce maternal distress in neonatal intensive care: a mixed methods pilot study/td>
Mgonja (2017)	Postpartum Depression Screening at Well-Child Appointments: A Quality Improvement Project/td>
Mianaei (2014)	The effect of Creating Opportunities for Parent Empowerment program on maternal stress, anxiety, and participation in NICU wards in Iran/td>
Miklowitz (2015)	Mindfulness-Based Cognitive Therapy for Perinatal Women with Depression or Bipolar Spectrum Disorder/td>
Milgrom (2011)	Towards parenthood: an antenatal intervention to reduce depression, anxiety and parenting difficulties/td>
Milgrom (2015)	Feasibility study and pilot randomised trial of an antenatal depression treatment with infant follow-up/td>
Milgrom (2016)	Internet Cognitive Behavioral Therapy for Women With Postnatal Depression: A Randomized Controlled Trial of MumMoodBooster/td>
Miller (2004)	Improving the identification and treatment of postpartum depression in a managed care organization/td>
Miller (2008)	Interpersonal psychotherapy with pregnant adolescents: two pilot studies/td>
Miller (2009)	Beyond screening: assessment of perinatal depression in a perinatal care setting/td>
Miller (2012)	Now what? Effects of on-site assessment on treatment entry after perinatal depression screening/td>
Misri (2000)	The impact of partner support in the treatment of postpartum depression/td>
Misri (2006)	The impact of treatment intervention on parenting stress in postpartum depressed mothers: a prospective study/td>
Misri (2013)	Factors impacting decisions to decline or adhere to antidepressant medication in perinatal women with mood and anxiety disorders/td>
Misund (2013)	Long-term risk of mental health problems in women experiencing preterm birth: a longitudinal study

	of 29 mothers/td>
Misund (2014)	Mental health in women experiencing preterm birth/td>
Moayedoddin (2013)	The impact of brief psychotherapy centred on parenthood on the anxio-depressive symptoms of mothers during the perinatal period/td>
Mojrian (2018)	The Effect of Supportive Counseling on the Symptoms of Acute Stress Disorder Following Emergency Cesarean Section/td>
Morrell (2011)	Training health visitors in cognitive behavioural and person-centred approaches for depression in postnatal women as part of a cluster randomised trial and economic evaluation in primary care: the PoNDER trial/td>
Morris (1987)	Group psychotherapy for prolonged postnatal depression/td>
Morton (2013)	My Time, My Space (an arts-based group for women with postnatal depression): a project report/td>
Mouradian (2013)	Art-based occupation group reduces parent anxiety in the neonatal intensive care unit: a mixed-methods study/td>
Mugweni (2019)	"I'll look after the kids while you go and have a shower": an evaluation of a service to address mild to moderate maternal perinatal mental health problems/td>
Mundorf (2018)	Reducing the Risk of Postpartum Depression in a Low-Income Community Through a Community Health Worker Intervention/td>
Munoz (2007)	Prevention of Postpartum Depression in Low-Income Women: Development of the Mamas y Bebes/Mothers and Babies Course/td>
Murphey (2017)	Psychological Distress in Healthy Low-Risk First-Time Mothers during the Postpartum Period: An Exploratory Study/td>
Muzik (2012)	Mindfulness yoga during pregnancy for psychiatrically at-risk women: preliminary results from a pilot feasibility study/td>
Muzik (2015)	Mom Power: preliminary outcomes of a group intervention to improve mental health and parenting among high-risk mothers/td>
Myors (2014)	'My special time': Australian women's experiences of accessing a specialist perinatal and infant mental health service/td>
Myors (2018)	'Modelling a Secure-Base' for Women with Complex Needs: Attachment-Based Interventions Used by Perinatal and Infant Mental Health Clinicians/td>
Nagle (2018)	Women's views and experiences of having their mental health needs considered in the perinatal period/td>
Nasiri (2015)	A comparative study of the effects of problem-solving skills training and relaxation on the score of self-esteem in women with postpartum depression/td>
Navaie-Waliser (2000)	Social support and psychological functioning among high-risk mothers: the impact of the Baby Love Maternal Outreach Worker Program/td>
Navidian (2017)	The effect of home-based supportive-educational counseling on primigravidas' postpartum stress/td>
Navodani (2019)	Common maternal health problems among Australian-born and migrant women: A prospective cohort study/td>
Naysmith (2015)	Development and outcomes of a therapeutic group for women with postnatal depression/td>
Ngai (2009)	The effects of a childbirth psychoeducation program on learned resourcefulness, maternal role competence and perinatal depression: a quasi-experiment/td>
Ngai (2011)	Psychosocial factors and maternal wellbeing: An exploratory path analysis/td>
Ngai (2012)	Learned resourcefulness, social support, and perinatal depression in chinese mothers/td>
Nguyen (2017)	Impact of preconceptional micronutrient supplementation on maternal mental health during pregnancy and postpartum: results from a randomized controlled trial in Vietnam/td>
Nieminen (2016)	Internet-provided cognitive behaviour therapy of posttraumatic stress symptoms following childbirth-a randomized controlled trial/td>
Ningrum (2018)	Stress management intervention for women with multiple roles: Case study of working women in post-partum period/td>
Nishi (2019)	Differences between Japan and Taiwan in the treatment of pregnant women with depressive symptoms by omega-3 fatty acids: An open-label pilot study/td>
Njoku (2017)	Translating Evidence of Skin-to-Skin and Rooming-in to Practice/td>
Noonan (2019)	Public health nurses' perinatal mental health training needs: A cross sectional survey/td>
Norouzi (2013)	The impact of kangaroo care and music on maternal state anxiety/td>
Novosad (2003)	The breathing bear: an intervention for crying babies and their mothers/td>
Nyatsanza (2016)	Filling the treatment gap: developing a task sharing counselling intervention for perinatal depression

	in Khayelitsha, South Africa/td>
O'Hara (2019)	A placebo controlled treatment trial of sertraline and interpersonal psychotherapy for postpartum depression/td>
Okronipa (2018)	Maternal supplementation with small-quantity lipid-based nutrient supplements during pregnancy and lactation does not reduce depressive symptoms at 6 months postpartum in Ghanaian women: a randomized controlled trial/td>
Olayiwola (2013)	Living Smart, Living Fit: a patient-centered program to improve perinatal outcomes in a community health center population/td>
Olhaberry (2015)	Video-feedback intervention in mother-baby dyads with depressive symptomatology and relationship difficulties/td>
O'Mahen (2008)	Preferences and perceived barriers to treatment for depression during the perinatal period/td>
O'Mahen (2013)	Internet-based behavioral activation--treatment for postnatal depression (Netmums): a randomized controlled trial/td>
O'Mahen (2013)	A pilot randomized controlled trial of cognitive behavioral therapy for perinatal depression adapted for women with low incomes/td>
O'Mahen (2014)	Netmums: a phase II randomized controlled trial of a guided Internet behavioural activation treatment for postpartum depression/td>
Onozawa (2001)	Infant massage improves mother-infant interaction for mothers with postnatal depression/td>
Orengo-Aguayo (2016)	Depression treatment delivered at the point-of-care: a qualitative assessment of the views of low-income US mothers/td>
Ormsby (2018)	Women's experiences of having depression during pregnancy and receiving acupuncture treatment- A qualitative study/td>
Ortiz (2014)	Antenatal psychosomatic programming to reduce postpartum depression risk and improve childbirth outcomes: a randomized controlled trial in Spain and France/td>
Oswalt (2013)	Reducing anxiety among children born preterm and their young mothers/td>
Pan (2019)	Assessing the effectiveness of mindfulness-based programs on mental health during pregnancy and early motherhood - a randomized control trial/td>
Panyayong (2013)	Postpartum depression among Thai women: a national survey/td>
Paris (2011)	Evaluating a home-based dyadic intervention: Changes in postpartum depression, maternal perceptions, and mother-infant interactions/td>
Parker (2015)	Predictors of post-natal depression are shaped distinctly by the measure of 'depression'/td>
Peindl (2004)	Identifying depression in the first postpartum year: guidelines for office-based screening and referral/td>
Perez-Blasco (2013)	Effects of a mindfulness-based intervention on psychological distress, well-being, and maternal self-efficacy in breast-feeding mothers: results of a pilot study/td>
Petch (2019)	Prediction of recurrent preterm delivery in asymptomatic women- an anxiety reducing measure?/td>
Phua (2017)	Positive maternal mental health during pregnancy associated with specific forms of adaptive development in early childhood: Evidence from a longitudinal study/td>
Polman (2007)	Effect of a single bout of exercise on the mood of pregnant women/td>
Posmontier (2016)	Telephone-Administered Interpersonal Psychotherapy by Nurse-Midwives for Postpartum Depression/td>
Posmontier (2019)	Incorporating Interpersonal Psychotherapy for Postpartum Depression Into Social Work Practice in Israel/td>
Powell (2018)	Mental health and well-being in parents of excessively crying infants: Prospective evaluation of a support package/td>
Premji (2019)	Examining postpartum depression screening effectiveness in well child clinics in Alberta, Canada: A study using the All Our Families cohort and administrative data/td>
Prevatt (2018)	Peer-support intervention for postpartum depression: Participant satisfaction and program effectiveness/td>
Prevatt (2018)	Facilitators and Barriers to Disclosure of Postpartum Mood Disorder Symptoms to a Healthcare Provider/td>
Preyde (2003)	Effectiveness of a parent "buddy" program for mothers of very preterm infants in a neonatal intensive care unit/td>
Price (2019)	A pilot feasibility study of mindfulness childbirth education for women with a history of sexual trauma/td>
Priest (2003)	Stress debriefing after childbirth: a randomised controlled trial/td>

Pritchett (2017)	Women's experiences of exercise as a treatment for their postnatal depression: A nested qualitative study/td>
Pugh (2016)	A Randomised Controlled Trial of Therapist-Assisted, Internet-Delivered Cognitive Behavior Therapy for Women with Maternal Depression/td>
Puryear (2019)	Outcomes of Implementing Routine Screening and Referrals for Perinatal Mood Disorders in an Integrated Multi-site Pediatric and Obstetric Setting/td>
Quispel (2014)	Withdrawal from mental and psychosocial care during pregnancy/td>
Rabiei (2014)	Fordyce happiness program and postpartum depression/td>
Rahman (2007)	Challenges and opportunities in developing a psychological intervention for perinatal depression in rural Pakistan--a multi-method study/td>
Ramos-Marcuse (2010)	Stability of maternal depressive symptoms among urban, low-income, African American adolescent mothers/td>
Ramphos (2019)	Responding to women's needs and preferences in an online program to prevent postpartum depression/td>
Reay (2003)	Implementation and effectiveness of interpersonal psychotherapy in a community mental health service/td>
Reay (2006)	Group interpersonal psychotherapy for postnatal depression: a pilot study/td>
Reid (2002)	A two-centred pragmatic randomised controlled trial of two interventions of postnatal support/td>
Reilly (2019)	Insights into implementation of routine depression screening and psychosocial assessment in a private hospital setting: A qualitative study/td>
Reinstein (2019)	The best perinatal depression screening: Is self-administered PHQ2 more feasible than a nurse-administered one?/td>
Reupert (2011)	Programmes for parents with a mental illness/td>
Ribeiro (2018)	Music therapy intervention in cardiac autonomic modulation, anxiety, and depression in mothers of preterms: randomized controlled trial/td>
Robichaud (2009)	The effects of an exercise intervention on the psychological well-being of postpartum women/td>
Robinson (1982)	Screening for depression and anxiety in the post-natal period: acceptance or rejection of a subsequent treatment offer/td>
Rock (2019)	A Stepped-Collaborative Perinatal Depression Model/td>
Rollans (2013)	'We just ask some questions...' the process of antenatal psychosocial assessment by midwives/td>
Rollans (2016)	Partner Involvement: Negotiating the Presence of Partners in Psychosocial Assessment as Conducted by Midwives and Child and Family Health Nurses/td>
Rossiter (2012)	Supporting depressed mothers at home: their views on an innovative relationship-based intervention/td>
Rotheram-Fuller (2018)	Maternal patterns of antenatal and postnatal depressed mood and the impact on child health at 3-years postpartum/td>
Rowan (2012)	Outcomes from implementing systematic antepartum depression screening in obstetrics/td>
Rowe (2010)	The contribution of Australian residential early parenting centres to comprehensive mental health care for mothers of infants: evidence from a prospective study/td>
Rowe (2017)	A complex postnatal mental health intervention: Australian translational formative evaluation/td>
Ruiz (2019)	Pilot Study of the Mastery Lifestyle Intervention/td>
Russomagno (2019)	Improving Postpartum Depression Screening and Referral in Pediatric Primary Care/td>
Saenz (2009)	Psychological stress of parents of preterm infants enrolled in an early discharge programme from the neonatal intensive care unit: a prospective randomised trial/td>
Salonen (2014)	Impact of an internet-based intervention on Finnish mothers' perceptions of parenting satisfaction, infant centrality and depressive symptoms during the postpartum year/td>
Sampson (2016)	A problem-solving therapy intervention for low-income, pregnant women at risk for postpartum depression/td>
San (2019)	Screening for perinatal depression and predictors of underscreening: findings of the Born in Queensland study/td>
Sawyer (2019)	The Effectiveness of an App-Based Nurse-Moderated Program for New Mothers With Depression and Parenting Problems (eMums Plus): Pragmatic Randomized Controlled Trial/td>
Schaar (2013)	A nurse-led initiative to improve obstetricians' screening for postpartum depression/td>
Schmied (2016)	Antenatal psychosocial risk status and Australian women's use of primary care and specialist mental health services in the year after birth: a prospective study/td>

Scholle (2003)	Addressing depression in obstetrics/gynecology practice/td>
Seeley (2019)	Mediation analyses of Internet-facilitated cognitive behavioral intervention for maternal depression/td>
Segre (2010)	Listening visits: an evaluation of the effectiveness and acceptability of a home-based depression treatment/td>
Segre (2012)	Depression screening of perinatal women by the Des Moines Healthy Start Project: program description and evaluation/td>
Segre (2013)	Perinatal depression screening in healthy start: an evaluation of the acceptability of technical assistance consultation/td>
Segre (2015)	Depression treatment for impoverished mothers by point-of-care providers: A randomized controlled trial/td>
Segre (2016)	Depression Management by NICU Nurses: Mothers' Views/td>
Selkirk (2006)	The longitudinal effects of midwife-led postnatal debriefing on the psychological health of mothers/td>
Sezen (2019)	Group art therapy for the management of fear of childbirth/td>
Shahnazi (2015)	The impact of education intervention on the Health Belief Model constructs regarding anxiety of nulliparous pregnant women/td>
Shahtaheri (2016)	The impact of stress reduction program based on mindfulness and conscious yoga on depression, perceived stress and quality of life in pregnant women/td>
Sharan (2006)	Early screening of postpartum depression using the Edinburgh Postnatal Depression Scale/td>
Sharma (2008)	Missed bipolarity and psychiatric comorbidity in women with postpartum depression/td>
Sharp (2010)	A pragmatic randomised controlled trial to compare antidepressants with a community-based psychosocial intervention for the treatment of women with postnatal depression: the RESPOND trial/td>
Shaw (2013)	Prevention of traumatic stress in mothers with preterm infants: a randomized controlled trial/td>
Sheeber (2017)	Mom-net: Evaluation of an internet-facilitated cognitive behavioral intervention for low-income depressed mothers/td>
Sheeder (2009)	Screening for postpartum depression at well-child visits: is once enough during the first 6 months of life?/td>
Shenai (2019)	Integrated Brief Intervention for PTSD and Substance Use in an Antepartum Unit/td>
Sherman (2018)	Diagnosis of Postpartum Depression and Timing and Types of Treatment Received Differ for Women with Private and Medicaid Coverage/td>
Shields (1997)	Impact of midwife-managed care in the postnatal period: An exploration of psychosocial outcomes/td>
Shivakumar (2014)	Screening to Treatment: Obstacles and Predictors in Perinatal Depression (STOP-PPD) in the Dallas Healthy Start program/td>
Shorey (2013)	The effectiveness of a Postnatal Psychoeducation Programme on outcomes of first-time mothers in Singapore/td>
Shorey (2015)	A randomized controlled trial of the effectiveness of a postnatal psychoeducation programme on self-efficacy, social support and postnatal depression among primiparas/td>
Shulman (2018)	Feasibility of a mindfulness-based cognitive therapy group intervention as an adjunctive treatment for postpartum depression and anxiety/td>
Sikander (2019)	Delivering the Thinking Healthy Programme for perinatal depression through volunteer peers: A cluster randomised controlled trial in Pakistan/td>
Silverstein (2011)	Problem-solving education to prevent depression among low-income mothers of preterm infants: a randomized controlled pilot trial/td>
Silverstein (2018)	Engaging mothers with depressive symptoms in care: Results of a randomized controlled trial in head start/td>
Simavli (2014)	Effect of music therapy during vaginal delivery on postpartum pain relief and mental health/td>
Simeone (2015)	Psychological impact of first-trimester prevention for preeclampsia on anxiety/td>
Siu (2011)	Pathway to care for postnatal depression after implementation of the Comprehensive Child Development Service in a regional hospital in Hong Kong/td>
Skoog (2019)	'Happy that someone cared'-Non-native-speaking immigrant mothers' experiences of participating in screening for postpartum depression in the Swedish child health services/td>
Small (2000)	Randomised controlled trial of midwife led debriefing to reduce maternal depression after operative childbirth/td>
Smith (2004)	Screening for and detection of depression, panic disorder, and PTSD in public-sector obstetric

	clinics/td>
Smith (2012)	Health Literacy and Depression in the Context of Home Visitation/td>
Smith (2018)	Evaluating a stepped care protocol for postpartum depression in a pediatric primary care clinic/td>
SmithBattle (2017)	Moms growing together: Piloting action methods and expressive arts in a therapeutic group for teen mothers/td>
Sorg (2019)	Improving Postpartum Depression Screening in Pediatric Primary Care: A Quality Improvement Project/td>
Sorsdahl (2015)	Feasibility and Preliminary Responses to a Screening and Brief Intervention Program for Maternal Mental Disorders Within the Context of Primary Care/td>
Steinberg (1999)	Characteristics and treatment of women with antenatal and postpartum depression/td>
Stevens (2018)	Examining the effectiveness of a coordinated perinatal mental health care model using an intersectional-feminist perspective/td>
Stewart (2017)	The impact of maternal diet fortification with lipid-based nutrient supplements on postpartum depression in rural Malawi: a randomised-controlled trial/td>
Stramrood (2012)	The patient observer: eye-movement desensitization and reprocessing for the treatment of posttraumatic stress following childbirth/td>
Surjaningrum (2018)	Personal attributes and competencies required by community health workers for a role in integrated mental health care for perinatal depression: voices of primary health care stakeholders from Surabaya, Indonesia/td>
Surkan (2012)	Impact of a health promotion intervention on maternal depressive symptoms at 15 months postpartum/td>
Swanson (2018)	An open-label pilot study of a home wearable light therapy device for postpartum depression/td>
Symon (2017)	The joy of parenting: infant sleep intervention to improve maternal emotional well-being and infant sleep/td>
Ta (2017)	Postpartum Traditions, Mental Health, and Help-Seeking Considerations Among Vietnamese American Women: a Mixed-Methods Pilot Study/td>
Tabeshpour (2017)	A double-blind, randomized, placebo-controlled trial of saffron stigma (<i>Crocus sativus</i> L.) in mothers suffering from mild-to-moderate postpartum depression/td>
Tabor (1987)	Psychological impact of amniocentesis on low-risk women/td>
Tachibana (2019)	Integrated mental health care in a multidisciplinary maternal and child health service in the community: the findings from the Suzaka trial/td>
Tandon (2005)	Formative evaluation of home visitors' role in addressing poor mental health, domestic violence, and substance abuse among low-income pregnant and parenting women/td>
Tandon (2011)	Preventing perinatal depression in low-income home visiting clients: a randomized controlled trial/td>
Teissedre (2003)	Postnatal depression: a study of the predictive effects of postnatal anxiety/td>
Teychenne (2018)	Feasibility and acceptability of a home-based physical activity program for postnatal women with depressive symptoms: A pilot study/td>
Thippeswamy (2018)	Help-seeking patterns in women with postpartum severe mental illness: a report from southern India/td>
Thitipitchayanant (2018)	Effectiveness of self-empowerment-affirmation-relaxation(Self-EAR) program for postpartum blues mothers: A randomize controlled trial/td>
Thoppil (2005)	Early intervention for perinatal depression/td>
Thorpe (1993)	A study of the use of the Edinburgh Postnatal Depression Scale with parent groups outside the postpartum period/td>
Timpano (2011)	Efficacy of a prevention program for postpartum obsessive-compulsive symptoms/td>
Toler (2018)	Screening for postpartum anxiety: A quality improvement project to promote the screening of women suffering in silence/td>
Tragea (2014)	A randomized controlled trial of the effects of a stress management programme during pregnancy/td>
Traube (2017)	Development of a Quality Improvement Initiative to Screen for Postpartum Depression/td>
Tripathy (2010)	Effect of a participatory intervention with women's groups on birth outcomes and maternal depression in Jharkhand and Orissa, India: a cluster-randomised controlled trial/td>
Tsai (2014)	Antenatal depression case finding by community health workers in South Africa: feasibility of a mobile phone application/td>
Tubay (2019)	The Effects of Group Prenatal Care on Infant Birthweight and Maternal Well-Being: A Randomized Controlled Trial/td>
Turkstra (2017)	An economic evaluation alongside a randomised controlled trial on psycho-education counselling

	intervention offered by midwives to address women's fear of childbirth in Australia/td>
Uebelacker (2016)	A pilot randomized controlled trial comparing prenatal yoga to perinatal health education for antenatal depression/td>
Ugarriza (2004)	Group therapy and its barriers for women suffering from postpartum depression/td>
Ugarriza (2006)	Telecare for women with postpartum depression/td>
Upshur (2016)	Using Prenatal Advocates to Implement a Psychosocial Education Intervention for Posttraumatic Stress Disorder during Pregnancy: Feasibility, Care Engagement, and Predelivery Behavioral Outcomes/td>
van der Ham (2013)	A collaborative approach to perinatal and infant mental health service delivery in Australia/td>
van der Zee-van (2017)	Post-Up Study: Postpartum Depression Screening in Well-Child Care and Maternal Outcomes/td>
Van Lieshout (2019)	Public health nurse delivered group cognitive behavioral therapy (CBT) for postpartum depression: A pilot study/td>
Van Ravesteyn (2018)	Group-based multicomponent treatment to reduce depressive symptoms in women with co-morbid psychiatric and psychosocial problems during pregnancy: A randomized controlled trial/td>
Vargas-Terrones (2019)	Physical exercise programme during pregnancy decreases perinatal depression risk: a randomised controlled trial/td>
Venkatesh (2016)	Implementation of universal screening for depression during pregnancy: feasibility and impact on obstetric care/td>
Verdeli (2016)	Community Norms and Human Rights: Supervising Haitian Colleagues on Interpersonal Psychotherapy (IPT) With a Depressed and Abused Pregnant Woman/td>
Victoria (1994)	The Latin American Multicenter Trial on psychosocial support during pregnancy: methodology and baseline comparability. Latin American Network for Perinatal and Reproductive Research (LANPER)/td>
Vieten (2008)	Effects of a mindfulness-based intervention during pregnancy on prenatal stress and mood: results of a pilot study/td>
Vieten (2018)	The mindful moms training: development of a mindfulness-based intervention to reduce stress and overeating during pregnancy/td>
Vigod (2019)	Transcranial direct current stimulation (tDCS) for depression in pregnancy: A pilot randomized controlled trial/td>
Vigod (2019)	A patient decision aid for antidepressant use in pregnancy: Pilot randomized controlled trial/td>
Viveiros (2018)	Barriers and facilitators of accessing perinatal mental health services: The perspectives of women receiving continuity of care midwifery/td>
Wahlbeck (2018)	Gaining hope and self-confidence-An interview study of women's experience of treatment by art therapy for severe fear of childbirth/td>
Wajid (2019)	Adversity in childhood and depression in pregnancy/td>
Walton (2014)	Decisional conflict among women considering antidepressant medication use in pregnancy/td>
Warriner (2018)	An evaluation of mindfulness-based childbirth and parenting courses for pregnant women and prospective fathers/partners within the UK NHS (MBCP-4-NHS)/td>
Webster (2003)	IDentify, Educate and Alert (IDEA) trial: an intervention to reduce postnatal depression/td>
Weinreb (2014)	What happens to mental health treatment during pregnancy? Women's experience with prescribing providers/td>
Weinreb (2018)	Postpartum outcomes of a pilot prenatal care-based psychosocial intervention for PTSD during pregnancy/td>
Welch (2016)	Depression and anxiety symptoms of mothers of preterm infants are decreased at 4 months corrected age with Family Nurture Intervention in the NICU/td>
Werner (2016)	PREPP: postpartum depression prevention through the mother-infant dyad/td>
Wheatley (1999)	'Just because I like it doesn't mean it has to work': Personal experiences of an antenatal psychosocial intervention designed to prevent postnatal depression/td>
White (2015)	Antenatal parenting support for vulnerable women/td>
Wickberg (2005)	Using the EPDS in routine antenatal care in Sweden: A naturalistic study/td>
Williams (2008)	Missouri Ozark women's experiences of living with postpartum depression symptoms/td>
Woolhouse (2014)	Antenatal mindfulness intervention to reduce depression, anxiety and stress: a pilot randomised controlled trial of the MindBabyBody program in an Australian tertiary maternity hospital/td>
Wright (2017)	Computerised cognitive-behavioural therapy for depression in adolescents: feasibility results and 4-month outcomes of a UK randomised controlled trial/td>

Wroblewski (2004)	Implementing a comprehensive postpartum depression support program/td>
Wu (2019)	Impact of a peer virtual community on pregnant women's well-being: A repeated-measure and quasi-experimental study/td>
Yaman (2015)	Posttraumatic stress and experiences of parents with a newborn in the neonatal intensive care unit/td>
Yawn (2009)	Concordance of Edinburgh Postnatal Depression Scale (EPDS) and Patient Health Questionnaire (PHQ-9) to assess increased risk of depression among postpartum women/td>
Yawn (2010)	Practice benefit from participating in a practice-based research network study of postpartum depression: a national research network (NRN) report/td>
Yawn (2012)	TRIPPD: a practice-based network effectiveness study of postpartum depression screening and management/td>
Yazdanimehr (2016)	The Effect of Mindfulness-integrated Cognitive Behavior Therapy on Depression and Anxiety among Pregnant Women: a Randomized Clinical Trial/td>
Ying (2017)	The occurrence of depression and nursing intervention for mothers of preterm infants/td>
Yonkers (2009)	Depression screening of perinatal women: an evaluation of the healthy start depression initiative/td>
Zayas (2004)	Adapting psychosocial intervention research to urban primary care environments: a case example/td>
Zhang (2015)	A mindfulness-based intervention for pregnant African-American women/td>
Zhang (2016)	The impact of social support on postpartum depression: The mediator role of self-efficacy/td>
Zhao (2019)	Effects of antenatal depression screening and intervention among Chinese high-risk pregnant women with medically defined complications: A randomized controlled trial/td>
Zlotnick (2011)	An interpersonally based intervention for low-income pregnant women with intimate partner violence: a pilot study/td>
Zlotnick (2018)	A randomized controlled trial of a computer-based brief intervention for victimized perinatal women seeking mental health treatment/td>
Zourladani (2011)	The effect of a low impact exercise training programme on the well-being of Greek postpartum women: A randomised controlled trial/td>
Not primary research	
Andriotti (2017)	ASSERT trial - How to assess the safety and efficacy of a high frequency rTMS in postpartum depression ? A multicenter, double blinded, randomized, placebo-controlled clinical trial/td>
Battle (2010)	Prenatal yoga and depression during pregnancy/td>
Battle (2015)	Physical Activity as an Intervention for Antenatal Depression: Rationale for Developing Tailored Exercise Programs for Pregnant Women with Depression/td>
Baumgartner (2015)	Mental health screenings for couples at churches in Nigeria: a strategy for enhancing community-based maternal mental health services in low-resource settings/td>
Beck (1995)	Screening methods for postpartum depression/td>
Boyce (2019)	Screening for perinatal depression: is it enough?/td>
Buist (2002)	To screen or not to screen--that is the question in perinatal depression/td>
Buist (2002)	Mental health in pregnancy: The sleeping giant/td>
Canady (2018)	Coordinated, community support drives effort to address postpartum depression/td>
Caramlau (2011)	Mums 4 Mums: structured telephone peer-support for women experiencing postnatal depression. Pilot and exploratory RCT of its clinical and cost effectiveness/td>
Dennis (2006)	Counselling and cognitive behavioural therapy reduce anxiety and depression in women with postnatal depression/td>
Felder (2019)	Implementing the USPSTF Recommendations on Prevention of Perinatal Depression-Opportunities and Challenges/td>
Fonte (2005)	Treating Postpartum Depression in Immigrant Muslim Women/td>
Freeman (2019)	Perinatal Depression: Recommendations for Prevention and the Challenges of Implementation/td>
Haas (2016)	Yoga for Prenatal Depression/td>
Haynes (2018)	Reaching women with perinatal mental illness at the booking-in appointment/td>
Holden (1991)	Postnatal depression: its nature, effects, and identification using the Edinburgh Postnatal Depression scale/td>
Hughes (2017)	Identification and Treatment of Adolescents With Perinatal Anxiety and Depression/td>
Hynie (2015)	Community resilience and community interventions for post-natal depression: Reflecting on maternal mental health in Rwanda/td>

Keefe (2016)	Addressing access barriers to services for mothers at risk for perinatal mood disorders: A social work perspective/td>
Kenyon (2012)	Evaluation of Lay Support in Pregnant women with Social risk (ELSIPS): a randomised controlled trial/td>
Lenz (2018)	Mindfulness-based Stress Reduction in Pregnancy: An App-Based Programme to Improve the Health of Mothers and Children (MINDFUL/PMI Study)/td>
Logsdon (2005)	Bioecological model for guiding social support research and interventions with pregnant adolescents/td>
Logsdon (2006)	Raising the awareness of primary care providers about postpartum depression/td>
Marcano (2017)	A bespoke mobile application for the longitudinal assessment of depression and mood during pregnancy: protocol of a feasibility study/td>
Milgrom (2011)	Does postnatal depression screening work? Throwing out the bathwater, keeping the baby/td>
Milgrom (2014)	Feasibility and efficacy of an internet treatment for postnatal depression utilising a behavioural activation approach/td>
Pedrina (2004)	Group Therapy with Mothers and Babies in Postpartum Crises: Preliminary Evaluation of A Pilot Project/td>
Posmontier (2010)	The role of midwives in facilitating recovery in postpartum psychosis/td>
Sharma (2018)	Should all women with postpartum depression be screened for bipolar disorder?/td>
Shorey (2015)	A randomized controlled trial of the effectiveness of a postnatal psychoeducation programme on outcomes of primiparas: study protocol/td>
Silverstein (2011)	Preventing depression of mothers of preterm infants/td>
Sit (2009)	Best practices: an emerging best practice model for perinatal depression care/td>
Smith (2010)	The design of a community lifestyle programme to improve the physical and psychological well-being of pregnant women with a BMI of 30 kg/m ² or more/td>
SmithBattle (2016)	Teen Mothers' Mental Health/td>
Springate (2006)	Mental health providers' self-reported expertise and treatment of perinatal depression/td>
Terrazas (2018)	Moving beyond depression screening: integrating perinatal depression treatment into OB/GYN practices/td>
Yawn (2012)	Postpartum Depression: Screening, Diagnosis, and Management Programs 2000 through 2010/td>
Zelkowitz (2009)	The Cues and Care trial: a randomized controlled trial of an intervention to reduce maternal anxiety and improve developmental outcomes in very low birthweight infants/td>