

Reference

- Bayley A, Milosevic M, Blend R, Logue J, Gospodarowicz M, Boxen I, *et al.* A prospective study of factors predicting clinically occult spinal cord compression in patients with metastatic prostate carcinoma. *Cancer* 2001;**92**:303–10¹⁰⁷
- Bernat JL, Greenberg ER, Barrett J. Suspected epidural compression of the spinal cord and cauda equina by metastatic carcinoma. Clinical diagnosis and survival. *Cancer* 1983;**51**:1953–7¹⁰⁸
- Chaichana KL, Pendleton C, Wolinsky JP, Gokaslan ZL, Sciubba DM. Vertebral compression fractures in patients presenting with metastatic epidural spinal cord compression. *Neurosurgery* 2009;**65**:267–74¹⁰⁹
- Fisher CG, DiPaola CP, Ryken TC, Bilsky MH, Shaffrey CI, Berven SH, *et al.* A novel classification system for spinal instability in neoplastic disease: an evidence-based approach and expert consensus from the Spine Oncology Study Group. *Spine* 2010;**35**:E1221–9¹¹⁰
- Goldman JM, Ash CM, Souhami RL, Geddes DM, Harper PG, Spiro SG, *et al.* Spinal cord compression in small cell lung cancer: a retrospective study of 610 patients. *Br J Cancer* 1989;**59**:591–3¹¹¹
- Harrison KM, Muss HB, Ball MR, McWhorter M, Case D. Spinal cord compression in breast cancer. *Cancer* 1985;**55**:2839–44¹¹²
- Helweg-Larsen S, Sorensen PS, Kreiner S. Prognostic factors in metastatic spinal cord compression: a prospective study using multivariate analysis of variables influencing survival and gait function in 153 patients. *Int J Radiat Oncol Biol Phys* 2000;**46**:1163–9¹¹³
- Helweg-Larsen S, Hansen SW, Sorensen PS. Second occurrence of symptomatic metastatic spinal cord compression and findings of multiple spinal epidural metastases. *Int J Radiat Oncol Biol Phys* 1995;**33**:595–8¹¹⁴
- Huddart RA, *et al.* Spinal cord compression in prostate cancer: treatment outcome and prognostic factors. *Radiother Oncol* 1997;**44**:229–36¹¹⁵
- Husband DJ, Grant KA, Romaniuk CS. MRI in the diagnosis and treatment of suspected malignant spinal cord compression. *Br J Radiol* 2001;**74**:15–23¹¹⁶
- Klekamp J, Samii H. Surgical results for spinal metastases. *Acta Neurochir* 1998;**140**:957–67¹¹⁷
- Kuban DA, el-Mahdi AM, Sigfred SV, Schellhammer PF, Babb TJ. Characteristics of spinal cord compression in adenocarcinoma of prostate. *Urology* 1986;**28**:364–9¹¹⁸
- Levack P, Graham J, Collie D, Grant R, Kidd J, Kunkler I, *et al.* Don't wait for a sensory level—listen to the symptoms: a prospective audit of the delays in diagnosis of malignant cord compression. *Clin Oncol (R Coll Radiol)* 2002;**14**:472–80¹¹⁹
- Loblack DA, Perry J, Chambers A, Laperriere NJ. Systematic review of the diagnosis and management of malignant extradural spinal cord compression: the cancer care Ontario practice guidelines initiative's neuro-oncology disease site group. *J Clin Oncol* 2005;**23**:2028–37⁶²
- Lu C, Stomper PC, Drislane FW, Wen PY, Block CC, Humphrey CC. Suspected spinal cord compression in breast cancer patients: a multidisciplinary risk assessment. *Breast Cancer Res Treat* 1998;**51**:121–31¹²⁰
- Lu C, Gonzalez RG, Jolesz FA, Wen PY, Talcott JA. Suspected spinal cord compression in cancer patients: a multidisciplinary risk assessment. *J Support Oncol* 2005;**3**:305–12¹²¹
- McCloskey EV, Spector TD, Eyres KS, Fern ED, O'Rourke N, Vasikaran SM, *et al.* The assessment of vertebral deformity – a method for use in population studies and clinical-trials. *Osteoporos Int* 1993;**3**:138–47¹²²
- Oka H, Kondoh T, Seichi A, Hozumi T, Nakamura K. Incidence and prognostic factors of Japanese breast cancer patients with bone metastasis. *J Orthop Sci* 2006;**11**:13–19¹²³
- Plunkett TA, Smith P, Rubens RD. Risk of complications from bone metastases in breast cancer: implications for management. *Eur J Cancer* 2000;**36**:476–82²⁴
- Rose PS, Laufer I, Boland PJ, Hanover A, Bilsky MH, Yamada J, *et al.* Risk of fracture after single fraction image-guided intensity-modulated radiation therapy to spinal metastases. *J Clin Oncol* 2009;**27**:5075–9⁸⁸
- Roth SE, Mousavi P, Finkelstein J, Chow E, Kreder H, Whyne CM. Metastatic burst fracture risk prediction using biomechanically based equations. *Clin Orthop Relat Res* 2004;**419**:83–90¹²⁴

Reference

- Sekine I, Nokihara H, Yamamoto N, Kunitoh H, Ohe Y, Tamura T. Risk factors for skeletal-related events in patients with non-small cell lung cancer treated by chemotherapy. *Lung Cancer* 2009;**65**:219–22¹²⁵
- Shah AN, Pietrobon R, Richardson WJ, Myers BS. Patterns of tumor spread and risk of fracture and epidural impingement in metastatic vertebrae. *J Spinal Disord Tech* 2003;**16**:83–9¹²⁶
- Snyder BD, Hipp JA, Nazarian A. *Non-invasive prediction of fracture risk due to benign and metastatic skeletal defects*. Warrendale, PA: Materials Research Society; 2005. pp. 39–51¹²⁷
- Snyder BD, Cordio MA, Nazarian A, Kwak SD, Chang DJ, Entezari V, et al. Noninvasive prediction of fracture risk in patients with metastatic cancer to the spine. *Clin Cancer Res* 2009;**15**:7676–83¹²⁸
- Soerdjbalie-Maikoe V, Pelger RCM, Nijeholt GABL, Arndt JW, Zwinderman AH, Bril H, et al. Bone scintigraphy predicts the risk of spinal cord compression in hormone-refractory prostate cancer. *Eur J Nucl Med Mol Imaging* 2004;**31**:958–63¹²⁹
- Sun J-M, Ahn JS, Lee S, Kim JA, Lee J, Park YH, et al. Predictors of skeletal-related events in non-small cell lung cancer patients with bone metastases. *Lung Cancer* 2011;**71**:89–93¹³⁰
- Talcott JA, Stomper PC, Drislane FW, Wen PY, Block CC, Humphrey CC, et al. Assessing suspected spinal cord compression: a multidisciplinary outcomes analysis of 342 episodes. *Support Care Cancer* 1999;**7**:31–8.¹³¹
- Taneichi H, Kaneda K, Takeda N, Abumi K, Satoh S. Risk factors and probability of vertebral body collapse in metastases of the thoracic and lumbar spine. *Spine* 1997;**22**:239–45⁸⁹
- Venkitaraman R, Sohaib SA, Barbachano Y, Parker CC, Khoo V, Huddart RA, et al. Detection of occult spinal cord compression with magnetic resonance imaging of the spine. *Clin Oncol (R Coll Radiol)* 2007;**19**:528–31¹³²
- Venkitaraman R, Sohaib SA, Barbachano Y, Parker CC, Huddart RA, Horwich A, et al. Frequency of screening magnetic resonance imaging to detect occult spinal cord compromise and to prevent neurological deficit in metastatic castration-resistant prostate cancer. *Clin Oncol (R Coll Radiol)* 2010;**22**:147–52¹³³
-