Type I:

	Health scenario A	Health scenario B
	You live in <b>H</b> for <b>T</b> and then die	You live in full health for VT and then die
Which scenario do you		
think is better?		

Type II:

	Health scenario A	Health scenario B
	<b>P</b> lives in <b>H</b> for <b>T</b> and then dies	<b>P</b> lives in full health for <b>VT</b> and then dies
Which scenario do you		
think is better?		

Type III:

	Health scenario A	Health scenario B
	You live for L in full health followed	You live in full health for $(L + VT)$ and
	by $\mathbf{T}$ in $\mathbf{H}$ and then die	then die
Which scenario do you		
think is better?		

Type IV:

	Health scenario A	Health scenario B
	<b>P</b> lives for <b>L</b> in full health followed	<b>P</b> lives in full health for $(L + VT)$ and then
	by <b>T</b> in <b>H</b> and then dies	die
Which scenario do you		
think is better?		

Type V:

	Health scenario A	Health scenario B
	You live in <b>H</b> for <b>T</b> and then die. You	You live in full health for VT and then
	have ${f S}$ with your own health/life during	die
	this time	
Which scenario do		
you think is better?		

Type VI:

	Health scenario A	Health scenario B
	You live in the following health state for T	
	and then die.	
	Unable to walk about	
	Unable to wash or dress yourself	
	Unable to do your usual activities	You will die immediately
	Extreme pain or discomfort	
	Extreme anxiety or depression	
Which scenario do		
you think is better?		

## Type VII:

	Health scenario A	Health scenario B
	Slight problems in walking about	Slight problems in walking about
	Moderate problems washing or dressing yourself	Severe problems washing or dressing yourself
	Severe problems doing your usual activities	Unable to do your usual activities
	Severe pain or discomfort	Extreme pain or discomfort
	Severely anxious or depressed	Not anxious or depressed
	Live for 5 years and then die	Live for 10 years and then die
Which scenario		
do you think is		
better?		

Type VIII:

## HEALTH SCENARIO A

## HEALTH SCENARIO B

	You live in full health for 5 years	
	Then you live for 5 years with the following:	
	Slight problems in walking about	
	No problems washing or dressing yourself	You live in <b>full health</b> for 9 years
	Moderate problems doing your usual activities	
	Slight pain or discomfort	
	Not anxious or depressed	
	Then you die	Then you die
Which scenario		
do you think is		
better?		

## Type IX:

You live in <b>full health</b> for 4 years
Then you die
Then you die
Then you die