

**Sedation in intensive care: Full text screening form**

**Assessor initials:**

**Date:**




**Study ID** (Surname of first author + year of publication)

**Type of study**

Q1. Is the study either:

An RCT in which people are randomised to receive either dexmedetomidine or clonidine (as intervention and comparator) OR

An RCT in which people are randomised to receive either dexmedetomidine or clonidine (as intervention) and propofol or benzodiazepines (as comparator)

Yes	Unclear	No
		
Go to next section		Exclude




**Participants in the study**

Q2. Are the participants all of the following:

Adults AND

In ICU AND

Mechanically ventilated/require mechanical ventilation

Yes	Unclear	No
		
Go to next section		Exclude

**Outcomes reported**

Q3. Does the study report any of the following:

Mortality

Duration of mechanical ventilation

Ventilator free days

Length of ICU stay

Adverse events (incl. rate of hypotension/hypertension/bradycardia/respiratory depression/delirium/coma/unplanned or accidental removal of lines or catheters)

Unpleasant side effects (e.g. unpleasant memories, diarrhea, constipation)

Duration of weaning

Time in target sedation range

Proportion of patients in target sedation range




Discharge readiness

Extubation readiness

Length of hospital stay

QoL

Costs

Yes	Unclear	No
		
Go to next section		Exclude

**Reasons for decision**

Include    Unclear    Exclude