

## Data Extraction Form: HTA 11/95/03 Interventions to prevent postnatal depression

<b>Initials of first data extractor:</b>		<b>Initials of second data extractor:</b>	
<b>Ref ID:</b>			
	<b>Citation</b>	<b>Author contact details</b>	
1.			
<b>Characteristics</b>		<b>Options</b>	
Intervention type		<input type="checkbox"/> Complementary and alternative medicines (e.g. music, massage, aromatherapy, Japanese herbal medicine) <input type="checkbox"/> Maternity care organisation <input type="checkbox"/> Mind-body interventions (e.g. acupuncture, autogenic training, auto-suggestion, biofeedback, guided imagery, hypnosis, hypnotherapy, meditation, prayer, sleep deprivation, tai-chi, yoga) <input type="checkbox"/> Pharmacological (e.g. antidepressants, calcium, omega-3 supplements, hormone therapy, thyroxine) <input type="checkbox"/> Psychological / psychotherapeutic / psychoeducational <input type="checkbox"/> Social support (e.g. exercise, lay support, peer support) <input type="checkbox"/> Other:	
Intervention mode		<input type="checkbox"/> Group <input type="checkbox"/> Individual <input type="checkbox"/> On-line <input type="checkbox"/> Other:	
Intervention provider		<input type="checkbox"/> Doula <input type="checkbox"/> Health visitor <input type="checkbox"/> Midwife <input type="checkbox"/> Nurse <input type="checkbox"/> Psychiatrist <input type="checkbox"/> Psychologist <input type="checkbox"/> Other:	
Intervention duration		<input type="checkbox"/> Single contact <input type="checkbox"/> Multiple contact	
Intervention time		<input type="checkbox"/> Antenatal only <input type="checkbox"/> During labour <input type="checkbox"/> Antenatal and postnatal <input type="checkbox"/> Postnatal only	
Sample selection criteria		<input type="checkbox"/> <b>Universal Preventive Interventions:</b> targeted at a whole population group not identified on the basis of increased risk <input type="checkbox"/> <b>Selective Preventive Interventions:</b> for population subgroups whose risk of developing postnatal depression is higher than average (e.g. psychosocial risk factors) <input type="checkbox"/> <b>Indicated Preventive Interventions:</b> for high-risk women identified as having a predisposition for PND but who do not meet diagnostic criteria for PND.	
<b>RCT details</b>			
<b>Country:</b> <input type="checkbox"/> Australia <input type="checkbox"/> Canada <input type="checkbox"/> China <input type="checkbox"/> France <input type="checkbox"/> Hong Kong <input type="checkbox"/> India <input type="checkbox"/> Italy <input type="checkbox"/> Japan <input type="checkbox"/> Korea <input type="checkbox"/> Mexico <input type="checkbox"/> Netherlands <input type="checkbox"/> Norway <input type="checkbox"/> South Africa <input type="checkbox"/> Sweden <input type="checkbox"/> Taiwan <input type="checkbox"/> UK <input type="checkbox"/> US <input type="checkbox"/> Other:			
<b>Study setting:</b> <input type="checkbox"/> Antenatal clinic <input type="checkbox"/> Home visits <input type="checkbox"/> Labour / delivery ward <input type="checkbox"/> Postnatal ward <input type="checkbox"/> Primary care <input type="checkbox"/> Other:			
<b>Number of centres:</b>			
<b>Recruitment:</b> <input type="checkbox"/> Pregnancy <input type="checkbox"/> During labour <input type="checkbox"/> Postnatally			
<b>Funding:</b> <input type="checkbox"/> Government <input type="checkbox"/> Private <input type="checkbox"/> Scholarship / award <input type="checkbox"/> Other:			
<b>Aim of the study</b>			
<b>Intervention</b> (same order as in the paper) Provide description of experimental interventions and controls (e.g. pharmacological, social support, mind-body intervention, CAMs, maternity care, location, number and length of sessions, number per group, professional background making diagnosis)			
<b>Intervention 1:</b> <b>Intervention 2 (if applicable):</b> <b>Control:</b>			

Participant recruitment				
<b>Recruitment dates:</b>				
Total number of women randomised (n) / screened (n):      /      %				
<b>Inclusion criteria:</b>				
<b>Exclusion criteria:</b> (e.g. risk, history, condition, antenatal anxiety, antenatal depression, social support, life events, domestic violence, method for identification)				
Baseline characteristics				
	<b>All (Total sample)</b> Mean (SD) / n (%)	<b>Intervention 1</b> Mean (SD) / n (%)	<b>Intervention 2</b> Mean (SD) / n (%)	<b>Control</b> Mean (SD) / n (%)
Age in years: Mean (SD) or range				
Number randomised				
EPDS				
Other measure				
Race/ethnicity				
Socio-demographic characteristics				
Education				
Diagnosis at baseline				
Populations of antenatal women				
Populations of early postnatal women				
Co-therapy or medication: (Specify dose if applicable)				
Outcomes summary				
1. Maternal outcomes (e.g. depressive symptoms, morbidity, mortality): <input type="checkbox"/> Yes <input type="checkbox"/> No				
2. Neonatal, post-neonatal and infant outcomes (e.g. Apgar, morbidity, mortality, injury, development): <input type="checkbox"/> Yes <input type="checkbox"/> No				
3. Family outcomes (e.g. dyadic relationship, discord, separation / divorce, abuse, violence): <input type="checkbox"/> Yes <input type="checkbox"/> No				
4. Process outcomes (e.g. uptake, did not receive intervention, discontinued intervention, number of sessions received): <input type="checkbox"/> Yes <input type="checkbox"/> No				
5. Cost of use of service outcomes: <input type="checkbox"/> Yes <input type="checkbox"/> No				
Primary outcome:				
Secondary outcomes:				
Timing of outcome assessments (e.g. 6 weeks postnatally, 12 weeks, 6 months, 12 months):				
Total length of follow up:				
Did not receive allocated intervention: n=				
Number of participants				
	<b>All (Total sample)</b>	<b>Intervention 1</b>	<b>Intervention 2</b>	<b>Control</b>
Randomised				
Losses to follow-up/drop out/sample attrition	Time: n:	Time: n:	Time: n:	Time: n:
Number analysed	Time: n:	Time: n:	Time: n:	Time: n:
Results: Maternal outcomes				
Report n, Mean (SD) for each time of assessment. Report all categorical outcomes in a separate table or provide details in the notes column				
<b>Outcome</b> (Note whether it is actual score or change score)	<b>Intervention 1</b> Time: n: Mean (SD)	<b>Intervention 2</b> Time: n: Mean (SD)	<b>Control</b> Time: n: Mean (SD)	<b>Categorical outcomes / Notes</b>
Anxiety measure:				
BDI				
CES-D				
Diagnostic interviews:				
DSM-IV				
EPDS				

GHQ				
Hopkins Scale				
ICD-10				
Kellner symptom questionnaire				
MADRS				
Maternal dissatisfaction with intervention				
Maternal morbidity:				
Maternal mortality				
Maternal perceived support:				
Maternal-infant attachment:				
Mood measure:				
Other depressive symptoms:				
POMS				
SCID				
Self-harm:				
Stress measures:				
Suicide attempts				
Others:				
<b>Results: Neonatal, post-neonatal and infant outcomes</b>				
<i>Report n, Mean (SD) for each time of assessment. Report all categorical outcomes in a separate table or provide details in the notes column</i>				
<b>Outcome</b> <i>(Note whether it is actual score or change score)</i>	<b>Intervention 1</b> <i>Time:</i> <i>n:</i> <i>Mean (SD)</i>	<b>Intervention 2</b> <i>Time:</i> <i>n:</i> <i>Mean (SD)</i>	<b>Control</b> <i>Time:</i> <i>n:</i> <i>Mean (SD)</i>	<b>Categorical outcomes / Notes</b>
Apgar				
Child abuse:				
Infant developmental assessments:				
Infant health parameters:				
Injury:				
Morbidity:				
Mortality				
Neglect:				
Others:				
<b>Results: Family outcomes</b>				
<i>Report n, Mean (SD) for each time of assessment. Report all categorical outcomes in a separate table or provide details in the notes column</i>				
<b>Outcome</b> <i>(Note whether it is actual score or change score)</i>	<b>Intervention 1</b> <i>Time:</i> <i>n:</i> <i>Mean (SD)</i>	<b>Intervention 2</b> <i>Time:</i> <i>n:</i> <i>Mean (SD)</i>	<b>Control</b> <i>Time:</i> <i>n:</i> <i>Mean (SD)</i>	<b>Categorical outcomes / Notes</b>
Abuse:				
Dyadic relationship:				
Marital discord:				
PSI				
Separation / divorce				
Violence:				
Others:				
<b>Results: Process outcomes, uptake</b>				
<i>(e.g. did not receive intervention, discontinued intervention, number of sessions received)</i>				
<i>Report n, Mean (SD) for each time of assessment. Report all categorical outcomes in a separate table or provide details in the notes column</i>				
<b>Outcome</b> <i>(Note whether it is actual score or change score)</i>	<b>Intervention 1</b> <i>Mean (SD)</i>	<b>Intervention 2</b> <i>Mean (SD)</i>	<b>Control</b> <i>Mean (SD)</i>	<b>Categorical outcomes / Notes</b>
Please specify:				
<b>Authors' conclusion</b>				
<b>Reviewers' conclusion</b>				