

Reviewer ID:

Data extraction date:

GENERAL INFORMATION	
Report title:	
First author / contact details	
Publication year	
Publication status:	Full-text paper <input type="checkbox"/> Conference abstract <input type="checkbox"/> Personal communication <input type="checkbox"/> Other unpublished reports <input type="checkbox"/>
Journal yy:vol(issue):pp	
Language (if non-English):	
Study IDs of any linked reports:	
Study funding sources <i>(including role of funders)</i>	
Possible conflicts of interest <i>(for study authors)</i>	
STUDY ELIGIBILITY	
Type of study: RCT <input type="checkbox"/> Non-randomised comparative study <input type="checkbox"/> Case series <input type="checkbox"/>	
Type of intervention: Dietary / lifestyle: Vitamin B6 <input type="checkbox"/> Vitamin B12 <input type="checkbox"/> Ginger <input type="checkbox"/> Acupuncture <input type="checkbox"/> Acupressure <input type="checkbox"/> Hypnosis <input type="checkbox"/> Antiemetic drugs: Antihistamines <input type="checkbox"/> Dopamine antagonists <input type="checkbox"/> 5-HT receptor antagonists <input type="checkbox"/> Corticosteroids <input type="checkbox"/> Doxylamine-Pyridoxine <input type="checkbox"/> Other (details below) <input type="checkbox"/> Intravenous fluids: <input type="checkbox"/> Enteral and total parenteral nutrition Enteral feeding <input type="checkbox"/> Total parenteral nutrition <input type="checkbox"/> Other Intervention <input type="checkbox"/> _____ Comparator: No treatment <input type="checkbox"/> Treatment as usual (details below) <input type="checkbox"/> Alternative intervention (details below) <input type="checkbox"/> _____ Comparator not applicable:	

Women with severe symptoms	<input type="checkbox"/>
<b>Participants:</b>	
Gestational age ≤20 weeks <input type="checkbox"/>	
Symptom severity:	
PUQE score ≥ 13 <input type="checkbox"/>	Rhodes score ≥ 33 <input type="checkbox"/>
Author defined scale (provide details below) <input type="checkbox"/>	
Percentage experiencing symptoms >80% <input type="checkbox"/>	
<b>Primary outcomes:</b>	
<b>Severity of symptoms:</b>	
PUQE <input type="checkbox"/>	Rhodes Index <input type="checkbox"/>
Visual Analogue scales <input type="checkbox"/>	McGill Nausea Questionnaire <input type="checkbox"/>
Nausea and Vomiting of Pregnancy Instrument <input type="checkbox"/>	
<b>Secondary outcomes:</b>	
<b>Maternal-physical:</b>	
Admission/readmission rate <input type="checkbox"/>	Length of hospital stay <input type="checkbox"/>
Antiemetic / other medication use <input type="checkbox"/>	Amount/duration IV fluid administration <input type="checkbox"/>
Enteral/total Parenteral nutrition <input type="checkbox"/>	Adverse events <input type="checkbox"/>
Economic costs (hospital/medical care) <input type="checkbox"/>	Adverse pregnancy outcomes <input type="checkbox"/>
Weight loss <input type="checkbox"/>	Therapeutic termination of pregnancy <input type="checkbox"/>
Other author defined NVP scale <input type="checkbox"/>	
<b>Maternal –psychosocial:</b>	
Quality of life (eg. SF-12/SF-36 score) <input type="checkbox"/>	General Health Questionnaire <input type="checkbox"/>
Pregnancy specific QoL instrument <input type="checkbox"/>	NVP specific questionnaire <input type="checkbox"/>
Satisfaction with care <input type="checkbox"/>	Direct costs to woman/family <input type="checkbox"/>
Time lost from work <input type="checkbox"/>	Edinburgh post natal depression score <input type="checkbox"/>
<b>Fetal/Neonatal:</b>	
Congenital abnormality <input type="checkbox"/>	Low birth weight (<2.5kg) <input type="checkbox"/>
Small for gestational age (<10 <sup>th</sup> centile) <input type="checkbox"/>	Pre-term birth (<37 weeks gestation) <input type="checkbox"/>
5 minute APGAR <input type="checkbox"/>	Stillbirth/IUD <input type="checkbox"/>
Neonatal death <input type="checkbox"/>	Spontaneous miscarriage <input type="checkbox"/>
Admission to special care baby unit <input type="checkbox"/>	Long term infant outcomes <input type="checkbox"/>
<b>INCLUDE</b> <input type="checkbox"/>	<b>EXCLUDE</b> <input type="checkbox"/>
<b>Reasons for exclusion:</b>	

**DO NOT PROCEED IF STUDY EXCLUDED FROM REVIEW**

ADDITIONAL STUDY INFORMATION			
Population and setting			
	Intervention	Comparator	Location in text
Population description			
Setting (including country / location / social context etc)			
Inclusion exclusion criteria			
Method/s of recruitment			
Informed consent obtained	Yes <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/>	Yes <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/>	
Notes:			
Methods			
	Descriptions as stated in report / paper		Location in text
Aim of study			
Design (no of arms)			
Unit of allocation (by individuals, cluster / groups)			
Start date			
End date			
Total study duration			
Ethical approval needed / obtained for study	Yes <input type="checkbox"/>	No <input type="checkbox"/>	Unclear <input type="checkbox"/>
Notes:			
Participant characteristics			
	Intervention	Comparator	Location in text
Number of patients enrolled:			
- Randomised (RCTs only), n (%)			
- Included (RCTs only), n (%)			
- Completed, n (%)			

- Available for follow-up, n (%)			
- Withdrew/lost to follow-up, with reasons, n (%)			
- Number analysed, n (%)			
Age (mean/median, SD/range)			
Ethnicity, n (%)			
Smoking status (give n (%) of smokers)			
BMI at baseline, (mean/median, SD/range)			
Weight at baseline (mean/median, SD range)			
Singleton pregnancy only (if no, give n (%))			
Gestational age at onset (week: mean/median, SD/range)			
Gestational age at primary admission (week: mean/median, SD/range)			
Gestational age at study entry / randomisation (week: mean/median, SD/range)			
Primiparas only (if no, give %)			
Obstetric history (previous NVP)			
Pre-existing medical conditions (please specify)			
NVP / HG Severity:			
- PUQE (Mean/median, SD/range)			
- Rhodes Index (Mean/median, SD/range)			
- McGill Nausea Questionnaire (Mean/median, SD/range)			

- Nausea and Vomiting of Pregnancy Instrument (Mean/median, SD/range)			
- Other scale (details)			
Other baseline characteristics (please specify):			

**INTERVENTION GROUPS** (*copy and paste table for each intervention group and comparator*)

**Intervention Group 1**

	Description in text	Location in text
Group name		
No. randomised to group		
Theoretical basis (include key references)		
Description of intervention (include sufficient detail for replication eg content, dose, components)		
Duration of treatment period		
Timing of treatment (eg frequency, duration of each episode)		
Delivery (eg mechanism, medium, intensity, fidelity)		
Providers (eg number, profession, training, gender / ethnicity / age if relevant)		
Co-interventions		
Economic variables (eg intervention cost, changes in other others as result of intervention)		

Notes:

**OUTCOMES** *(copy and paste table for each outcome)***Outcome 1**

	Description as stated in report / paper	Location in text
Outcome name		
Time points measured		
Time points reported		
Outcome definition (with diagnostic criteria if relevant)		
Personal measuring/ reporting		
Unit of measurement (if relevant)		
Scales: upper and lower limits (indicate whether high or low score is good)		
Is outcome/tool validated?	Yes <input type="checkbox"/> No <input type="checkbox"/> Unclear <input type="checkbox"/>	
Imputations of missing data (eg assumptions made for ITT analysis)		
Assumed risk estimate (eg baseline or population risk note in background)		
Power		

Notes:

**RESULTS****Dichotomous outcomes**

	Description as stated in report/paper	Location in text
Comparison		
Outcome		
Subgroup		
Timepoint (specify whether from start or end of intervention)		
Results	Intervention	Comparison

	No. events	No. participants		No. events	No. participants		
No. missing participants and reasons							
No. participants moved from other group and reasons							
Any other results reported							
Unit of analysis (by individuals, cluster/groups or body parts)							
Statistical methods used							
Notes:							
<b>Continuous outcome</b>							
	<b>Description as stated in report/paper</b>						<b>Location in text</b>
Comparison							
Outcome							
Subgroup							
Timepoint (specify whether from start or end of intervention)							
Post-intervention or change from baseline?							
Results	Intervention			Comparison			
	Mean	SD (or other variance)	No. participants	Mean	SD (or other variance)	No. participants	

No. missing participants and reasons				
No. participants moved from other group and reasons				
Any other results reported				
Unit of analysis (individuals, cluster/groups or body parts)				
Statistical methods used				
Notes:				
<b>Other outcome</b>				
	<b>Description as stated in report/paper</b>			<b>Location in text</b>
Comparison				
Outcome				
Subgroup				
Timepoint (specify whether from start or end of intervention)				
Results	Intervention result	SD (or other variance)	Control result	SD (or other variance)
	Overall results		SE (or other variance)	
No. participants	Intervention		Control	
No. missing participants and reasons				
No. participants moved from other group and reasons				



Any other results reported		
Unit of analysis (by individuals, cluster/groups or body parts)		
Statistical methods used		
Notes:		
Conclusion as reported by the authors of the study		
Additional information and comments		