

Full record citation	Reason
A randomised controlled trial to examine the efficacy of e-cigarettes compared with nicotine replacement therapy, when used within the UK stop smoking service (Project record)2015; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000774/frame.html .	Ongoing study
AboElNaga HH. Electronic cigarettes: not an advantageous alternative to conventional smoking in asthma. <i>Egyptian Journal of Chest Diseases and Tuberculosis</i> . 2018;67(4):429-34.	Wrong comparator
Abramson Cancer Center of the University of P. Extended Duration Varenicline for Smoking Among Cancer Patients: A Clinical Trial. https://ClinicalTrials.gov/show/NCT02851901 ; 2012.	Ongoing study
Actrn, Nosa V GM. TALANOA Samoa: A randomised controlled trial to evaluate the efficacy of a cessation support programme for smokers delivered via radio. 2013.	Wrong or missing intervention
Actrn. An open-label randomised pragmatic policy trial examining effectiveness of short-term use of Nicotine Replacement Therapy (NRT) vs short- or long-term use of NRT vs short- or long-term use of NRT or electronic nicotine delivery systems for smoking cessation in cigarette smokers.	Participants received same drug
Actrn. Improving radiotherapy outcomes with smoking cessation: Feasibility trial in head and neck cancer patients. 2014.	Ongoing study
Agency for Healthcare R, Quality, Agency for Healthcare R, Quality. Treating Tobacco Use and Dependence: 2008 Update. Content last reviewed June 2015. Rockville, MD.; 2008.	Wrong study design
Ahmadi J, Ashkani H, Ahmadi M, Ahmadi N. Twenty-four week maintenance treatment of cigarette smoking with nicotine gum, clonidine and naltrexone. <i>Journal of Substance Abuse Treatment</i> . 2003;24(3):251-5.	Wrong comparator
AhnAllen CG, Bidwell LC, Tidey JW. Cognitive effects of very low nicotine content cigarettes, with and without nicotine replacement, in smokers with schizophrenia and controls. <i>Nicotine & Tobacco Research</i> . 2015;17(5):510-4.	Wrong study design
Ahnallen CG, Bidwell LC, Tidey JW. Cognitive effects of very low nicotine content cigarettes, with and without nicotine replacement, in smokers with schizophrenia and controls. <i>Nicotine and Tobacco Research</i> . 2015;17(5):510-4.	Wrong study design
Akers L, Severson HH, Andrews JA, Lichtenstein E. Cost-effectiveness of self-help smokeless tobacco cessation programs. <i>Nicotine & Tobacco Research</i> . 2007;9(9):907-14.	Wrong or missing intervention
Akhavan S, Nguyen LC, Chan V, Saleh J, Bozic KJ. Impact of Smoking Cessation Counseling Prior to Total Joint Arthroplasty. <i>Orthopedics</i> . 2017;40(2):E323-E8.	Awaiting classification
Akhu-Zaheya LM, Shiyab WY. The effect of short message system (SMS) reminder on adherence to a healthy diet, medication, and cessation of smoking among adult patients with cardiovascular diseases. <i>Int J Med Inf</i> . 2017;98:65-75.	Wrong study design
Albedah AM, Khalil MK, Khalil AA, Elolemy AT. Use of the target group index survey to evaluate the cigarette smoking profile in Saudi Arabia. <i>Saudi Medical Journal</i> . 2011;32(10):1055-9.	Wrong study design
Albert Einstein College of Medicine I, National Institute on Drug A. Varenicline Smoking Cessation Treatment for Methadone Maintenance Patients. https://ClinicalTrials.gov/show/NCT01378858 ; 2011.	Participants received same drug
Albert Einstein College of Medicine I. Smoking Cessation Treatment for Methadone Maintenance Patients. https://ClinicalTrials.gov/show/NCT01027754 ; 2009.	Ongoing study
Al-Delaimy WK, Myers MG, Leas EC, Strong DR, Hofstetter CR. E-cigarette use in the past and quitting behavior in the future: a population-based study. <i>Am J Public Health</i> . 2015;105(6):1213-9.	Wrong study design

Al-Halabi S, Fernandez-Artamendi S, Diaz-Mesa EM, Garcia-Alvarez L, Florez G, Martinez-Santamaria E, et al. Tobacco and cognitive performance in schizophrenia patients: the design of the COGNICO study. <i>Adicciones</i> . 2016;724.	Ongoing study
Al-Halabi S, Fernandez-Artamendi S, Diaz-Mesa EM, Garcia-Alvarez L, Florez G, Santamaria EM, et al. Tobacco and cognitive performance in schizophrenia patients: The design of the COGNICO study. <i>Tabaco y rendimiento cognitivo en pacientes con esquizofrenia: Diseno del estudio COGNICO</i> . 2017;29(1):6-12.	Ongoing study
Allen A, Vogel RI, Meier E, Anderson A, Jensen J, Severson HH, et al. Gender differences in snus versus nicotine gum for cigarette avoidance among a sample of US smokers. <i>Drug & Alcohol Dependence</i> . 2016;168:8-12.	Wrong comparator
Alliance for Clinical Trials in O, National Cancer I, Pfizer. Smoking Cessation Prior to Surgery to Reduce Surgical Complications in Patients With Newly Diagnosed Lung Cancer Who Smoke Cigarettes. https://ClinicalTrials.gov/show/NCT02856581 ; 2016.	Ongoing study
Alliance for Clinical Trials in O, National Cancer I. Comparison of Nicotine Inhaler and/or Bupropion in Helping People to Stop Smoking and Prevent the Recurrence of Smoking. https://ClinicalTrials.gov/show/NCT00033592 ; 2002.	Ongoing study
Alonso Fernandez M, Franco Vidal A, Lopez Sampedro P, Garcia Lavandera J. [Middle-term effectiveness of a support program for smokers implemented in primary care]. <i>Atencion Primaria</i> . 2002;30(9):541-6.	Wrong study design
Alterman AI, Gariti P, Cook TG, Cnaan A. Nicodermal patch adherence and its correlates. <i>Drug and Alcohol Dependence</i> . 1999;53(2):159-65.	Participants received same drug
Alterman AI, Gariti P, Mulvaney F. Short- and long-term smoking cessation for three levels of intensity of behavioral treatment. <i>Psychology of Addictive Behaviors</i> . 2001;15(3):261-4.	Participants received same drug
American University of Beirut Medical C, Pfizer. Varenicline (Chantix) Treatment for Waterpipe Smoking Cessation. https://ClinicalTrials.gov/show/NCT02575183 ; 2016.	Ongoing study
An L, Zhu SH, Nelson D, Arikian N, Nugent S, Partin M, et al. Telestop: A randomized trial of increased access to behavioral and pharmacological therapy for smoking cessation. <i>Nicotine & Tobacco Research</i> . 2005;7(4):689-.	Cannot locate full text
An LC, Zhu SH, Nelson DB, Arikian NJ, Nugent S, Partin MR, et al. Benefits of telephone care over primary care for smoking cessation: A randomized trial. <i>Archives of Internal Medicine</i> . 2006;166(5):536-42.	Wrong study design
Andrew Howe AJ, Shand J, Menown I. Real world evaluation of varenicline therapy within a secondary care smoking cessation programme. <i>European Journal of Preventive Cardiology</i> . 2012;1):S40.	Cannot locate full text
Andrews JA, Severson HH, Lichtenstein E, Gordon JS, Barckley MF. Evaluation of a dental office tobacco cessation program: Effects on smokeless tobacco use. <i>Annals of Behavioral Medicine</i> . 1999;21(1):48-53.	Wrong or missing intervention
Angersbach D. Varenicline for smoking cessation is tolerated and effective. [German]. <i>Medizinische Monatsschrift fur Pharmazeuten</i> . 2013;36(4):165-7.	Wrong study design
Anonymous. Varenicline plus nicotine replacement therapy more effective than varenicline alone. <i>Prescriber</i> . 2014;25(19):31-.	Cannot locate full text
Anonymous. Varenicline plus NRT better than varenicline alone for stopping smoking. <i>Drug and Therapeutics Bulletin</i> . 2014;52(9):100.	Wrong study design
Aparici M, Fernandez Gonzalez AL, Alegria E. [Clonidine in the treatment of tobacco withdrawal. A comparison with nicotine chewing gum]. <i>Revista Clinica Espanola</i> . 1994;194(6):453-6.	Wrong comparator

Aristotle University Of T, Ltd AAISCS, National Association Of General Practitioners In B. Combined Bio- and Neuro- Feedback vs. Varenicline Use for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02991781 ; 2017.	Participants received same drug
Arterburn D, Sofer T, Boudreau DM, Bogart A, Westbrook EO, Theis MK, et al. Long-Term Weight Change after Initiating Second-Generation Antidepressants. <i>Journal of Clinical Medicine</i> . 2016;5(4).	Wrong comparator
Asfar T, Klesges RC, Sanford SD, Sherrill-Mittleman D, Robison LL, Hudson MM, et al. Trial design: The St. Jude Children's Research Hospital Cancer Survivors Tobacco Quit Line study. <i>Contemporary Clinical Trials</i> . 2010;31(1):82-91.	Linked to excluded study
Assistance Publique - Hôpitaux de P, Ministry of Health F, National Cancer Institute F. Smoking Cessation for Patients With Acute Coronary Syndrome (ACS). https://ClinicalTrials.gov/show/NCT00987597 ; 2008.	Participants received same drug
Assistance Publique - Hôpitaux de P. Determination of Carcinogens Nicotine in Electronic Cigarettes Users. https://ClinicalTrials.gov/show/NCT02897401 ; 2016.	Ongoing study
Association for the Japanese Academy of Maxillofacial I. Tobacco Cessation Intervention Study for Oral Diseases. https://ClinicalTrials.gov/show/NCT02737176 ; 2016.	Ongoing study
Asvat Y, Cao DC, Africk JJ, Matthews A, King A. Feasibility and Effectiveness of a Community-Based Smoking Cessation Intervention in a Racially Diverse, Urban Smoker Cohort. <i>American Journal of Public Health</i> . 2014;104:S620-S7.	Wrong or missing intervention
Attwood AS, Williams T, Adams S, McClernon FJ, Munafo MR. Effects of varenicline and cognitive bias modification on neural response to smoking-related cues: study protocol for a randomized controlled study. 2014;15:391.	Ongoing study
Aubin HJ, Peiffer G, Stuebner-Delbarre A, Vicaut E, Jeanpetit Y, Solesse A, et al. The French Observational Cohort of Usual Smokers (FOCUS) cohort: French smokers perceptions and attitudes towards smoking cessation. <i>Bmc Public Health</i> . 2010;10.	Wrong or missing intervention
Augusta U, National Cancer I. The Impact of a Tobacco Control Intervention in African-American Families. https://ClinicalTrials.gov/show/NCT01164306 ; 2007.	Ongoing study
Aung MN, Yuasa M, Lorga T, Moolphate S, Fukuda H, Kitajima T, et al. Evidence-based new service package vs. routine service package for smoking cessation to prevent high risk patients from cardiovascular diseases (CVD): Study protocol for randomized controlled trial. <i>Trials</i> . 2013;14 (1) (no pagination)(419).	Ongoing study
Aveyard P, Brown K, Saunders C, Alexander A, Johnstone E, Munafo MR, et al. Weekly versus basic smoking cessation support in primary care: a randomised controlled trial. <i>Thorax</i> . 2007;62(10):898-903.	Wrong or missing intervention
Baker A, Borland R, Bonevski B, Castle DJ, Williams J, Segan C, et al. Quitlink: Accessible smoking cessation support for people living with severe and enduring mental illness. <i>Schizophrenia Bulletin</i> . 2018;44 (Supplement 1):S151.	Wrong or missing intervention
Balmford J, Borland R, Hammond D, Cummings KM. Adherence to and reasons for premature discontinuation from stop-smoking medications: data from the ITC Four-Country Survey. <i>Nicotine & Tobacco Research</i> . 2011;13(2):94-102.	Wrong study design
Balmford J, Borland R, Hammond D, Cummings KM. Adherence to and reasons for premature discontinuation from stop-smoking medications: data from the ITC Four-Country Survey. <i>Nicotine & Tobacco Research</i> . 2011;13(2):94-102.	Wrong study design
Barrett SP, Campbell ML, Temporale K, Good KB. The acute effect of Swedish-style snus on cigarette craving and self-administration in male and female smokers. <i>Human Psychopharmacology</i> . 2011;26(1):58-62.	Wrong study design

Barrett SP, Darredeau C. The acute effects of nicotine on the subjective and behavioural responses to denicotinized tobacco in dependent smokers. <i>Behavioural Pharmacology</i> . 2012;23(3):221-7.	Wrong study design
Barrueco M, Jimenez Ruiz C, Palomo L, Torrecilla M, Romero P, Riesco JA. Veracity of smokers' reports of abstinence at smoking cessation clinics. [Spanish]. <i>Archivos de Bronconeumologia</i> . 2005;41(3):135-40.	Wrong study design
Bastian LA, Fish LJ, Peterson BL, Biddle AK, Garst J, Lyna P, et al. Proactive recruitment of cancer patients' social networks into a smoking cessation trial. <i>Contemporary Clinical Trials</i> . 2011;32(4):498-504.	Wrong study design
Batra A, Collins SE, Schroeter M, Eck S, Torchalla I, Buchkremer G. A cluster-randomized effectiveness trial of smoking cessation modified for at-risk smoker subgroups. <i>Journal of Substance Abuse Treatment</i> . 2010;38(2):128-40.	Wrong comparator
Beard E, Aveyard P, Brown J, West R. Assessing the association between the use of NRT for smoking reduction and attempts to quit smoking using propensity score matching. <i>Drug and Alcohol Dependence</i> . 2012;126(3):354-61.	Wrong study design
Beard E, Aveyard P, McNeill A, Michie S, Fidler JA, Brown J, et al. Mediation analysis of the association between use of NRT for smoking reduction and attempts to stop smoking. <i>Psychology and Health</i> . 2012;27(9):1118-33.	Wrong study design
Beard E, McNeill A, Aveyard P, Fidler J, Michie S, West R. Association between use of nicotine replacement therapy for harm reduction and smoking cessation: A prospective study of English smokers. <i>Tobacco control</i> . 2013;22(2):118-22.	Wrong study design
Beard E, McNeill A, Aveyard P, Fidler J, Michie S, West R. Use of nicotine replacement therapy for smoking reduction and during enforced temporary abstinence: a national survey of English smokers. <i>Addiction</i> . 2011;106(1):197-204.	Wrong study design
Beard E, West R. Use of nicotine replacement therapy for smoking reduction and temporary abstinence: an update of beard et al. (2011). <i>Addiction (Abingdon, England)</i> . 2012;107(6):1186-7.	Cannot locate full text
Beard E, West R. Use of nicotine replacement therapy for smoking reduction and temporary abstinence: an update of beard et al. (2011). <i>Addiction</i> . 2012;107(6):1186-7.	Wrong study design
Becker KM, Rose JE, Albino AP. A randomized trial of nicotine replacement therapy in combination with reduced-nicotine cigarettes for smoking cessation. <i>Nicotine and Tobacco Research</i> . 2008;10(7):1139-48.	Wrong study design
Benmarhnia T, Pierce JP, Leas E, White MM, Strong DR, Noble ML, et al. Can E-Cigarettes and Pharmaceutical Aids Increase Smoking Cessation and Reduce Cigarette Consumption? Findings From a Nationally Representative Cohort of American Smokers. <i>American Journal of Epidemiology</i> . 2018;187(11):2397-404.	Wrong study design
Berg KM, Jorenby DE, Baker TB, Fiore MC. Triple smoking cessation therapy with varenicline, nicotine patch and nicotine lozenge: A pilot study to assess tolerability, satisfaction and end-of-treatment quit rates. <i>Journal Of Smoking Cessation</i> . 2017:No-Specified.	Participants received same drug
Berg KM, Jorenby DE, Fiore M, Baker T. Triple smoking cessation therapy with varenicline, nicotine patch and nicotine lozenge: A pilot study to assess tolerability. <i>Journal of General Internal Medicine</i> . 2017;32 (2 Supplement 1):S364.	Participants received same drug
Berger VW. Randomization, permuted blocks, masking, allocation concealment, and selection bias in the Tobacco Quit Line Study. <i>Contemporary Clinical Trials</i> . 2010;31(3):201-.	Linked to excluded study
Berlin I, Covey LS, Jiang HP, Hamer D. Lack of effect of D2 dopamine receptor TaqI A polymorphism on smoking cessation. <i>Nicotine & Tobacco Research</i> . 2005;7(5):725-8.	Wrong or missing intervention

Berlin I, Jacob N, Coudert M, Perriot J, Schultz L, Rodon N. Adjustment of nicotine replacement therapies according to saliva cotinine concentration: the ADONIS trial-a randomized study in smokers with medical comorbidities. <i>Addiction</i> . 2011;106(4):833-43.	Participants received same drug
Bills A, Giang K, La Spisa S, Buckley K, Tsu L, Early N, et al. Impact of a pharmacist-only clinic versus a pharmacist-led multidisciplinary clinic on tobacco cessation rates in a homeless population. <i>Pharmacotherapy</i> . 2015;35 (11):e290.	Wrong study design
Binnington MJ, Zhu AZX, Renner CC, Lanier AP, Hatsukami DK, Benowitz NL, et al. CYP2A6 and CYP2B6 genetic variation and its association with nicotine metabolism in South Western Alaska Native people. <i>Pharmacogenetics and Genomics</i> . 2012;22(6):429-40.	Wrong or missing intervention
Bize R, Stoianov R, Ruffieux C, Ghali W, Paccaud F, Cornuz J. Effectiveness of a low-intensity smoking cessation intervention for hospitalized patients. <i>European Journal of Cancer Prevention</i> . 2006;15(5):464-70.	Participants received same drug
Blak B, Hards M, Thompson M, Dattani H. Impact of the UK public smoking ban on the prescribing of smoking cessation therapy in primary care-a thin database study. <i>Value in Health</i> . 2010;13 (3):A203.	Cannot locate full text
Blak BT, Hards M, Dattani H, Thompson M. Prescribing behaviour of smoking cessation products in the UK - A thin database study. <i>Pharmacoepidemiology and Drug Safety</i> . 2010;19:S282.	Cannot locate full text
Blak BT, Wilson K, Metcalfe M, Maguire A, Hards M. Evaluation of varenicline as an aid to smoking cessation in UK general practice - a THIN database study. <i>Current Medical Research & Opinion</i> . 2010;26(4):861-70.	Wrong study design
Boland VC, Mattick RP, Siahpush M, Barker D, Doran CM, Martire KA, et al. Factors associated with Quitline and pharmacotherapy utilisation among low-socioeconomic status smokers. <i>Addictive Behaviors</i> . 2019;89:113-20.	Linked to excluded study
Bondy SJ, Diemert LM, Victor JC, McDonald PW, Cohen JE. Assessing the reach of nicotine replacement therapy as a preventive public health measure. <i>Chronic Diseases and Injuries in Canada</i> . 2012;33(1):19-28.	Wrong study design
Borland R, Balmford J, Bishop N, Segan C, Piterman L, McKay-Brown L, et al. In-practice management versus quitline referral for enhancing smoking cessation in general practice: a cluster randomized trial. <i>Family Practice</i> . 2008;25(5):382-9.	Wrong or missing intervention
Borland R, Balmford J, Segan C, Livingston P, Owen N. The effectiveness of personalized smoking cessation strategies for callers to a Quitline service. <i>Addiction</i> . 2003;98(6):837-46.	Wrong study design
Borland R, Partos TR, Cummings KM. Systematic biases in cross-sectional community studies may underestimate the effectiveness of stop-smoking medications. <i>Nicotine & Tobacco Research</i> . 2012;14(12):1483-7.	Wrong study design
Borland R, Partos TR, Cummings KM. Systematic biases in cross-sectional community studies may underestimate the effectiveness of stop-smoking medications. <i>Nicotine and Tobacco Research</i> . 2012;14(12):1483-7.	Wrong study design
Boshier A, Wilton LV, Shakir SA. Evaluation of the safety of bupropion (Zyban) for smoking cessation from experience gained in general practice use in England in 2000. <i>European Journal of Clinical Pharmacology</i> . 2003;59(10):767-73.	Wrong study design
Boshier A, Wilton LV, Shakir SAW. Evaluation of the safety of bupropion (Zyban) for smoking cessation from experience gained in general practice use in England in 2000. <i>European Journal of Clinical Pharmacology</i> . 2003;59(10):767-73.	Wrong study design
Boston U, Harvard School of Public H, The Miriam H, Boston Housing A. Intervention Study to Increase Smoking Cessation Rates Among Public Housing Residents. https://ClinicalTrials.gov/show/NCT01651611 ; 2011.	Participants received same drug

Botteri E, de Lange T, Tonstad S, Berstad P. Exploring the effect of a lifestyle intervention on cancer risk: 43-year follow-up of the randomized Oslo diet and antismoking study. <i>J Intern Med.</i> 2018;22:22.	Wrong or missing intervention
Bottorff JL, Johnson JL, Moffat B, Fofonoff D, Budz B, Groening M. Synchronizing clinician engagement and client motivation in telephone counseling. <i>Qualitative Health Research.</i> 2004;14(4):462-77.	Wrong study design
Boyle RG, Enstad C, Asche SE, Thoele MJ, Sherwood NE, Severson HH, et al. A randomized controlled trial of Telephone Counseling with smokeless tobacco users: The ChewFree Minnesota study. <i>Nicotine & Tobacco Research.</i> 2008;10(9):1433-40.	Wrong or missing intervention
Boyle RG, Pronk NP, Enstad CJ. A randomized trial of telephone counseling with adult moist snuff users. <i>American Journal of Health Behavior.</i> 2004;28(4):347-51.	Wrong or missing intervention
Boyle RG, Solberg LI, Magnan S, Davidson G, Alesci NL. Does insurance coverage for drug therapy affect smoking cessation? <i>Health affairs (Project Hope).</i> 2002;21(6):162-8.	Wrong study design
Brandt CJ, Christiansen L, Christoffersen G, Ellegaard H, Joensen M, Madsen A, et al. Effect of diagnosis of "smoker's lung". <i>Lancet.</i> 1997;349(9047):253-.	Wrong or missing intervention
Bredie SJH, Fouwels AJ, Wollersheim H, Schippers GM. Effectiveness of Nurse Based Motivational Interviewing for smoking cessation in high risk cardiovascular outpatients: A randomized trial. <i>European Journal of Cardiovascular Nursing.</i> 2011;10(3):174-9.	Wrong or missing intervention
Brensilver M, Heinzerling KG, Swanson AN, Shoptaw SJ. A retrospective analysis of two randomized trials of bupropion for methamphetamine dependence: suggested guidelines for treatment discontinuation/augmentation. <i>Drug & Alcohol Dependence.</i> 2012;125(1-2):169-72.	Wrong population
Brentwood Biomedical Research I. Nicotine Receptor Density & Response to Nicotine Patch: Pt 2 Extended Treatment. https://ClinicalTrials.gov/show/NCT02676375 ; 2014.	Ongoing study
Brody AL, Zorick T, Hubert R, Hellemann GS, Balali S, Kawasaki SS, et al. Combination Extended Smoking Cessation Treatment Plus Home Visits for Smokers With Schizophrenia: A Randomized Controlled Trial. <i>Nicotine & Tobacco Research.</i> 2016;3:3.	Participants received same drug
Brose LS, Hitchman SC, Brown J, West R, McNeill A. Is the use of electronic cigarettes while smoking associated with smoking cessation attempts, cessation and reduced cigarette consumption? A survey with a 1-year follow-up. <i>Addiction.</i> 2015;110(7):1160-8.	Wrong study design
Brown J, Beard E, Kotz D, Michie S, West R. Real-world effectiveness of e-cigarettes when used to aid smoking cessation: a cross-sectional population study. <i>Addiction (Abingdon, England).</i> 2014;109(9):1531-40.	Wrong study design
Brown J, Beard E, Kotz D, Michie S, West R. Real-world effectiveness of e-cigarettes when used to aid smoking cessation: a cross-sectional population study. <i>Addiction.</i> 2014;109(9):1531-40.	Wrong study design
Brown U, National Institute on Drug A. Nicotine and Sensorimotor Replacement for Smoking in Smokers With Schizophrenia. https://ClinicalTrials.gov/show/NCT01213524 ; 2005.	Ongoing study
Browne J, Aslam N, Jones E, Flint B, Dzingai J, Fabris G, et al. Stop smoking as treatment for COPD: Quitinterventions of higher intensity and duration are required. <i>Thorax.</i> 2011;66:A117-A8.	Wrong study design

Buchanan LM, El-Banna M, White A, Moses S, Siedlik C, Wood M. An exploratory study of multicomponent treatment intervention for tobacco dependency. <i>Journal of Nursing Scholarship</i> . 2004;36(4):324-30.	Participants received same drug
Buck D, Morgan A. Smoking and quitting with the aid of nicotine replacement therapies in the English adult population. Results from the Health Education Monitoring Survey 1995. <i>European Journal of Public Health</i> . 2001;11(2):211-7.	Wrong study design
Budney AJ. Clinical findings in cannabis research: Implications for practice: Clinical approaches to co-occurring cannabis and tobacco use. <i>American Journal on Addictions</i> . 2017;26 (3):233.	Ongoing study
Bullen C, Verbiest M, Galea-Singer S, Kurdziel T, Laking G, Newcombe D, et al. The effectiveness and safety of combining varenicline with nicotine e-cigarettes for smoking cessation in people with mental illnesses and addictions: study protocol for a randomised-controlled trial. <i>BMC Public Health</i> . 2018;18(1):596.	Ongoing study
Buller DB, Halperin A, Severson HH, Borland R, Slater MD, Bettinghaus EP, et al. Effect of nicotine replacement therapy on quitting by young adults in a trial comparing cessation services. <i>Journal of Public Health Management & Practice</i> . 2014;20(2):E7-E15.	Wrong study design
Burling TA, Burling AS, Latini D. A controlled smoking cessation trial for substance-dependent inpatients. <i>Journal of Consulting and Clinical Psychology</i> . 2001;69(2):295-304.	Participants received same drug
Bushnell FKL, Forbes B, Goffaux J, Dietrich M, Wells N. Smoking cessation in military personnel. <i>Military Medicine</i> . 1997;162(11):715-9.	Participants received same drug
Butler H, Brown U. Mood and Anti-craving Effects of Varenicline in Psychiatric Inpatients. https://ClinicalTrials.gov/show/NCT00525928 ; 2007.	Study withdrawn
Butler H, Brown U. Study of Mood Effects of Varenicline (Chantix) in Depressed Outpatient Smokers. https://ClinicalTrials.gov/show/NCT00525837 ; 2007.	Wrong study design
Butler H, National Cancer I. Varenicline or Nicotine Patch and Nicotine Gum in Helping Smokers in a Methadone Treatment Program Stop Smoking. https://ClinicalTrials.gov/show/NCT00790569 ; 2008.	Ongoing study
Cabezas C, Advani M, Puente D, Rodriguez-Blanco T, Martin C. Effectiveness of a stepped primary care smoking cessation intervention: cluster randomized clinical trial (ISTAPS study)2017; 106(9):[1696-706 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/996/CN-01260996/frame.html .	Wrong study design
Calabro KS, Marani S, Yost T, Segura J, Mullin-Jones M, Nelson S, et al. Project SUCCESS: Results from a Randomized Controlled Trial. <i>ISRN Public Health</i> . 2012;2012:Article ID 913713.	Participants received same drug
Calhoun PS, Datta S, Olsen M, Smith VA, Moore SD, Hair LP, et al. Comparative effectiveness of an Internet-based smoking cessation intervention versus clinic-based specialty care for veterans. <i>Journal of Substance Abuse Treatment</i> . 2016;69:19-27.	Participants received same drug
Calhoun PS, Wilson SM, Hicks TA, Thomas SP, Dedert EA, Hair LP, et al. Racial and Sociodemographic Disparities in Internet Access and eHealth Intervention Utilization Among Veteran Smokers. <i>Journal of Racial and Ethnic Health Disparities</i> . 2017;4(5):846-53.	Linked to excluded study
Caponnetto P, Auditore R, Russo C, Cappello GC, Polosa R. Impact of an electronic cigarette on smoking reduction and cessation in schizophrenic smokers: A prospective 12-month pilot study. <i>International Journal of Environmental Research and Public Health</i> . 2013;10(2):446-61.	Wrong study design
Caponnetto P, Polosa R, Auditore R, Minutolo G, Signorelli M, Maglia M, et al. Smoking cessation and reduction in schizophrenia (SCARIS) with e-cigarette: study protocol for a randomized control trial. <i>Trials [Electronic Resource]</i> . 2014;15:88.	Ongoing study

Carandang R, Sims J, Rordorf G. Nicotine replacement therapy is not associated with increased vasospasm in aneurysmal subarachnoid hemorrhage. <i>Neurocritical Care</i> . 2010;13:S197.	Wrong study design
Carandang RA, Barton B, Rordorf GA, Ogilvy CS, Sims JR. Nicotine replacement therapy after subarachnoid hemorrhage is not associated with increased vasospasm. <i>Stroke</i> . 2011;42(11):3080-6.	Wrong study design
Carmody TP, Delucchi K, Simon JA, Duncan CL, Solkowitz SN, Huggins J, et al. Expectancies Regarding the Interaction Between Smoking and Substance Use in Alcohol-Dependent Smokers in Early Recovery. <i>Psychology of Addictive Behaviors</i> . 2012;26(2):358-63.	Linked to excluded study
Carpenter MJ, Hughes JR, Keely JP. Effect of smoking reduction on later cessation: a pilot experimental study. <i>Nicotine & Tobacco Research</i> . 2003;5(2):155-62.	Participants received same drug
Carroll AJ, Veluz-Wilkins AK, Blazekovic S, Kalhan R, Leone FT, Wileyto EP, et al. Cancer-related disease factors and smoking cessation treatment: Analysis of an ongoing clinical trial. <i>Psycho-Oncology</i> . 2018;27(2):471-6.	Ongoing study
Carroll DM, Wagener TL, Stephens LD, Brame LS, Thompson DM, Beebe LA. The relationship between nicotine metabolism and nicotine and carcinogen exposure among American Indian commercial cigarette smokers and electronic nicotine delivery system users. <i>Addictive Behaviors</i> . 2018;92:58-63.	Wrong comparator
Casalino L, Griffo R, Barbara C, Zappulla T, Clavario P. Efficacy of varenicline therapy in the "board the bus and quit" smoking cessation program: Six-month results. <i>European Heart Journal</i> . 2011;32:222.	Wrong study design
Case Comprehensive Cancer C, American Cancer Society I, Alere W, North Carolina Tobacco Q. Enhancing Quitline Services for African American Smokers. https://ClinicalTrials.gov/show/NCT03064971 ; 2017.	Participants received same drug
Castaldelli-Maia JM, Loreto AR, Guimaraes-Pereira BBS, Carvalho CFC, Gil F, Frallonardo FP, et al. Smoking cessation treatment outcomes among people with and without mental and substance use disorders: An observational real-world study. <i>European Psychiatry: the Journal of the Association of European Psychiatrists</i> . 2018;52:22-8.	Wrong study design
Castellet JMC, Doniga IF, Laborda MQ, Torrecilla BS, Agudo LS. Design and early results of a proactive telephone-based smoking cessation treatment compared with face-to-face group sessions. <i>Medicina Clinica</i> . 2007;128(7):247-50.	Participants received same drug
Castro A, Valerio J, Correia A, Guimaraes M, Pascoal I. Smoking cessation-What determines adherence to pharmacological treatment. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2011;38(no pagination).	Cannot locate full text
Cather C, Dyer MA, Burrell HA, Hoepfner B, Goff DC, Evins AE. An open trial of relapse prevention therapy for smokers with schizophrenia. <i>Journal of Dual Diagnosis</i> . 2013;9(1):87-93.	Wrong study design
Center HLMC, Research I, James, Esther King Biomedical Research P, Pfizer. Translating Extinction Research to Improve Pharmacotherapy for Tobacco Dependence. https://ClinicalTrials.gov/show/NCT02202499 ; 2014.	Ongoing study
Center HLMC, Research I, National Cancer I. Smoking Relapse-Prevention Intervention for Cancer Patients. https://ClinicalTrials.gov/show/NCT01630161 ; 2012.	Ongoing study
Center MDAC, National Cancer I. Project Impact 2: A Culturally Tailored Smoking Cessation Intervention for Latino Smokers. https://ClinicalTrials.gov/show/NCT02596711 ; 2015.	Participants received same drug
Center MDAC, National Cancer I. Smoking Cessation for Low-Income Adults. https://ClinicalTrials.gov/show/NCT00948129 ; 2010.	Ongoing study

Center MDAC, National Cancer I. Smoking Cessation Intervention During Low Dose CT (LDCT) Screening for Lung Cancer. https://ClinicalTrials.gov/show/NCT03059940 ; 2017.	Participants received same drug
Center MDAC, National Cancer I. Smoking Cessation Treatment for Head & Neck Cancer Patients. https://ClinicalTrials.gov/show/NCT01098955 ; 2010.	Participants received same drug
Center MDAC, National Institute of Mental H. Smoking Cessation for Depressed Smokers. https://ClinicalTrials.gov/show/NCT00494728 ; 2007.	Participants received same drug
Center MDAC, National Institute on Drug A. Behavioral Activation Therapy for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02697227 ; 2016.	Participants received same drug
Center MDAC, National Institute on Minority H, Health D, Department of H, Human S, National Institutes of H. Determinants of Smoking Cessation Among Latinos. https://ClinicalTrials.gov/show/NCT01008748 ; 2009.	Participants received same drug
Center MDAC, Pharmaceuticals NAL, Pfizer. Reward Sensitivity and Pharmacotherapy for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02162849 ; 2015.	Ongoing study
Center MDAC, Pharmaceuticals NAL, Pfizer. Varenicline and Combined Nicotine Replacement Therapy (NRT) for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02271919 ; 2015.	Ongoing study
Centre for A, Mental H, Canadian Cancer Society Research I, Ottawa Heart Institute Research C. Personalized Dosing of Nicotine Replacement (NRT to Effect). https://ClinicalTrials.gov/show/NCT03000387 ; 2017.	Ongoing study
Centre for A, Mental H, Canadian Institutes of Health R. Effect of Varenicline on Reactivity to Smoking and Drinking Cues. https://ClinicalTrials.gov/show/NCT00873535 ; 2008.	Ongoing study
Centre for A, Mental H, Canadian Tobacco Control Research I. Testing a Full Substitution Therapy Approach As Treatment of Tobacco Dependence. https://ClinicalTrials.gov/show/NCT00390923 ; 2007.	Study terminated
Centre for A, Mental H, Global Research Awards for Nicotine D. The MATCH (Medication Aids for Tobacco Cessation and Health) Study. https://ClinicalTrials.gov/show/NCT02146911 ; 2014.	Ongoing study
Centre for A, Mental H, Ontario Lung A. Varenicline in Drug Treatment. https://ClinicalTrials.gov/show/NCT01286584 ; 2011.	Ongoing study
Centre for A, Mental H, Ontario Ministry of H, Long Term C, Pfizer. Stop Smoking Therapy for Ontario Patients. https://ClinicalTrials.gov/show/NCT00352781 ; 2006.	Wrong study design
Centre for A, Mental H, Ontario Ministry of H, Long Term C. Phase I Study: Stop Smoking Therapy for Ontario Patients (STOP). https://ClinicalTrials.gov/show/NCT00356993 ; 2005.	Wrong study design
Centre for A, Mental H, Ontario Ministry of H, Long Term C. STOP Study: Effectiveness of Zyban in a Clinical Population. https://ClinicalTrials.gov/show/NCT01015170 ; 2009.	Wrong study design
Centre for A, Mental H, Pfizer. Effect of Varenicline on Cognitive Function in Cigarette Smokers With Schizophrenia. https://ClinicalTrials.gov/show/NCT01093365 ; 2010.	Ongoing study
Centre for A, Mental H, Pfizer. Effects of Varenicline on Plasticity in Schizophrenia. https://ClinicalTrials.gov/show/NCT01934023 ; 2013.	Ongoing study

Centre for A, Mental H, Pfizer. Effects of Varenicline on Smoking Lapse in Smokers With and Without Schizophrenia. https://ClinicalTrials.gov/show/NCT01850953 ; 2013.	Ongoing study
Centre for A, Mental H. Efficacy and Neural Correlates of Personalized Treatment With Transdermal Nicotine Replacement. https://ClinicalTrials.gov/show/NCT02439944 ; 2015.	Ongoing study
Chakravorty BJ. A product substitution approach to adolescent smokeless tobacco cessation [thesis]. Dissertation Abstracts International. 1992;53(6-B):2808-9.	Wrong population
Chan SS, Leung DY, Abdullah AS, Lo SS, Yip AW, Kok WM, et al. Smoking-cessation and adherence intervention among Chinese patients with erectile dysfunction. American Journal of Preventive Medicine. 2010;39(3):251-8.	Participants received same drug
Chang E, Fung LC, Li CS, Lin TC, Tam L, Tang C, et al. Offering acupuncture as an adjunct for tobacco cessation: a community clinic experience. Health promotion practice. 2013;14(5 Supplement):80S-7S.	Wrong comparator
Chen J, Chen Y, Chen P, Liu Z, Luo H, Cai S. Effectiveness of individual counseling for smoking cessation in smokers with chronic obstructive pulmonary disease and asymptomatic smokers. Experimental and Therapeutic Medicine. 2014;7(3):716-20.	Wrong or missing intervention
Cheung YT, Leung JP, Cheung CK, Li WH, Wang MP, Lam TH. Motivating smokers at outdoor public smoking hotspots to have a quit attempt with a nicotine replacement therapy sample: study protocol for a randomized controlled trial. 2016;17(1):355.	Ongoing study
Choi K, Forster J. Characteristics Associated With Awareness, Perceptions, and Use of Electronic Nicotine Delivery Systems Among Young US Midwestern Adults. American Journal of Public Health. 2013;103(3):556-61.	Wrong study design
Choi K, Forster JL. Authors' Response. American Journal of Preventive Medicine. 2014;46(6):e58-e9.	Linked to excluded study
Choi K, Forster JL. Beliefs and experimentation with electronic cigarettes: a prospective analysis among young adults. 2014;46(2):175-8.	Wrong study design
Choi W, Beebe L, Nazir N, Kaur B, Hopkins M, Talawyma M, et al. All Nations Breath of Life: a Randomized Trial of Smoking Cessation for American Indians 2017; 51(5 Supplement3):[743-51 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/284/CN-01245284/frame.html .	Wrong study design
Choi WS, Beebe LA, Nazir N, Kaur B, Hopkins M, Talawyma M, et al. All Nations Breath of Life: a Randomized Trial of Smoking Cessation for American Indians. American Journal of Preventive Medicine. 2016;51(5):743-51.	Wrong study design
Choi WS, Faseru B, Beebe LA, Greiner AK, Yeh H-W, Shireman TI, et al. Culturally-Tailored Smoking Cessation for American Indians: Study protocol for a randomized controlled trial. Trials. 2011;12.	Linked to excluded study
Chorti M, Poulianiti K, Jamurtas A, Kostikas K, Tzatzarakis M, Vynias D, et al. Effects of active and passive electronic and tobacco cigarette smoking on lung function. Toxicology Letters. 2012;211:S64-S.	Wrong study design
Chou KJ, Chen HK, Hung CH, Chen TT, Chen CM, Wu BJ. Readiness to quit as a predictor for outcomes of smoking-reduction programme with transdermal nicotine patch or bupropion in a sample of 308 patients with schizophrenia. European Archives of Psychiatry & Clinical Neuroscience. 2015;265(3):249-57.	Awaiting classification
Chou KR, Chen R, Lee JF, Ku CH, Lu RB. The effectiveness of nicotine-patch therapy for smoking cessation in patients with schizophrenia. International Journal of Nursing Studies. 2004;41(3):321-30.	Wrong comparator

Chouinard M-C, Robichaud-Ekstrand S. The effectiveness of a nursing inpatient smoking cessation program in individuals with cardiovascular disease. Nursing research. 2005;54(4):243-54.	Participants received same drug
Christenhusz L, Pieterse M, Seydel E, van der Palen J. Prospective determinants of smoking cessation in COPD patients within a high intensity or a brief counseling intervention. Patient Education & Counseling. 2007;66(2):162-6.	Participants received same drug
Christensen T, Welsh E, Faseru B. Profile of e-cigarette use and its relationship with cigarette quit attempts and abstinence in Kansas adults. Preventive Medicine. 2014;69:90-4.	Wrong study design
Christiansen BA, Carbin J, TerBeek E, Fiore MC. Helping Smokers with Severe Mental Illness Who Do Not Want to Quit. Substance Use & Misuse. 2018;53(6):949-62.	Participants received same drug
Chui CY, Thomas D, Taylor S, Bonevski B, Abramson MJ, Paul E, et al. Factors associated with nicotine replacement therapy use among hospitalised smokers. Drug & Alcohol Review. 2018;37(4):514-9.	Linked to excluded study
Cigrang JA, Severson HH, Peterson AL. Pilot evaluation of a population-based health intervention for reducing use of smokeless tobacco. Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco. 2002;4(1):127-31.	Wrong or missing intervention
Claudio Mendes P, University of Sao P. Effect of Smoking Cessation on Clinical and Microbiological Outcomes of the Non-surgical Periodontal Therapy. https://ClinicalTrials.gov/show/NCT02744417 ; 2010.	Ongoing study
Cohen SJ, Christen AG, Katz BP. Counseling medical and dental patients about cigarette smoking: The impact of nicotine gum and chart reminders. American Journal of Public Health. 1987;77(3):313-6.	Wrong study design
Cohen SJ, Stookey GK, Katz BP, Drook CA, Christen AG. HELPING SMOKERS QUIT - A RANDOMIZED CONTROLLED TRIAL WITH PRIVATE-PRACTICE DENTISTS. Journal of the American Dental Association. 1989;118(1):41-5.	Wrong study design
Cohen SJ, Stookey GK, Katz BP, Drook CA, Smith DM. ENCOURAGING PRIMARY CARE PHYSICIANS TO HELP SMOKERS QUIT - A RANDOMIZED, CONTROLLED TRIAL. Annals of Internal Medicine. 1989;110(8):648-52.	Awaiting classification
Cohrs S, Rodenbeck A, Riemann D, Szagun B, Jaehne A, Brinkmeyer J, et al. Impaired sleep quality and sleep duration in smokers-results from the German Multicenter Study on Nicotine Dependence. Addiction Biology. 2014;19(3):486-96.	Wrong or missing intervention
Colby H, Sprick J, Pham G, Cooke W, Fogt D, Rickards C. Cerebral blood flow regulation following inhalation of nicotine via electronic cigarettes. FASEB Journal Conference: Experimental Biology. 2015;29(no pagination).	Cannot locate full text
Coleman T, Cooper S, Thornton JG, Grainge MJ, Watts K, Britton J, et al. A Randomized Trial of Nicotine-Replacement Therapy Patches in Pregnancy. New England Journal of Medicine. 2012;366(9):808-18.	Wrong population
Coleman T, McEwen A, Bauld L, Ferguson J, Lorgelly P, Lewis S. Protocol for the Proactive Or Reactive Telephone Smoking CeSsation Support (PORTSSS) trial. Trials. 2009;10.	Linked to excluded study
Collins B, Lepore S. Babies Living Safe & Smokefree: randomized controlled trial of a multilevel multimodal behavioral intervention to reduce low-income children's tobacco smoke exposure 2017; 17(1):[249 p.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/679/CN-01410679/frame.html .	Participants received same drug

Colvin A, Riege V, Xie H. Prescribing patterns and access to tobacco cessation therapy using a multipractice program and pharmacy-driven clinical practice guideline. <i>Journal of the American Pharmacists Association</i> . 2011;51 (5):640.	Cannot locate full text
Company RJRV, Celerion, Company RAIS. Comparison of Nicotine Plasma Concentrations and Subjective Effects for Three Menthol Electronic Cigarettes vs Combustible Menthol Cigarettes and Nicotine Gum. https://ClinicalTrials.gov/show/NCT02664012 ; 2016.	Ongoing study
Cook JW, Collins LM, Fiore MC, Smith SS, Fraser D, Bolt DM, et al. Comparative effectiveness of motivation phase intervention components for use with smokers unwilling to quit: A factorial screening experiment. <i>Addiction</i> . 2015:No Pagination Specified.	Wrong study design
Cooper TV, Taylor T, Murray A, DeBon MW, Weg MWV, Klesges RC, et al. Differences between intermittent and light daily smokers in a population of US military recruits. <i>Nicotine & Tobacco Research</i> . 2010;12(5):465-73.	Linked to excluded study
Cortada EB, Munoz J, Pinilla CF, Garcia-Aranda VL, Torrell JMR, de Burgoa VS, et al. Profile of the patient who wants to quit smoking seen in the Primary Care and Specialized consultations in Spain. <i>Revista Clinica Espanola</i> . 2009;209(4):160-7.	Wrong study design
Courtney K, Ray L. The effects of varenicline and naltrexone, alone and in combination, on neural responses to alcohol cues in heavy drinking smokers 2013; 37:[248a p.]. Available from: http://onlinelibrary.wiley.com/doi/10.1002/1471-2575.tb027176 .	Cannot locate full text
Cox LS, Cupertino A-P, Mussulman LM, Nazir N, Greiner KA, Mahnken JD, et al. Design and baseline characteristics from the KAN-QUIT disease management intervention for rural smokers in primary care. <i>Preventive Medicine</i> . 2008;47(2):200-5.	Linked to excluded study
Creighton U. e-Cigarettes Versus NRT Gum for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01925781 ; 2013.	Study terminated
Croghan IT, Hurt RD, Dakhil SR, Croghan GA, Sloan JA, Novotny PJ, et al. Randomized comparison of a nicotine inhaler and bupropion for smoking cessation and relapse prevention. <i>Mayo Clinic Proceedings</i> . 2007;82(2):186-95.	Wrong study design
Cservenka A, Courtney KE, Ghahremani DG, Ray LA. Effect of varenicline, naltrexone, and their combination, on alcohol cue-related ventral striatum response in heavy drinking smokers. <i>Alcoholism: Clinical and Experimental Research</i> . 2016;40:263A.	Cannot locate full text
Cummings SR. An evaluation of a behavioral change intervention for smokeless tobacco use [thesis]. <i>Dissertation Abstracts International</i> . 1996;56(12-B):6692.	Wrong or missing intervention
Cupertino AP, Mahnken JD, Richter K, Cox LS, Casey G, Resnicow K, et al. Long-term engagement in smoking cessation counseling among rural smokers. <i>Journal of Health Care for the Poor and Underserved</i> . 2007;18(4):39-51.	Linked to excluded study
Currie SR, Karltn J, Lussier D, De Denus E, Brown D, El-Guebaly N. Outcome from a community-based smoking cessation program for persons with serious mental illness. <i>Community Mental Health Journal</i> . 2008;44(3):187-94.	Participants received same drug
Da Penha Sales M, Aguiar F, Oliveira MI. Efficacy of varenicline among smokers enrolled on a public health tobacco control program (HM-TCP) in the real-life setting. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2015;46(no pagination).	Cannot locate full text
Daejeon U. Trial of Treatment Protocol for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02768025 ; 2016.	Participants received same drug
Dahne J, Wahlquist AE, Boatright AS, Garrett-Mayer E, Fleming DO, Davis R, et al. Nicotine replacement therapy sampling via primary care: Methods from a pragmatic cluster randomized clinical trial. <i>Contemporary Clinical Trials</i> . 2018;72:1-7.	Ongoing study

Dalmau R, Castro A, Marin M, Torres F, Valbuena S, Araujo A, et al. Smoking cessation intervention after acute myocardial infarction, predictors of relapse analysed. <i>European Journal of Preventive Cardiology</i> . 2012;1):S24.	Cannot locate full text
Danaher BG, Hart LG, McKay HG, Severson HH. Measuring participant rurality in web-based interventions. <i>BMC Public Health</i> . 2007;7.	Linked to excluded study
Danaher BG, Severson HH, Andrews JA, Tyler MS, Lichtenstein E, Woolley TG, et al. Randomized Controlled Trial of MyLastDip: A Web-Based Smokeless Tobacco Cessation Program for Chewers Ages 14-25. <i>Nicotine & Tobacco Research</i> . 2013;15(9):1502-10.	Wrong or missing intervention
Danaher BG, Severson HH, Crowley R, van Meter N, Tyler MS, Widdop C, et al. Randomized controlled trial examining the adjunctive use of nicotine lozenges with MyLastDip: An eHealth smokeless tobacco cessation intervention. <i>Internet Interventions</i> . 2015;2(1):69-76.	Wrong comparator
Danan ER, Fu SS, Clothier BA, Noorbaloochi S, Hammett PJ, Widome R, et al. The Equity Impact of Proactive Outreach to Smokers: Analysis of a Randomized Trial. <i>American Journal of Preventive Medicine</i> . 2018;55(4):506-16.	Linked to excluded study
Dartmouth-Hitchcock Medical C, National Institute on Drug A. Targeting Tobacco Cessation During Treatment for Cannabis Use Disorders. https://ClinicalTrials.gov/show/NCT01834794 ; 2013.	Participants received same drug
Das S, Hickman N, Prochaska J. Treating Smoking in Adults With Co-occurring Acute Psychiatric and Addictive Disorders 2017; 11(4):[273-9 pp.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/020/CN-01412020/frame.html .	Linked to excluded study
Davidescu L, Jurca R, Ulmeanu R. Value of adding behavioral-cognitive therapy to standard treatment in smoking cessation programme: Results of smoking cessation center oradea on 7 years. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2014;44(no pagination).	Cannot locate full text
Dawkins L, Turner J, Crowe E. Nicotine derived from the electronic cigarette improves time-based prospective memory in abstinent smokers. <i>Psychopharmacology</i> . 2013;227(3):377-84.	Wrong study design
Dawood B, Mulder J, Reynald L. An audit of smoking prevalence and cessation amongst patients admitted to the respiratory unit in peninsula health. <i>Respirology</i> . 2015;20:84.	Wrong study design
De La Garza R, Yoon J, Yammine L, Holst M, Salas R. Preliminary evaluation of the effects of electronic cigarettes versus own cigarette on withdrawal, craving, and smoking severity in tobacco-dependent volunteers. <i>Neuropsychopharmacology</i> . 2016;41:S380.	Wrong comparator
De La Red Gallego H, Gonzalez Silva Y, Montero Carretero T, Delgado De Paz A, Sanchez Anorga MF, Canibano Maroto E, et al. Results of a smoking cessation program in primary care. <i>European Psychiatry</i> . 2016;33:S297-S8.	Wrong or missing intervention
de Moor JS, Puleo E, Ford JS, Greenberg M, Hodgson DC, Tyc VL, et al. Disseminating a smoking cessation intervention to childhood and young adult cancer survivors: baseline characteristics and study design of the partnership for health-2 study. <i>Bmc Cancer</i> . 2011;11.	Linked to excluded study
Decker KP, Peglow S. Nicotine replacement treatment in patients with comorbid medical or psychiatric problems during substance use rehabilitation. <i>American Journal on Addictions</i> . 2013;22 (3):315-6.	Wrong comparator
Dockrell M, Morrison R, Bauld L, McNeill A. E-Cigarettes: Prevalence and Attitudes in Great Britain. <i>Nicotine & Tobacco Research</i> . 2013;15(10):1737-44.	Wrong study design

Donny EC, Jones M. Prolonged exposure to denicotinized cigarettes with or without transdermal nicotine. Drug and Alcohol Dependence. 2009;104(1-2):23-33.	Wrong or missing intervention
Doran C, Stafford J, Shanahan M, Mattick RP. Characteristics of Australian smokers using bupropion and nicotine-replacement therapies. Expert Review of Pharmacoeconomics and Outcomes Research. 2007;7(1):41-7.	Wrong study design
Doran CM, Valenti L, Robinson M, Britt H, Mattick RP. Smoking status of Australian general practice patients and their attempts to quit. Addictive Behaviors. 2006;31(5):758-66.	Wrong study design
Douptcheva N, Gmel G, Studer J, Deline S, Etter JF. Use of electronic cigarettes among young Swiss men. Journal of Epidemiology and Community Health. 2013;67(12):1075-6.	Wrong study design
dr. maria m, University of M, Florida International U. Patch Study - Intervention for HIV Positive Smokers. https://ClinicalTrials.gov/show/NCT02982772 ; 2016.	Ongoing study
Drehmer JE, Hipple B, Nabi-Burza E, Ossip DJ, Chang YC, Rigotti NA, et al. Proactive enrollment of parents to tobacco quitlines in pediatric practices is associated with greater quitline use: a cross-sectional study. BMC Public Health. 2016;16.	Wrong study design
Drovandi AD, Teague P-A, Glass BD, Malau-Aduli B. A randomised controlled trial evaluating the effectiveness and tolerability of step-up and step-down varenicline therapy for smoking cessation: A study protocol. Journal Of Smoking Cessation. 2017:No-Specified.	Ongoing study
D'Ruiz CD, Graff DW, Robinson E. Reductions in biomarkers of exposure, impacts on smoking urge and assessment of product use and tolerability in adult smokers following partial or complete substitution of cigarettes with electronic cigarettes. BMC Public Health. 2016;16.	Wrong comparator
Du D, Nides M, Borders J, Selmani A, Wawerczak W. Comparison of nicotine oral soluble film and nicotine lozenge on efficacy in relief of smoking cue-provoked acute craving after a single dose of treatment in low dependence smokers. Psychopharmacology. 2014;231(22):4383-91.	Wrong comparator
Duffy SA, Ronis DL, Titler MG, Blow FC, Jordan N, Thomas PL, et al. Dissemination of the nurse-administered Tobacco Tactics intervention versus usual care in six Trinity community hospitals: study protocol for a comparative effectiveness trial. Trials. 2012;13:125.	Wrong or missing intervention
Duffy SA, Ronis DL, Valenstein M, Lambert MT, Fowler KE, Gregory L, et al. A tailored smoking, alcohol, and depression intervention for head and neck cancer patients. Cancer Epidemiology Biomarkers & Prevention. 2006;15(11):2203-8.	Wrong study design
Duffy SA, Scheumann AL, Fowler KE, Darling-Fisher C, Terrell JE. Perceived Difficulty Quitting Predicts Enrollment in a Smoking-Cessation Program for Patients With Head and Neck Cancer. Oncology Nursing Forum. 2010;37(3):349-56.	Linked to excluded study
Duke U, Intelligent Automation I, National Institute on Drug A. Stage Ib Trial of mSMART for Smoking Cessation Medication Adherence. https://ClinicalTrials.gov/show/NCT02635919 ; 2016.	Participants received same drug
Duke U, Intelligent Automation I, National Institute on Drug A. Stage Ib Trial of mSMART With Varenicline. https://ClinicalTrials.gov/show/NCT03069768 ; 2017.	Participants received same drug
Duke U, National Institute of A, Infectious D. Effects of E-cigarettes (ECIGs) on Pulmonary Inflammation and Behavior in HIV Infected Smokers. https://ClinicalTrials.gov/show/NCT02848586 ; 2016.	Wrong comparator
Duke U, National Institute on Drug A, Pfizer. Adaptive Pharmacotherapy for Tobacco Dependence. https://ClinicalTrials.gov/show/NCT02501265 ; 2017.	Ongoing study
Duke U, National Institute on Drug A, Philip Morris Usa I. Combination Bupropion Varenicline for Smoking Cessation in Male Smokers. https://ClinicalTrials.gov/show/NCT01806779 ; 2013.	Ongoing study

Duke U, National Institute on Drug A. Mechanistic Evaluations of Pre-Cessation Therapies for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01406223 ; 2011.	Ongoing study
Duke U, Philip Morris Usa I. Tailored Smoking Cessation Treatment for LIVE FOR LIFE® Participants. https://ClinicalTrials.gov/show/NCT01560507 ; 2012.	Ongoing study
Duke U, Shire. Attention Deficit Hyperactivity Disorder (ADHD) Smoking Cessation Study. https://ClinicalTrials.gov/show/NCT00736255 ; 2007.	Participants received same drug
Ebbert JO, Little MA, Klesges RC, Bursac Z, Johnson KC, Thomas F, et al. Step Care treatment for smoking cessation. <i>Health Education Research</i> . 2017;32(1):1-11.	Wrong study design
Ebbert JO, Severson HH, Croghan IT, Danaher BG, Schroeder DR. Comparative effectiveness of the nicotine lozenge and tobacco-free snuff for smokeless tobacco reduction. <i>Addictive Behaviors</i> . 2013;38(5):2140-5.	Wrong comparator
Ebrahimi SJA, Ebrahimzadeh L, Bidarpour F, Gharibi F. Efficacy of different methods combined with nicotine patch therapy for smoking cessation. <i>Research Journal of Pharmaceutical, Biological and Chemical Sciences</i> . 2015;6(2):69-74.	Wrong comparator
Edmunds M, Conner H, Jones C, Gorayeb R, Waranch HR. Evaluation of a multicomponent group smoking cessation program. <i>Preventive Medicine</i> . 1991;20(3):404-13.	Wrong comparator
Ellerbeck EF, Mahnken JD, Cupertino AP, Cox LS, Greiner A, Mussulman LM, et al. Effect of Varying Levels of Disease Management on Smoking Cessation A Randomized Trial. <i>Annals of Internal Medicine</i> . 2009;150(7):437-+.	Wrong study design
Emmons KM, Puleo E, Sprunck-Harrild K, Ford J, Ostroff JS, Hodgson D, et al. Partnership for Health-2, A Web-Based Versus Print Smoking Cessation Intervention for Childhood and Young Adult Cancer Survivors: Randomized Comparative Effectiveness Study. <i>Journal of Medical Internet Research</i> . 2013;15(11).	Participants received same drug
Esmat I, Kassim D. Comparative study between transdermal nicotine and melatonin patches on postoperative pain relief after laparoscopic cholecystectomy, a double-blind, placebo-controlled trial 2017; 32(3):[299-307 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/191/CN-01246191/frame.html .	Wrong population
Essilini A, Abadie D, Montastruc F, Fulda V, Gouraud A, Yelehe-Okouma M, et al. Drug-induced panic attacks: Analysis of the French pharmacovigilance database. <i>Drug Safety</i> . 2015;38 (10):1009.	Wrong comparator
Etter JF, Bullen C. A longitudinal study of electronic cigarette users. 2014;39(2):491-4.	Wrong study design
Etter J-F, Bullen C. Electronic cigarette: users profile, utilization, satisfaction and perceived efficacy. <i>Addiction</i> . 2011;106(11):2017-28.	Wrong study design
Etter JF, Eissenberg T. Dependence levels in users of electronic cigarettes, nicotine gums and tobacco cigarettes. <i>Drug and Alcohol Dependence</i> . 2015;147:68-75.	Wrong study design
Euctr IT. Efficacy and safety of smoking cessation with varenicline tartrate in diabetic smokers: a double-blind, placebo-controlled, randomized trial. 2009.	Ongoing study
Faessel H, Ravva P, Williams K. Pharmacokinetics, Safety, and Tolerability of Varenicline in Healthy Adolescent Smokers: A Multicenter, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study. <i>Clinical Therapeutics</i> . 2009;31(1):177-89.	Wrong population
Fagerstrom KO, Hughes JR, Rasmussen T, Callas PW. Randomised trial investigating effect of a novel nicotine delivery device (Eclipse) and a nicotine oral inhaler on smoking behaviour, nicotine and carbon monoxide exposure, and motivation to quit. <i>Tobacco Control</i> . 2000;9(3):327-33.	Wrong study design
Fagerstrom KO, Schneider NG, Lunell E. EFFECTIVENESS OF NICOTINE PATCH AND NICOTINE GUM AS INDIVIDUAL VERSUS COMBINED TREATMENTS FOR TOBACCO WITHDRAWAL SYMPTOMS. <i>Psychopharmacology</i> . 1993;111(3):271-7.	Wrong study design

Farris S, Leyro T, Allan N, Overup C, Schmidt N, Zvolensky M. Distress intolerance during smoking cessation treatment 2017; 85:[33-42 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/979/CN-01215979/frame.html .	Participants received same drug
Farsalinos KE, Tsiapras D, Kyrzopoulos S, Savvopoulou M, Voudris V. Acute effects of using an electronic nicotine-delivery device (electronic cigarette) on myocardial function: Comparison with the effects of regular cigarettes. <i>BMC Cardiovascular Disorders</i> . 2014;14 (no pagination)(78).	Wrong comparator
Faseru B, Ellerbeck EF, Catley D, Gajewski BJ, Scheuermann TS, Shireman TI, et al. Changing the default for tobacco-cessation treatment in an inpatient setting: study protocol of a randomized controlled trial. <i>Trials [Electronic Resource]</i> . 2017;18(1):379.	Participants received same drug
Fatemi SH, Stary JM, Hatsukami DK, Murphy SE. A double-blind placebo-controlled cross over trial of bupropion in smoking reduction in schizophrenia. <i>Schizophrenia Research</i> . 2005;76(2-3):353-6.	Wrong study design
Fda. Drug Safety Communication: FDA revises description of mental health side effects of the stop-smoking medicines Chantix (varenicline) and Zyban (bupropion) to reflect clinical trial findings. www.fda.gov . 2016;[Available from: https://www.fda.gov/Drugs/DrugSafety/ucm532221.htm].	Wrong study design
Fda. Drug Safety Communication: FDA updates label for stop smoking drug Chantix (varenicline) to include potential alcohol interaction, rare risk of seizures, and studies of side effects on mood, behavior, or thinking. www.fda.gov . 2015;[Available from: https://www.fda.gov/Drugs/DrugSafety/ucm436494.htm].	Wrong study design
Fenech MA, Reid C. A smoking cessation intervention for people with chronic hepatitis C: A randomised controlled trial. <i>Hepatology</i> . 2012;56:229A.	Cannot locate full text
Ferguson J, Docherty G, Bauld L, Lewis S, Lorgelly P, Boyd KA, et al. Effect of offering different levels of support and free nicotine replacement therapy via an English national telephone quitline: Randomised controlled trial. <i>BMJ: British Medical Journal</i> . 2012;344(7854):1-13.	Wrong study design
Ferguson J. Effect of offering different levels of support and free nicotine replacement therapy via an English national telephone quitline: randomised controlled trial (vol 344, e1696, 2012). <i>British Medical Journal</i> . 2012;345.	Wrong study design
Ferguson SG, Walters JA, Lu W, Wells GP, Schuz N. Examination of the mechanism of action of two pre-quit pharmacotherapies for smoking cessation. <i>BMC public health</i> . 2015;15:1268.	Ongoing study
Fernandes F, Prado G, Santos U, Lotufo JP, Fukuoka M, Souza E. Anxiety and depression measured by HASD questionnaire are related to failure in 4 week smoking cessation program. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2011;38(no pagination).	Cannot locate full text
Fernandez MA, Vidal AF, Lopez Sampedro P, Garcia Lavandera J. Middle-term effectiveness of a support program for smokers implemented in primary care. [Spanish]. <i>Atencion Primaria</i> . 2002;30(9):541-6.	Wrong comparator
Finger R, Tamm M, Seiffert B, Kuster M, Meyer A, Stolz D. Determinants of change in quality of life after smoking cessation of health care employees. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2011;38(no pagination).	Cannot locate full text
Flocke SA, Lewis S, Seeholzer E, Gill I, Antognoli E, Rose JC, et al. Electronic medical record documentation of tobacco cessation support at eight community safety-net clinics with a high prevalence of tobacco use. <i>J Eval Clin Pract</i> . 2018;20:20.	Wrong study design
Florou A, Eleftheriou K, Patentakis G, Sachlas A, Michailidis D, Gratziou C. Smoking cessation effectiveness in smokers with obstructive respiratory disease. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2011;38(no pagination).	Cannot locate full text
Folan PA, Spatarella A, Jacobsen D, Kohn N, Fardellone C, Talwar A. Self-efficacy intervention and quit rates among participants in a tobacco dependence treatment program. <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2014;189(no pagination).	Cannot locate full text
Fraser D, Borland R, Gartner C. Protocol for a randomised pragmatic policy trial of nicotine products for quitting or long-term substitution in smokers. <i>BMC public health</i> . 2015;15:1026.	Ongoing study

Fraser D, Kobinsky K, Smith SS, Kramer J, Theobald WE, Baker TB. Five population-based interventions for smoking cessation: a MOST trial. <i>Translational Behavioral Medicine</i> . 2014;4(4):382-90.	Wrong study design
French National Institute for H, Medical Research-French National Agency for Research on A, Viral H, Pfizer. Efficacy and Safety of Varenicline Among HIV-infected Patients. https://ClinicalTrials.gov/show/NCT00918307 ; 2009.	Ongoing study
Frikart M, Etienne S, Cornuz J, Zellweger JP. Five-day plan for smoking cessation using group behaviour therapy. <i>Swiss Medical Weekly</i> . 2003;133(3-4):39-43.	Wrong or missing intervention
Frye MA, Ebbert JO, Prince CA, Lineberry TW, Geske JR, Patten CA. A feasibility study of varenicline for smoking cessation in bipolar patients with subsyndromal depression. <i>Journal of Clinical Psychopharmacology</i> . 2013;33(6):821-3.	Participants received same drug
Fu S, Rothman A, Vock D, Lindgren B, Almirall D, Begnaud A, et al. Program for lung cancer screening and tobacco cessation: study protocol of a sequential, multiple assignment, randomized trial 2017; 60:[86-95 pp.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/179/CN-01394179/frame.html .	Wrong study design
Fu S, Ryn M, Sherman S, Burgess D, Noorbaloochi S, Clothier B, et al. Proactive tobacco treatment and population-level cessation: a pragmatic randomized clinical trial 2014; 174(5):[671-7 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/030/CN-00988030/frame.html .	Wrong or missing intervention
Fu SS, van Ryn M, Burgess DJ, Nelson D, Clothier B, Thomas JL, et al. Proactive tobacco treatment for low income smokers: study protocol of a randomized controlled trial. <i>BMC public health</i> . 2014;14:337.	Participants received same drug
Fu SS, Van Ryn M, Nelson D, Burgess DJ, Thomas JL, Saul J, et al. Proactive tobacco treatment offering free nicotine replacement therapy and telephone counselling for socioeconomically disadvantaged smokers: A randomised clinical trial. <i>Thorax</i> . 2016;71(5):446-53.	Participants received same drug
Fucito LM, Zweben A, O'Malley SS. Sleep disturbance in alcohol dependent individuals: The relation with cigarette smoking. <i>Alcoholism: Clinical and Experimental Research</i> . 2015;39:83A.	Cannot locate full text
Fukui M, Suzuki K, Matsunaga T, Oh S, Takamochi K. The Importance of Smoking Cessation on Surgical Outcome in Primary Lung Cancer. <i>Ann Thorac Surg</i> . 2019;02:02.	Wrong or missing intervention
Gallego SS, Villanueva SH, Martinez MN, Martinez AS, Lobato IH, Vehils SM. NICOTINE REPLACEMENT THERAPY ACCEPTANCE IN PSYCHIATRIC INPATIENTS. <i>Revista Rol De Enfermeria</i> . 2018;41(6):422-6.	Awaiting classification
Galvin KT. A nurse-led health education programme for smoking cessation: Evaluation using the theory of planned behaviour. <i>Dissertation Abstracts International Section C: Worldwide</i> . 2018;75(4-C):No-Specified.	Awaiting classification
Gambaryan M, Kalinina A, Grigoryan T, Vigodin V. Implementing smoking cessation strategies in primary health care in Russia: Opportunities and challenges. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2012;40(no pagination).	Cannot locate full text
Gansky SA, Ellison JA, Rudy D, Bergert N, Letendre MA, Nelson L, et al. Cluster-randomized controlled trial of an athletic trainer-directed spit (smokeless) tobacco intervention for collegiate baseball athletes: Results after 1 year. <i>Journal of Athletic Training</i> . 2005;40(2):76-87.	Wrong or missing intervention
Garcia FIS, Derefinko KJ, Bursac Z, Hand S, Klesges RC. Planning a Change Easily (PACE): A randomized controlled trial for smokers who are not ready to quit. <i>Contemporary Clinical Trials</i> . 2018;68:14-22.	Ongoing study

Garvey AJ, Kalman D, Hoskinson RA, Kinnunen T, Armour CD, Copp S, et al. Effects of extended-duration counseling vs. shorter-duration counseling after 1.5 years of follow-up (POS4-49). Society for Research on Nicotine and Tobacco 18th Annual Meeting March 13-16, 2012, Houston, Texas 2012. 2012:139.	Participants received same drug
Gehricke J-G, Hong N, Whalen CK, Steinhoff K, Wigal TL. Effects of transdermal nicotine on symptoms, moods, and cardiovascular activity in the everyday lives of smokers and nonsmokers with attention-deficit/hyperactivity disorder. <i>Psychology of Addictive Behaviors</i> . 2009;23(4):644-55.	Wrong study design
Gehricke J-G, Whalen CK, Jamner LD, Wigal TL, Steinhoff K. The reinforcing effects of nicotine and stimulant medication in the everyday lives of adult smokers with ADHD: A preliminary examination. <i>Nicotine & Tobacco Research</i> . 2006;8(1):37-47.	Wrong study design
Gelderse Vallei H, Deventer Z, Wageningen U. Nicotine Replacement Therapy in the Intensive Care Unit. https://ClinicalTrials.gov/show/NCT01362959 ; 2012.	Ongoing study
Geller AC, Borrelli B, Davine JA, Burtner J, Heeren TC, Rees VW, et al. Factors Associated with Recent Use of Nicotine Replacement Therapy Among Multiethnic Smokers Residing in Public Housing. <i>Journal Of Smoking Cessation</i> . 2018;13(3):162-70.	Wrong study design
General Hospital of K, University of T. Smoking Cessation Intervention in Respiratory Inpatients. https://ClinicalTrials.gov/show/NCT02922387 ; 2012.	Ongoing study
George TP, Ziedonis DM, Feingold A, Pepper WT, Satterburg CA, Winkel J, et al. Nicotine transdermal patch and atypical antipsychotic medications for smoking cessation in schizophrenia. <i>American Journal of Psychiatry</i> . 2000;157(11):1835-42.	Participants received same drug
Georgetown U, Albert Einstein College of Medicine I, University of M. Positively Smoke Free on the Web (PSFW) for Smokers Living With HIV. https://ClinicalTrials.gov/show/NCT01570595 ; 2011.	Participants received same drug
Gershon AS, Campitelli MA, Hawken S, Victor JC, Sproule B, Kurdyak P, et al. The risk of cardiovascular and neuropsychiatric adverse events following varenicline use for tobacco cessation in Ontario, Canada: A self-controlled study. <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2015;191(no pagination).	Wrong study design
Ghiasi F, Farhang S, Farnam A, Safikhlanlou S. The short term effect of nicotine abstinence on visuospatial working memory in smoking patients with schizophrenia. <i>Nordic Journal of Psychiatry</i> . 2013;67(2):104-8.	Wrong comparator
Gilbody S, Peckham E, Man MS, Mitchell N, Li J, Becque T, et al. Bespoke smoking cessation for people with severe mental ill health (SCIMITAR): a pilot randomised controlled trial. <i>The Lancet Psychiatry</i> . 2015;2(5):395-402.	Wrong study design
Gillies MA, McKenzie CA, Whiteley C, Beale RJ, Tibby SM. Safety of nicotine replacement therapy in critically ill smokers: A retrospective cohort study. <i>Intensive Care Medicine</i> . 2012;38(10):1683-8.	Wrong comparator
Giovenco DP, Lewis MJ, Delnevo CD. Factors Associated with E-cigarette Use A National Population Survey of Current and Former Smokers. <i>American Journal of Preventive Medicine</i> . 2014;47(4):476-80.	Wrong study design
GlaxoSmithKline. Comparison of Pharmacokinetic Profiles of Two Nicotine Gum Formulations. https://ClinicalTrials.gov/show/NCT01847443 ; 2013.	Wrong study design
GlaxoSmithKline. Nicotine Lozenge Bioequivalence Study. https://ClinicalTrials.gov/show/NCT01536704 ; 2011.	Wrong study design
GlaxoSmithKline. Relapse Prevention Study in Newly Abstinent Smokers. https://ClinicalTrials.gov/show/NCT00793468 ; 2008.	Study withdrawn
Glostrup University Hospital C, The Danish Rheumatism A, The Novo Nordic F, Region Capital D. REU-stop - Effect of Intensive Smoking Cessation Intervention on Smoking Cessation and Disease Activity in Patients With Rheumatoid Arthritis. https://ClinicalTrials.gov/show/NCT02901886 ; 2016.	Ongoing study
Glover ED, Wang MQ, Glover PN. Development of a high school smokeless tobacco cessation manual. <i>Health Values</i> . 1994;18(2):28-33.	Wrong population

Goldade K, Des Jarlais D, Everson-Rose SA, Guo H, Thomas J, Gelberg L, et al. Knowing Quitters Predicts Smoking Cessation in a Homeless Population. American Journal of Health Behavior. 2013;37(4):517-24.	Linked to excluded study
Goldade K, Whembolua G, Thomas J, Eischen S, Guo H, Connett J, et al. Designing a smoking cessation intervention for the unique needs of homeless persons: a community-based randomized clinical trial 2017; 8(6):[744-54 pp.].	Participants received same drug
Goniewicz ML, Zielinska-Danch W. Electronic Cigarette Use Among Teenagers and Young Adults in Poland. Pediatrics. 2012;130(4):E879-E85.	Wrong study design
Gonzalez FJC, Castillo SG, Ruiz JC, Molina AIT, Crespo MG, Martinez Garcia AJ, et al. Continuous abstinence rates at 3, 6, 9 and 12 months for males in a tobacco cessation unit in Albacete University Hospital during 2010. European Respiratory Journal Conference: European Respiratory Society Annual Congress. 2014;44(no pagination).	Cannot locate full text
Gonzalez FJC, Garcia AM, Crespo MG, Calderon JP, Cano AM, Molina AIT, et al. Differences in results of two follow-up periods in a tobacco cessation unit in albacete university hospital. Chest Conference: CHEST World Congress. 2014;145(3 MEETING ABSTRACT).	Cannot locate full text
Gonzalez FJC, Ruiz JC, Castillo SG, Molina AIT, Crespo MG, Cano AM, et al. Results in a tobacco cessation unit in Albacete University Hospital during 2010. European Respiratory Journal Conference: European Respiratory Society Annual Congress. 2014;44(no pagination).	Cannot locate full text
Gonzalez FJC, Ruiz JC, Castillo SG, Tornero Molina AI, Crespo MG, Garcia AJM, et al. Continuous abstinence rates at 3, 6, 9, and 12 months for males, females, and overall sample in a tobacco cessation unit in Albacete University Hospital during 2010. Chest Conference: CHEST. 2014;146(4 MEETING ABSTRACT).	Cannot locate full text
Gonzalez Roz A, Secades Villa R, Weidberg S. Evaluating nicotine dependence levels in e-cigarette users. Adicciones. 2017;29(2):136-8.	Wrong comparator
Goodney P, Hoel A, Beck A, Tan TW, Hallett JJ, Newhall K, et al. A pilot randomized trial of a brief smoking cessation intervention for patients with peripheral arterial disease: the vascular physicians offer and report (VAPOR) trial 2017; 63(6 suppl. 1):[204s p.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/266/CN-01267266/frame.html .	Participants received same drug
Goodney P, Spangler E, Newhall K, Brooke B, Schanzer A, Tan TW, et al. Feasibility and pilot efficacy of a brief smoking cessation intervention delivered by vascular surgeons in the Vascular Physician Offer and Report (VAPOR) Trial 2017; Date of Publication: August 22. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/597/CN-01330597/frame.html .	Participants received same drug
Goodney PP, Spangler EL, Newhall K, Brooke BS, Schanzer A, Tan TW, et al. Feasibility and pilot efficacy of a brief smoking cessation intervention delivered by vascular surgeons in the Vascular Physician Offer and Report (VAPOR) Trial. Journal of Vascular Surgery. 2017;09:09.	Participants received same drug
Gordon JS, Akers L, Severson HH, Danaher BG, Boles SM. Successful participant recruitment strategies for an online smokeless tobacco cessation program. Nicotine & Tobacco Research. 2006;8:S35-S41.	Linked to excluded study
Gowin J, Vatsalya V, Momenan R, Schwandt M, Bartlett S, Heilig M, et al. Attenuation of amygdala response to alcohol cues by varenicline in heavy drinkers: Association with self-administration 2014; 38:[155a p.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/523/CN-01064523/frame.html .	Cannot locate full text
Gowin J, Vatsalya V, Westman J, Schwandt M, Momenan R, Hommer D, et al. Varenicline administration diminishes amygdala response and self-reported feelings of acute effects of alcohol in heavy drinkers 2014; 39:[S274 p.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/372/CN-01066372/frame.html .	Cannot locate full text

Graham AL, Burke MV, Jacobs MA, Cha S, Croghan IT, Schroeder DR, et al. An integrated digital/clinical approach to smoking cessation in lung cancer screening: study protocol for a randomized controlled trial. <i>Trials</i> . 2017;18.	Wrong or missing intervention
Grana RA, Popova L, Ling PM. A longitudinal analysis of electronic cigarette use and smoking cessation. 2014;174(5):812-3.	Wrong study design
Greaves C. Smoking cessation trial may be missing the point. <i>Thorax</i> . 2008;63(3):291-.	Linked to excluded study
Gross J, Stitzer ML, Maldonado J. NICOTINE REPLACEMENT - EFFECTS ON POSTCESSATION WEIGHT-GAIN. <i>Journal of Consulting and Clinical Psychology</i> . 1989;57(1):87-92.	Wrong population
Group Health C, University of M, University of California D. My Mobile Advice Program: A Randomized Pilot Feasibility Study. https://ClinicalTrials.gov/show/NCT02136498 ; 2014.	Participants received same drug
Groupe Hospitalier Paris Saint J. Study of the Immediate Cardiovascular Effects of Electronic Cigarette in Subjects With Cardiovascular Disease. https://ClinicalTrials.gov/show/NCT02777515 ; 2015.	Ongoing study
Grundey J, Amu R, Ambrus GG, Batsikadze G, Paulus W, Nitsche MA. Double dissociation of working memory and attentional processes in smokers and non-smokers with and without nicotine. <i>Psychopharmacology</i> . 2015;232(14):2491-501.	Wrong study design
Guillaumier A, Manning V, Wynne O, Gartner C, Borland R, Baker AL, et al. Electronic nicotine devices to aid smoking cessation by alcohol- and drug-dependent clients: protocol for a pilot randomised controlled trial. <i>Trials [Electronic Resource]</i> . 2018;19(1):415.	Ongoing study
Guillot CR, Stone MD, Geary BA, Kirkpatrick MG, Tidey JW, Cook JW, et al. Pharmacological, sensorimotor, and expectancy effects on tobacco withdrawal: A preliminary study. <i>Human Psychopharmacology</i> . 2015;30(5):364-71.	Wrong study design
Gupta PC, Aghi MB, Bhonsle RB, et al., Zaridze DG, Peto R. An intervention study of tobacco chewing and smoking habits for primary prevention of oral cancer among 12,212 Indian villagers. 1986:307-18.	Wrong study design
Haarmann H, Gossler A, Bonev S, Andreas S, Raupach T. A randomised controlled trial investigating the effects of varenicline on sympathovagal balance during acute nicotine withdrawal [Abstract]2014; 44(Suppl 58):[P322 p.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/281/CN-01078281/frame.html .	Cannot locate full text
Hagens P, Pieterse M, Valk P, Palen J. Effectiveness of intensive smoking reduction counselling plus combination nicotine replacement therapy in promoting long-term abstinence in patients with chronic obstructive pulmonary disease not ready to quit smoking: protocol of the REDUQ trial2017; 8:[248-57 pp.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/172/CN-01432172/frame.html .	Ongoing study
Hahn B, Harvey AN, Concheiro-Guisan M, Huestis MA, Holcomb HH, Gold JM. A test of the cognitive self-medication hypothesis of tobacco smoking in schizophrenia. <i>Biological Psychiatry</i> . 2013;74(6):436-43.	Wrong study design
Hajj M, Kheir N, Al MA, Al-Badriyeh D, Al KA, Mahfoud Z, et al. Assessing the effectiveness of a pharmacist-delivered smoking cessation program in the State of Qatar: study protocol for a randomized controlled trial2017; 16:[65 p.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/052/CN-01256052/frame.html .	Ongoing study
Hall SM, Reus VI, Munoz RF, Sees KL, Humfleet G, Hartz DT, et al. Nortriptyline and cognitive-behavioral therapy in the treatment of cigarette smoking. <i>Archives of General Psychiatry</i> . 1998;55(8):683-90.	Wrong or missing intervention

Halperin AC, McAfee TA, Jack LM, Catz SL, McClure JB, Deprey TM, et al. Impact of symptoms experienced by varenicline users on tobacco treatment in a real world setting. <i>Journal of Substance Abuse Treatment</i> . 2009;36(4):428-34.	Linked to excluded study
Hamari A, Toljamo T, Salmela M, Nieminen P. Randomized clinical study on smoking cessation in young adults. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2013;42(no pagination).	Ongoing study
Hammersley JJ, Gilbert DG, Rzetelny A, Rabinovich NE. Moderation of nicotine effects on covert orienting of attention tasks by poor placebo performance and cue validity. <i>Pharmacology, Biochemistry & Behavior</i> . 2016;149:9-16.	Wrong study design
Hammersley JJ, Rzetelny A, Gilbert DG, Rabinovich NE, Small SL, Huggenvik JI. Effects of nicotine on emotional distraction of attentional orienting: Evidence of possible moderation by dopamine type 2 receptor genotype. <i>Pharmacology Biochemistry and Behavior</i> . 2013;105:199-204.	Wrong study design
Han ES, Foulds J, Steinberg MB, Gandhi KK, West B, Richardson DL, et al. Characteristics and smoking cessation outcomes of patients returning for repeat tobacco dependence treatment. <i>International Journal of Clinical Practice</i> . 2006;60(9):1068-74.	Wrong study design
Hao R, Zhou JP, Ni L, Li QY, Shi GC. [Smoking abstinence rate and its associated factors between abrupt and gradual smoking cessation]. <i>Chung Hua Chieh Ho Ho Hu Hsi Tsa Chih</i> . 2017;40(12):898-902.	Awaiting classification
Harcouet L, Balanger S, Meunier N, Mourgues A, Grabar S, Haouili B, et al. Tobacco reduction in a prison of France. <i>Revue De Medecine Interne</i> . 2008;29(5):357-63.	Participants received same drug
Hart C, Gruer L, Bauld L. Does Smoking Reduction in Midlife Reduce Mortality Risk? Results of 2 Long-Term Prospective Cohort Studies of Men and Women in Scotland. <i>American Journal of Epidemiology</i> . 2013;178(5):770-9.	Wrong or missing intervention
Hartwell EE, Roche DJ, Ray LA. PHARMACOGENETICS OF NALTREXONE AND VARENICLINE IN HEAVY DRINKING SMOKERS. <i>Alcoholism-Clinical and Experimental Research</i> . 2014;38:223A-A.	Cannot locate full text
Hasan FM, Zagarins SE, Pischke KM, Saiyed S, Bettencourt AM, Beal L, et al. Hypnotherapy is more effective than nicotine replacement therapy for smoking cessation: Results of a randomized controlled trial. <i>Complementary Therapies in Medicine</i> . 2014;22(1):1-8.	Wrong comparator
Hatsukami DK, Ebbert JO, Edmonds A, Li C, Lin H, Le C, et al. Smokeless tobacco reduction: Preliminary study of tobacco-free snuff versus no snuff. <i>Nicotine & Tobacco Research</i> . 2008;10(1):77-85.	Wrong or missing intervention
Hatsukami DK, Hertzgaard LA, Vogel RI, Jensen JA, Murphy SE, Hecht SS, et al. Reduced nicotine content cigarettes and nicotine patch. <i>Cancer Epidemiology Biomarkers and Prevention</i> . 2013;22(6):1015-24.	Wrong comparator
Hatsukami DK, Severson H, Anderson A, Vogel RI, Jensen J, Broadbent B, et al. Randomised clinical trial of snus versus medicinal nicotine among smokers interested in product switching. <i>Tobacco control</i> . 2016;25(3):267-74.	Wrong comparator
Haustein Ko BALBWA. The effect of short-term or long-term reduction on smoking cessation; results from a placebo controlled smoking reduction study with the nicotine gum. <i>Nicotine & Tobacco Research</i> . 2003;5(2):278.	Cannot locate full text
Haustein KO, Voigt M, Haustein H, Meigen C. Treatment of Tobacco addiction with Nicotine - Experiences from the Smokers' Advisory Center in Erfurt. [German]. <i>Zeitschrift fur Allgemeinmedizin</i> . 2004;80(3):108-12.	Participants received same drug
Hayes, Inc. Bupropion for smoking cessation (Structured abstract)2011; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32011001319/frame.html .	Awaiting classification

Hayes, Inc. Chantix (Varenicline Tartrate) (Pfizer Inc.) for smoking cessation (Structured abstract)2010; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32011000793/frame.html .	Awaiting classification
Hayes, Inc. Nicotine replacement therapy for smoking cessation: nicotine gum and nicotine lozenge (Structured abstract)2011; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32011000896/frame.html .	Awaiting classification
Hayes, Inc. Nicotine replacement therapy for smoking cessation: nicotine nasal spray and nicotine inhaler (Structured abstract)2011; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32011000897/frame.html .	Awaiting classification
Hayes, Inc. Nicotine replacement therapy for smoking cessation: nicotine patch (Structured abstract)2011; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32011000898/frame.html .	Awaiting classification
Health UC, Donaghue Medical Research F. Contingency Management and Pharmacotherapy for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00683280 ; 2008.	Participants received same drug
Health UC, National Institute on Drug A. Contingency Management for Smoking Cessation in the Homeless. https://ClinicalTrials.gov/show/NCT01736982 ; 2012.	Participants received same drug
Heffner JL, Mull KE, McClure JB, Bricker JB. Positive Affect as a Predictor of Smoking Cessation and Relapse: Does It Offer Unique Predictive Value among Depressive Symptom Domains? Substance Use & Misuse. 2018;53(6):980-8.	Participants received same drug
Hennrikus D, Joseph AM, Lando HA, Duval S, Ukestad L, Kodl M, et al. Effectiveness of a smoking cessation program for peripheral artery disease patients: A randomized controlled trial. Journal of the American College of Cardiology. 2010;56(25):2105-12.	Wrong or missing intervention
Herbec AA, Chang Y, Tindle HA, Rigotti NA. Smokers' use of electronic cigarettes before, during, and in the month after hospitalization. Findings from the Helping HAND 2 Study. Addictive Behaviors. 2018;29:29.	Wrong study design
Holub M, Bodek D, Xie X, Estiverne C, Liu M, Feurdean M. Smoking cessation in hospitalized veterans. Chest. 2017;152 (4 Supplement 1):A1094.	Wrong study design
Hong H, Wang W, Chen L. Study on effect of 5A intervention method combined with varenicline for smoking cessation of COPD patients in stable stage. Chinese nursing research. 2015;29(2C):667-70.	Awaiting classification
Honisett TD. Nicotine replacement therapy for smokers admitted to intensive care. Intensive Critical Care Nursing. 2001;17(6):318-32.	Wrong study design
Hospital do C, Ministry of Health B. Cognitive Behaviour Therapy and Nicotine Replacement to Increase Tobacco Cessation. https://ClinicalTrials.gov/show/NCT01734330 ; 2010.	Ongoing study
Huber M, Weber R, Ledergerber B. Integration of structured counselling for smoking cessation in HIV outpatient care. HIV Medicine. 2009;10:130.	Cannot locate full text
Huet JD, Wewers M, Browning K, Harness J, Habash D, Ferketich A, et al. Body composition changes in HIV+ smokers undergoing smoking cessation. American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS. 2010;181(1 MeetingAbstracts).	Cannot locate full text
Hughes JR, Wadland WC, Fenwick JW, Lewis J, Bickel WK. EFFECT OF COST ON THE SELF-ADMINISTRATION AND EFFICACY OF NICOTINE GUM - A PRELIMINARY-STUDY. Preventive Medicine. 1991;20(4):486-96.	Participants received same drug

Humair JP, Tango R. Can e-cigarette help patients to reduce or stop smoking in primary care practice? Journal of General Internal Medicine. 2014;29:S480.	Wrong or missing intervention
Humair JP, Tango R. Electronic cigarette use for smoking reduction and cessation in primary care. Journal of General Internal Medicine. 2016;1):S877.	Wrong or missing intervention
Hunt N, Watkins L, McConnell O, Westall S, Holden C, Young-Jamieson J, et al. Effectiveness of smoking cessation advice following admission to hospital. European Respiratory Journal Conference: European Respiratory Society Annual Congress. 2012;40(no pagination).	Wrong comparator
Hurt RD, Offord KP, Lauger GG, Marusic Z, Fagerstrom KO, Enright PL, et al. Cessation of long-term nicotine gum use - A prospective, randomized trial. Addiction. 1995;90(3):407-13.	Wrong population
Icahn School of Medicine at Mount S, National Institute on Drug A. A Smoking Intervention Study Using Scheduled Gradual Reduction With Varenicline to Help With Cessation. https://ClinicalTrials.gov/show/NCT01772641 ; 2012.	Study terminated
Ikonomidis I, Kourea K, Pavlidis G, Tsougos E, Marinou M, Papadakis I, et al. Effect of smoking reduction on endothelial glucocalyx and arterial wave reflection during medically-aided smoking cessation program. Circulation Conference: American Heart Association. 2013;128(22 SUPPL. 1).	Cannot locate full text
Iliaz S, Onur ST, Uysal MA, Chousein EGU, Tanriverdi E, Bagci BA, et al. The Adverse Effects and Treatment Results of Smoking Cessation Pharmacotherapy During Fasting/Non-Fasting State. Substance Use & Misuse. 2017;52(8):969-73.	Wrong study design
International SRI, National Cancer I. Varenicline Tartrate With Telephone-Based Counseling and/or Internet-Based Counseling in Helping Adults Stop Smoking. https://ClinicalTrials.gov/show/NCT00301145 ; 2005.	Ongoing study
Isrctn. Evaluation of the impact of systematic delivery of cessation interventions on delivery of smoking cessation in secondary care. 2010.	Wrong study design
Isrctn. The efficacy of e-cigarettes compared with nicotine replacement therapy, when used within the UK stop smoking service. 2014.	Ongoing study
Jackson EA. Bupropion for smoking cessation. Journal of Family Practice. 1998;46(2):111-2.	Awaiting classification
Jacobsen LK, D'Souza DC, Mencil WE, Pugh KR, Skudlarski P, Krystal JH. Nicotine effects on brain function and functional connectivity in schizophrenia. Biological Psychiatry. 2004;55(8):850-8.	Wrong study design
Jiang Y, Maddison R, McRobbie H, Bullen C, Roberts V, Glover M, et al. Can exercise enhance smoking cessation outcomes? A pragmatic randomized controlled trial (fit2quit Study). Clinical Trials. 2012;9 (4):484-5.	Participants received same drug
Jiang Y, Maddison R, McRobbie H, Bullen C, Roberts V, Glover M, et al. Can exercise enhance smoking cessation outcomes? A pragmatic randomized controlled trial (fit2quit Study)2012; 9(4):[484-5 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/599/CN-00834599/frame.html .	Participants received same drug
Jimenez-Ruiz CA, Barrios M, Pena S, Cicero A, Mayayo M, Cristobal M, et al. Increasing the dose of varenicline in patients who do not respond to the standard dose. Mayo Clinic Proceedings. 2013;88(12):1443-5.	Wrong study design
Jimenez-Ruiz CA, Riesco Miranda JA, Altet Gomez N, Lorza Blasco JJ, Signes-Costa Minana J, Solano Reina S, et al. Treatment of smoking in smokers with chronic obstructive pulmonary disease. Sociedad Espanola de Neumologia y Cirugia Toracica (SEPAR). Archivos de Bronconeumologia. 2013;49(8):354-63.	Wrong study design
Jodar-Sanchez F, Carrasco Hernandez L, Nunez-Benjumea FJ, Mesa Gonzalez MA, Moreno Conde J, Parra Calderon CL, et al. Using the Social-Local-Mobile App for Smoking Cessation in the SmokeFreeBrain Project: Protocol for a Randomized Controlled Trial. JMIR Research Protocols. 2018;7(12):e12464.	Ongoing study

Johns Hopkins U, American Thoracic S. A Study of Novel Smoking Cessation Interventions in Current and Former Injection Drug Users. https://ClinicalTrials.gov/show/NCT01334736 ; 2011.	Wrong or missing intervention
Johns Hopkins U, Massachusetts General H. Trial of Integrated Smoking Cessation, Exercise and Weight Management in Serious Mental Illness: TRIUMPH. https://ClinicalTrials.gov/show/NCT02424188 ; 2015.	Ongoing study
Johns Hopkins U, National Institute on Drug A, University of Witwatersrand SA, Truth I. A Smoking Cessation Trial in HIV-infected Patients in South Africa. https://ClinicalTrials.gov/show/NCT01484340 ; 2014.	Ongoing study
Johns Hopkins U. Nicotine Dependence, Withdrawal and Replacement Therapy Assessed by PET Imaging. https://ClinicalTrials.gov/show/NCT01664741 ; 2012.	Ongoing study
Johns Hopkins U. Pre to Postoperative Smoking Cessation. https://ClinicalTrials.gov/show/NCT02683161 ; 2015.	Participants received same drug
Johns Hopkins U. Zonisamide Augmentation of Varenicline Treatment for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01685996 ; 2012.	Participants received same drug
Jones TE, Williams J. Craving control using nicotine replacement therapy in a teaching hospital. <i>Internal Medicine Journal</i> . 2012;42(3):317-22.	Wrong study design
Jorenby DE, Leischow SJ, Nides MA, Rennard SI, Johnston JA, Hughes AR, et al. A controlled trial of sustained - Release bupropion, a nicotine patch, or both for smoking cessation. <i>Revista Portuguesa de Pneumologia</i> . 2001;7(2):161-3.	Wrong study design
Jorenby DE, Smith SS, Fiore MC, Baker TB. Nicotine levels, withdrawal symptoms, and smoking reduction success in real world use: A comparison of cigarette smokers and dual users of both cigarettes and E-cigarettes. <i>Drug and Alcohol Dependence</i> . 2017;170:93-101.	Wrong comparator
Juarranz Sanz M, Soriano Llorca T, Reyes Martin G, Sanchez Sanchez MI. Efficacy of the treatment with substitute of nicotine in the smoking cessation from the Primary Health Care. [Spanish]. <i>MEDIFAM - Revista de Medicina Familiar y Comunitaria</i> . 1998;8(6):376-83.	Awaiting classification
Kalechstein AD, Mahoney JJ, Verrico CD, De La Garza R. Short-term, low-dose varenicline administration enhances information processing speed in methamphetamine-dependent users. <i>Neuropharmacology</i> . 2014;85:493-8.	Wrong study design
Kalkhoran S, Inman E, Kulesa Kelley JH, Ashburner JM, Rigotti NA. Proactive population health strategy to offer tobacco dependence treatment to smokers in primary care practice network. <i>Journal of General Internal Medicine</i> . 2018;33 (2 Supplement 1):312-3.	Wrong study design
Kalkhoran S, Kruse GR, Chang YC, Rigotti NA. Smoking-Cessation Efforts by US Adult Smokers with Medical Comorbidities. <i>American Journal of Medicine</i> . 2018;131(3).	Wrong study design
Kallan FV, Brandt CJ, Ellegaard H, Joensen MB, Sorknaes AD, Tougaard L. The diagnosis of "smoker's lung" encourages smoking cessation. <i>Ugeskrift for laeger</i> . 1997;159(44):6528-30.	Wrong or missing intervention
Kalman D, Hoskinson R, Sambamoorthi U, Garvey AJ. A prospective study of persistence in the prediction of smoking cessation outcome: results from a randomized clinical trial. <i>Addictive Behaviors</i> . 2010;35(2):179-82.	Participants received same drug
Katz MM, Harris SM, Polydorou S, Emmanouel MD, Grossman E. Enhanced smoking cessation services via on-site nicotine replacement therapy (NRT) in an opioid treatment program (OTP). <i>Journal of General Internal Medicine</i> . 2015;30:S532.	Cannot locate full text
Kawachi E, Imaizumi S, Takata K, Abe S, Matsuo Y, Yahiro E, et al. Effect of smoking cessation on HDL function. <i>European Heart Journal</i> . 2014;35:902.	Cannot locate full text

Kct. Effect of an electronic cigarette for smoking reduction and cessation in Korean male smokers: a randomized, controlled study. 2014.	Ongoing study
Killen JD, Robinson TN, Ammerman S, Hayward C, Rogers J, Stone C, et al. Randomized clinical trial of the efficacy of Bupropion combined with nicotine patch in the treatment of adolescent smokers. <i>Journal of Consulting and Clinical Psychology</i> . 2004;72(4):729-35.	Wrong population
Kimmel SE, Berlin JA, Miles C, Jaskowiak J, Carson JL, Strom BL. Risk of acute first myocardial infarction and use of nicotine patches in a general population. <i>Journal of the American College of Cardiology</i> . 2001;37(5):1297-302.	Wrong study design
King BA, Alam S, Promoff G, Arrazola R, Dube SR. Awareness and Ever-Use of Electronic Cigarettes Among U.S. Adults, 2010-2011. <i>Nicotine & Tobacco Research</i> . 2013;15(9):1623-7.	Wrong study design
Kisely SR, Wise M, Preston N, Malmgren S, Shannon P. A group intervention to reduce smoking in individuals with psychiatric disorder: Brief report of a pilot study. <i>Australian and New Zealand Journal of Public Health</i> . 2003;27(1):61-3.	Wrong study design
Kittle J, Lopes RD, Huang M, Marquess ML, Wilson MD, Ascher J, et al. Cardiovascular adverse events in the drug-development program of bupropion for smoking cessation: A systematic retrospective adjudication effort. <i>Clinical Cardiology</i> . 2017;40(10):899-906.	Wrong study design
Klesges RC, DeBon M, Vander Weg MW, Haddock CK, Lando HA, Relyea GE, et al. Efficacy of a tailored tobacco control program on long-term use in a population of US military troops. <i>Journal of Consulting and Clinical Psychology</i> . 2006;74(2):295-306.	Wrong study design
Klesges RC, Klesges LM, Vander Weg MW, DeBon M, Poston WSC, Ebbert J, et al. Characteristics of Air Force Personnel Who Choose Pharmacological Aids for Smoking Cessation Following an Involuntary Tobacco Ban and Tobacco Control Program. <i>Health Psychology</i> . 2007;26(5):588-97.	Linked to excluded study
Klesges RC, Krukowski RA, Klosky JL, Liu W, Srivastava DK, Boyett JM, et al. Efficacy of a tobacco quitline among adult survivors of childhood cancer. <i>Nicotine and Tobacco Research</i> . 2015;17(6):710-8.	Participants received same drug
Kleykamp BA, Jennings JM, Sams C, Weaver MF, Eissenberg T. The influence of transdermal nicotine on tobacco/nicotine abstinence and the effects of a concurrently administered cigarette in women and men. <i>Experimental & Clinical Psychopharmacology</i> . 2008;16(2):99-112.	Wrong study design
Klonizakis M, Crank H, Gumber A, Brose L. Smokers making a quit attempt using e-cigarettes with or without nicotine or prescription nicotine replacement therapy: impact on cardiovascular function (ISME-NRT) - a study protocol 2017; 17(1):[293 p.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/082/CN-01401082/frame.html .	Ongoing study
Kocak ND, Akturk UA. What factors influence non-adherence to smoking cessation support programs? <i>European Respiratory Journal Conference: European Respiratory Society International Congress, ERS</i> . 2017;50(Supplement 61).	Wrong study design
Kollins SH, English JS, Itchon-Ramos N, Chrisman AK, Dew R, O'Brien B, et al. A Pilot Study of Lis-Dexamfetamine Dimesylate (LDX/SPD489) to Facilitate Smoking Cessation in Nicotine-Dependent Adults With ADHD. <i>Journal of Attention Disorders</i> . 2014;18(2):158-68.	Wrong comparator
Kotz D, Brown J, West R. Effectiveness of varenicline versus nicotine replacement therapy for smoking cessation with minimal professional support: Evidence from an English population study. <i>Psychopharmacology</i> . 2014;231(1):37-42.	Wrong study design
Kotz D, Brown J, West R. Prospective cohort study of the effectiveness of smoking cessation treatments used in the "real world". <i>Mayo Clinic Proceedings</i> . 2014;89(10):1360-7.	Wrong study design
Kotz D, Brown J, West R. Prospective cohort study of the effectiveness of varenicline versus nicotine replacement therapy for smoking cessation in the "real world". <i>BMC public health</i> . 2014;14:1163.	Wrong study design
Kotz D, Brown J, West R. 'Real-world' effectiveness of smoking cessation treatments: a population study. <i>Addiction (Abingdon, England)</i> . 2014;109(3):491-9.	Wrong study design

Kotz D, Huibers MJH, West RJ, Wesseling G, van Schayck OCP. What mediates the effect of confrontational counselling on smoking cessation in smokers with COPD? Patient Education and Counseling. 2009;76(1):16-24.	Linked to excluded study
Kotz D, Simpson C, Viechtbauer W, Van Schayck OCP, West R, Sheikh A. Cardiovascular and neuropsychiatric safety of varenicline and bupropion compared with nicotine replacement therapy for smoking cessation: Study protocol of a retrospective cohort study using the QResearch general practice database. BMJ Open. 2014;4 (8) (no pagination)(e005281).	Cannot locate full text
Kotz D, Wesseling G, Huibers MJH, van Schayck OCP. Efficacy of confrontational counselling for smoking cessation in smokers with previously undiagnosed mild to moderate airflow limitation: study protocol of a randomized controlled trial. BMC Public Health. 2007;7:332.	Linked to excluded study
Kralikova E, Kubatova S, Truneckova K, Kmetova A, Hajek P. THE ELECTRONIC CIGARETTE: WHAT PROPORTION OF SMOKERS HAVE TRIED IT AND HOW MANY USE IT REGULARLY? Addiction. 2012;107(8):1529-.	Wrong study design
Krumpe P, Malani N, Adler J, Ramoorthy S, Asadi S, Corwin N, et al. EFFICACY OF TRANSDERMAL NICOTINE ADMINISTRATIONS AS AN ADJUNCT FOR SMOKING CESSATION IN HEAVILY NICOTINE ADDICTED SMOKERS. American Review of Respiratory Disease. 1989;139(4 PART 2):A337-A.	Awaiting classification
Kuhn L, Brown B, Wigle P, Tolle E, Espel M, Curington R. Targeting self-efficacy in a charitable pharmacy smoking cessation program for the underserved population. Journal of the American Pharmacists Association Conference. 2017;57(3).	Ongoing study
Landfeldt B, Kruse E, Westin A, Mattson K, Lojander J. Nicotine replacement treatment in heavy smokers nicotine nasal spray combined with nicotine patch in a double-blind controlled study. European Respiratory Journal. 1998;12(SUPPL. 28):154S-S.	Awaiting classification
Lando HA, Rolnick S, Klevan D, Roski J, Cherney L, Lauger G. Telephone support as an adjunct to transdermal nicotine in smoking cessation. American Journal of Public Health. 1997;87(10):1670-4.	Participants received same drug
Lasser KE, Quintiliani LM, Truong V, Xuan Z, Murillo J, Jean C, et al. Effect of patient navigation and financial incentives on smoking cessation among primary care patients at an urban safety-net hospital: A randomized clinical trial. JAMA Internal Medicine. 2017;177(12):1798-807.	Participants received same drug
Lasser KE, Quintiliani LM, Truong V, Xuan Z, Pbert L. Patient navigation to promote smoking cessation in primary care: Preliminary findings from an ongoing randomized controlled trial. Journal of General Internal Medicine. 2017;32 (2 Supplement 1):S266.	Participants received same drug
Laude JR, Bailey SR, Crew E, Varady A, Lembke A, McFall D, et al. Extended treatment for cigarette smoking cessation: a randomized control trial. Addiction. 2017;112(8):1451-9.	Wrong study design
Lazary J, Dome P, Faludi G. Significant factors on success rate of nicotine addiction treatment: A prospective study. European Neuropsychopharmacology. 2011;21:S594-S5.	Cannot locate full text
Li J, Bullen C, Newcombe R, Walker N, Walton D. The use and acceptability of electronic cigarettes among New Zealand smokers. New Zealand Medical Journal. 2013;126(1375):48-57.	Wrong study design
Liang J, Abramson M, Zwar N, Russell G, Holland A, Bonevski B, et al. Preferences and experiences of smokers attending Australian general practices. European Respiratory Journal Conference: European Respiratory Society International Congress, ERS. 2017;50(Supplement 61).	Wrong study design
Liang J, Abramson MJ, Zwar N, Russell G, Holland AE, Bonevski B, et al. Quitting experiences and smoking cessation preferences of smokers in Australian general practices. Respirology. 2017;22 (Supplement 2):34.	Wrong study design
Liang J, Abramson MJ, Zwar NA, Russell GM, Holland AE, Bonevski B, et al. Diagnosing COPD and supporting smoking cessation in general practice: Evidenceepractice gaps. Medical Journal of Australia. 2018;208(1):29-34.	Wrong study design

Ling AC, Elward DD, Barry S. A retrospective cohort study of the long term effectiveness of smoking cessation counselling. Thorax. 2011;66:A116-A7.	Wrong study design
Little SJ, Stevens VJ, Severson HH, Lichtenstein E. Effective smokeless tobacco intervention for dental hygiene patients. Journal of dental hygiene : JDH. 1992;66(4):185-90.	Linked to excluded study
Lopes NH, Buzo CG, Lima Santos PCJ, Pereira AC, Paulitch F, Abe TMO, et al. Nicotine replacement therapy was more effective in elderly people of a cardiovascular hospital. European Journal of Preventive Cardiology. 2014;1):S17.	Cannot locate full text
Loreto AR, Carvalho CFC, Frallonardo FP, Ismael F, de Andrade AG, Castaldelli-Maia JM. Smoking Cessation Treatment for Patients With Mental Disorders Using CBT and Combined Pharmacotherapy. Journal of Dual Diagnosis. 2017;13(4):238-46.	Wrong study design
Lou P, Zhu Y, Chen P, Zhang P, Yu J, Zhang N, et al. Supporting smoking cessation in chronic obstructive pulmonary disease with behavioral intervention: a randomized controlled trial. BMC Family Practice. 2013;14:91.	Wrong or missing intervention
Lu CC, Hsiao YC, Huang HW, Lin JY, Huang CL. Effects of a Nurse-Led, Stage-Matched, Tailored Program for Smoking Cessation in Health Education Centers: A Prospective, Randomized, Controlled Trial. Clin Nurs Res. 2018:1054773817754276.	Wrong or missing intervention
MacPherson L, Tull MT, Matusiewicz AK, Rodman S, Strong DR, Kahler CW, et al. Randomized Controlled Trial of Behavioral Activation Smoking Cessation Treatment for Smokers With Elevated Depressive Symptoms. Journal of Consulting and Clinical Psychology. 2010;78(1):55-61.	Participants received same drug
Maggie M Sweitzer P, National Institute on Drug A, Duke U. Neuroimaging Reward, Behavioral Treatment, and Smoking Cessation. https://ClinicalTrials.gov/show/NCT02927847 ; 2016.	Participants received same drug
Mahmarijan JJ, Moye LA, Nasser GA, Nagueh SF, Bloom MF, Benowitz NL, et al. Nicotine patch therapy in smoking cessation reduces the extent of exercise-induced myocardial ischemia. Journal of the American College of Cardiology. 1997;30(1):125-30.	Wrong study design
Mankani SK, Garabrant DH, Homa DM. Effectiveness of nicotine patches in a workplace smoking cessation program. An eleven-month follow-up study. Journal of Occupational & Environmental Medicine. 1996;38(2):184-9.	Participants received same drug
Manzoli L, Flacco ME, Fiore M, La Vecchia C, Marzuillo C, Gualano MR, et al. Electronic cigarettes efficacy and safety at 12 months: Cohort study. PLoS ONE. 2015;10 (6) (no pagination)(e0129443).	Wrong comparator
Manzoli L, La Vecchia C, Flacco ME, Capasso L, Simonetti V, Boccia S, et al. Multicentric cohort study on the long-term efficacy and safety of electronic cigarettes: study design and methodology. BMC Public Health. 2013;13:883.	Wrong comparator
Marcos T, Godas T, Corominas J. Nicotine replacement therapy versus gradual smoking withdrawal in smoking cessation. [Spanish]. Medicina Clinica. 2004;123(4):127-30.	Wrong comparator
Marcus BH, Lewis BA, Hogan J, King TK, Albrecht AE, Bock B, et al. The efficacy of moderate intensity exercise as an aid for smoking cessation in women: A randomized controlled trial. Nicotine and Tobacco Research. 2005;7(6):871-80.	Participants received same drug
Mardle T, Merrett S, Wright J, Percival F, Lockhart I, Marshall S. Real world evaluation of different smoking cessation service models in England. Value in Health. 2011;14 (7):A500.	Wrong study design

Mardle T, Merrett S, Wright J, Percival F, Lockhart I. Real world evaluation of three models of NHS smoking cessation service in England. BMC research notes. 2012;5:9.	Wrong study design
Marks DF, Sykes CM. Randomized controlled trial of cognitive behavioural therapy for smokers living in a deprived are of London: Outcome at one-year follow-up. Psychology, Health and Medicine. 2002;7(1):17-24.	Wrong or missing intervention
Marshall A, Raw M. NICOTINE CHEWING GUM IN GENERAL-PRACTICE - EFFECT OF FOLLOW UP APPOINTMENTS. British Medical Journal. 1985;290(6479):1397-8.	Participants received same drug
Masouredis CM, Hilton JF, Grady D, Gee L, Chesney M, Hengl L, et al. A spit tobacco cessation intervention for college athletes: three-month results. Advances in dental research. 1997;11(3):354-9.	Wrong or missing intervention
Massachusetts General H, National Alliance for Research on S, Depression, The Bowman Family F. Study of the Effects of Mecamylamine and Varenicline in Schizophrenia. https://ClinicalTrials.gov/show/NCT00455650 ; 2007.	Wrong population
Massachusetts General H, National Cancer I, National Institutes of H. Integrating Tobacco Treatment Into Cancer Care: A Randomized Controlled Comparative Effectiveness Trial. https://ClinicalTrials.gov/show/NCT01871506 ; 2013.	Ongoing study
Massachusetts General H, National Heart L, Blood I. Smoking Interventions for Hospital Patients. https://ClinicalTrials.gov/show/NCT01177176 ; 2010.	Wrong study design
Massachusetts General H, National Institute on Drug A, McLean H, GlaxoSmithKline. Effectiveness of GSK598809, a Selective D3 Antagonist, Added to CBT and NRT for Smoking Cessation and Relapse Prevention. https://ClinicalTrials.gov/show/NCT01188967 ; 2010.	Wrong or missing intervention
Massachusetts General H, Southern Methodist U, University of H, National Institute on Drug A. Integrated Treatment for Smoking Cessation & Anxiety in People With HIV Phase 1. https://ClinicalTrials.gov/show/NCT01393301 ; 2011.	Participants received same drug
Massachusetts General H. Chantix in Adult Smokers With Attention Deficit Hyperactivity Disorder (ADHD). https://ClinicalTrials.gov/show/NCT00907218 ; 2009.	Study terminated
Massachusetts General H. Extended Duration Nicotine Replacement Therapy and Bupropion in Smokers With Schizophrenia. https://ClinicalTrials.gov/show/NCT01576640 ; 2005.	Study terminated
Mayer SA, Chong JY, Ridgway E, Min KC, Commichau C, Bernardini GL. Delirium from nicotine withdrawal in neuro-ICU patients. Neurology. 2001;57(3):551-3.	Wrong study design
Mayo C. Varenicline and Alcohol in Inpatient Addictions Program (IAP). https://ClinicalTrials.gov/show/NCT01169610 ; 2011.	Wrong study design
Mayoral R, Gonzalez F, Ruiz J, Crespo M, Molina A, Lopez P, et al. Effectiveness of antismoking drug treatment in patients with psychiatric comorbidity compared with patients without this comorbidity. Journal of the Egyptian Public Health Association. 2017;92(2):61-7.	Wrong study design
McChargue DE, Collins FL, Jr., Cohen LM. Effect of non-nicotinic moist snuff replacement and lobeline on withdrawal symptoms during 48-h smokeless tobacco deprivation. Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco. 2002;4(2):195-200.	Wrong or missing intervention
McClure JB, Jack L, Deprey M, Catz S, McAfee T, Zbikowski S, et al. Canary in a coal mine? Interest in bupropion SR use among smokers in the COMPASS trial. Nicotine & Tobacco Research. 2008;10(12):1815-6.	Wrong study design

McClure JB, Swan GE, St John J, Fauver R, Javitz HS, Bergen AW, et al. Pharmacogenetic smoking cessation intervention in a health care setting: a pilot feasibility study. <i>Nicotine & Tobacco Research</i> . 2013;15(2):518-26.	Wrong study design
McFall M, Saxon AJ, Thompson CE, Yoshimoto D, Malte C, Straits-Troster K, et al. Improving the rates of quitting smoking for veterans with posttraumatic stress disorder. <i>American Journal of Psychiatry</i> . 2005;162(7):1311-9.	Wrong study design
McGain F, Durie ML, Bates S, Polmear CM, Meyer J, French CJ, et al. Smoking cessation therapy in Australian and New Zealand intensive care units: a multicentre point prevalence study. <i>Critical Care & Resuscitation</i> . 2018;20(1):68-73.	Awaiting classification
McGill U. Evaluating the Efficacy of E-Cigarette Use for Smoking Cessation (E3) Trial. https://ClinicalTrials.gov/show/NCT02417467 ; 2016.	Ongoing study
McKay-Brown L, Borland R, Balmford J, Segan CJ, Andrews C, Tasker C, et al. The challenges of recruiting and retaining GPs in research: Findings from a smoking cessation project. <i>Australian Journal of Primary Health</i> . 2007;13(1):61-7.	Linked to excluded study
McKee SA, Harrison ELR, Cummings KM, Borland R, Kahler CW, Fong GT, et al. Varenicline use is associated with reductions in alcohol consumption among smokers in the international tobacco control four country survey. <i>Alcoholism: Clinical and Experimental Research</i> . 2011;35:22A.	Wrong study design
McKee SA, Young-Wolff KC, Harrison ELR, Cummings KM, Borland R, Kahler CW, et al. Longitudinal Associations Between Smoking Cessation Medications and Alcohol Consumption Among Smokers in the International Tobacco Control Four Country Survey. <i>Alcoholism: Clinical and Experimental Research</i> . 2013;37(5):804-10.	Wrong study design
McLean H, Harvard Medical S. Cognitive Behavioral Therapy and the Nicotine Transdermal Patch for Cannabis Dependence and Nicotine Dependence. https://ClinicalTrials.gov/show/NCT01292642 ; 2009.	Participants received same drug
McNeil AB, Johnson, Johnson C, Personal Products W. Bioequivalence Between an Oral Nicotine Replacement Product and Nicorette® Gum. https://ClinicalTrials.gov/show/NCT01152749 ; 2009.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Bioequivalence Between Nicotine Replacement Products and Nicorette® Gum. https://ClinicalTrials.gov/show/NCT01113424 ; 2008.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Bioequivalence Between Two New Oral Nicotine Replacement Therapy Products and Nicorette® Microtab. https://ClinicalTrials.gov/show/NCT01238640 ; 2008.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Bioequivalence Between Two Oral Nicotine Sublingual Tablets, 2 mg and 4 mg. https://ClinicalTrials.gov/show/NCT01238627 ; 2007.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Comparative Pharmacokinetic Study of New Oral Nicotine Replacement Therapy Products. https://ClinicalTrials.gov/show/NCT01227720 ; 2009.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Early Effects of a New Oral Nicotine Replacement Product and NiQuitin™ Lozenge. https://ClinicalTrials.gov/show/NCT01075659 ; 2010.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Multiple-dose Nicotine Pharmacokinetics With a New Oral Nicotine Replacement Product. https://ClinicalTrials.gov/show/NCT01084707 ; 2009.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Multiple-dose Nicotine Pharmacokinetics With Three Oral Nicotine Replacement Products. https://ClinicalTrials.gov/show/NCT01321931 ; 2011.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Nicotine Pharmacodynamics With a New Oral Nicotine Replacement Therapy and Nicotine Gum. https://ClinicalTrials.gov/show/NCT01368016 ; 2011.	Wrong comparator
McNeil AB, Johnson, Johnson C, Personal Products W. Pharmacokinetics With a New Oral Nicotine Replacement Product and Nicorette® Gum. https://ClinicalTrials.gov/show/NCT01152736 ; 2008.	Wrong comparator

McNeil AB, Johnson, Johnson C, Personal Products W. Pharmacokinetics With Two New Oral Nicotine Replacement Products and Nicorette® Gum. https://ClinicalTrials.gov/show/NCT01152723 ; 2008.	Wrong comparator
McNeil AB, Johnson, Johnson C, Personal Products W. Pilot Study of a New Nicotine Replacement Therapy. https://ClinicalTrials.gov/show/NCT00764439 ; 2008.	Ongoing study
McNeil AB, Johnson, Johnson C, Personal Products W. Single-dose Nicotine Pharmacokinetics With a New Oral Nicotine Replacement Product. https://ClinicalTrials.gov/show/NCT01084603 ; 2009.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Single-dose Nicotine Pharmacokinetics With Four Oral Nicotine Replacement Products. https://ClinicalTrials.gov/show/NCT01234896 ; 2010.	Wrong study design
McNeil AB, Johnson, Johnson C, Personal Products W. Single-dose Nicotine Pharmacokinetics With Three Oral Nicotine Replacement Products. A Study in Healthy Smokers. https://ClinicalTrials.gov/show/NCT01234792 ; 2008.	Ongoing study
McNeil Consumer Healthcare Division of McNeil-Ppc I, GlaxoSmithKline, McNeil AB, Johnson, Johnson C, Personal Products W. Study to Determine the Efficacy and Safety of a Novel Nicotine Replacement Therapy. https://ClinicalTrials.gov/show/NCT02355665 ; 2015.	Wrong or missing intervention
McNutt MD. Examining weight gain in treatment-seeking African American smokers: A biopsychosocial approach. Dissertation Abstracts International: Section B: The Sciences and Engineering. 2017;78(3-B(E)):No-Specified.	Linked to excluded study
McRobbie H, Phillips A, Goniewicz ML, Smith KM, Knight-West O, Przulj D, et al. Effects of Switching to Electronic Cigarettes with and without Concurrent Smoking on Exposure to Nicotine, Carbon Monoxide, and Acrolein. <i>Cancer Prev Res (Phila)</i> . 2015;8(9):873-8.	Wrong study design
McRobbie H, Przulj D, Smith KM, Cornwall D. Complementing the Standard Multicomponent Treatment for Smokers With Denicotinized Cigarettes: A Randomized Trial. <i>Nicotine & Tobacco Research</i> . 2016;18(5):1134-41.	Wrong study design
Medical University of South C, National Cancer I. Gain-framed Messages and NRT for Lung Cancer Screening Patients. https://ClinicalTrials.gov/show/NCT03069924 ; 2017.	Ongoing study
Medical University of South C. Gender Differences in Response to Nicotine Replacement Therapy and De-Nicotinized Cigarettes. https://ClinicalTrials.gov/show/NCT00960778 ; 2009.	Wrong study design
Medical University of W. The Influence of Electronic and Traditional Cigarettes Smoking on Hemodynamic Parameters. https://ClinicalTrials.gov/show/NCT03037775 ; 2015.	Wrong study design
Meier-Lammermann E, Buttner H, Klupfel P, Trapp VE, Wellhofer G, Bolcskei PL. Development of a psychologically founded program for smoking cessation with adjuvant use of nicotine-containing chewing gum. [German]. <i>Pneumologie (Stuttgart, Germany)</i> . 1990;44 Suppl 1:116-7.	Wrong study design
Memorial Sloan Kettering Cancer C, Center MDAC. Smoking Cessation Intervention for Cancer Patients. https://ClinicalTrials.gov/show/NCT00575718 ; 2002.	Participants received same drug
Memorial Sloan Kettering Cancer C, Weill Medical College of Cornell U. Investigation of the Effects of Electronic Cigarettes on Vascular Health. https://ClinicalTrials.gov/show/NCT03041493 ; 2017.	Wrong comparator
Metelitsa VI. PHARMACOLOGICAL AGENTS AGAINST SMOKING. <i>Byulleten' Vsesoyuznogo Kardiologicheskogo Nauchnogo Tsentra Amn SSSR</i> . 1987;10(1):109-12.	Wrong study design
Metse A, Bowman J, Wye P, Stockings E, Adams M, Clancy R, et al. Evaluating the efficacy of an integrated smoking cessation intervention for mental health patients: Study protocol for a randomized controlled trial. <i>Asia-Pacific Journal of Clinical Oncology</i> . 2014;10:144.	Participants received same drug

Metse A, Bowman J, Wye P, Wiggers J. Mental health patients' receptivity to and uptake of a smoking cessation intervention, and reported changes in smoking behaviour. <i>Asia-Pacific Journal of Clinical Oncology</i> . 2014;10:144.	Participants received same drug
Metse A, Wiggers J, Wye P, Wolfenden L, Freund M, Clancy R, et al. Efficacy of a universal smoking cessation intervention initiated in inpatient psychiatry and continued post-discharge: a randomised controlled trial 2017; 51(4):[366-81 pp.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/129/CN-01374129/frame.html .	Participants received same drug
Metse AP, Bowman JA, Wye P, Stockings E, Adams M, Clancy R, et al. Evaluating the efficacy of an integrated smoking cessation intervention for mental health patients: study protocol for a randomised controlled trial. 2014;15:266.	Participants received same drug
Metse AP, Wiggers J, Wye P, Clancy R, Moore L, Adams M, et al. Uptake of smoking cessation aids by smokers with a mental illness. <i>Journal of Behavioral Medicine</i> . 2016;39(5):876-86.	Linked to excluded study
Miller MJ, Draugalis JR, Ortmeier BG, Leischow SJ. A retrospective analysis of the costs and consequences of a tobacco-cessation program for active duty service members. <i>Military Medicine</i> . 1996;161(7):420-4.	Wrong study design
Milton SHMC. Penn State TXT2Quit Study. https://ClinicalTrials.gov/show/NCT02367391 ; 2015.	Participants received same drug
Minneker E, Buchkremer G, Bents H. Predictors of successful smoking cessation. [German]. <i>Suchtgefahren</i> . 1989;35(2):91-102.	Awaiting classification
Minneker E, Buchkremer G, Block M. THE EFFECT OF DIFFERENT DOSAGES OF A TRANSDERMAL NICOTINE SUBSTITUTION SYSTEM ON THE SUCCESS RATE OF SMOKING CESSATION THERAPY. <i>Methods and Findings in Experimental and Clinical Pharmacology</i> . 1989;11(3):219-22.	Wrong study design
Mirza S, Bockorny B, Sittig R, Bastian L. Inpatient smoking cessation counseling and treatment for patients with lung cancer. <i>Chest Conference: CHEST</i> . 2014;146(4 MEETING ABSTRACT).	Wrong study design
Mitchell M, Leone F, Williams K. Varenicline for smoking cessation (Structured abstract) 2009; (4). Available from: http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32009100416/frame.html .	Awaiting classification
Moderation of nicotine effects on covert orienting of attention tasks by poor placebo performance and cue validity 2016. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/164/CN-01196164/frame.html .	Wrong study design
Mohiuddin SM, Mooss AN, Hunter CB, Grollmes TL, Cloutier DA, Hilleman DE. Intensive smoking cessation intervention reduces mortality in high-risk smokers with cardiovascular disease. <i>Chest</i> . 2007;131(2):446-52.	Wrong study design
Monarrez-Espino J, Galanti MR, Hansson J, Janszky I, Soderberg-Lofdal K, Moller J. Treatment With Bupropion and Varenicline for Smoking Cessation and the Risk of Acute Cardiovascular Events and Injuries: a Swedish Case-Crossover Study. <i>Nicotine & Tobacco Research</i> . 2018;20(5):606-13.	Wrong study design
Moolchan ET, Robinson ML, Ernst M, Cadet JL, Pickworth WB, Heishman SJ, et al. Safety and efficacy of the nicotine patch and gum for the treatment of adolescent tobacco addiction. <i>Pediatrics</i> . 2005;115(4):e407-e14.	Wrong population
Mooney ME, Poling J, Gonzalez G, Gonsai K, Kosten T, Sofuoglu M. Preliminary study of buprenorphine and bupropion for opioid-dependent smokers. <i>American Journal on Addictions</i> . 2008;17(4):287-92.	Wrong study design
Mooney ME, Schmitz JM, Allen S, Grabowski J, Pentel P, Oliver A, et al. Bupropion and naltrexone for smoking cessation: A double-blind randomized placebo-controlled clinical trial. <i>Clinical Pharmacology and Therapeutics</i> . 2016:344-52.	Wrong comparator

Morissette SB, Gulliver SB, Kamholz BW, Spiegel DA, Tiffany ST, Barlow DH. Transdermal nicotine during cue reactivity in adult smokers with and without anxiety disorders. <i>Psychology of Addictive Behaviors</i> . 2012;26(3):507-18.	Wrong study design
Morris CD, Waxmonsky JA, May MG, Tinkelman DG, Dickinson M, Giese AA. Smoking reduction for persons with mental illnesses: 6-month results from community-based interventions. <i>Community Mental Health Journal</i> . 2011;47(6):694-702.	Participants received same drug
Mueller SE, Petitjean SA, Wiesbeck GA. Cognitive behavioral smoking cessation during alcohol detoxification treatment: A randomized, controlled trial. <i>Drug and Alcohol Dependence</i> . 2012;126(3):279-85.	Wrong study design
Mundt MP, Baker TB, Fraser DL, Smith SS, Piper ME, Fiore MC. Paying Low-Income Smokers to Quit? The Cost Effectiveness of Incentivizing Tobacco Quit Line Engagement for Medicaid Recipients Who Smoke. <i>Value in Health</i> . 2018.	Linked to excluded study
Muramoto ML, Leischow SJ, Sherrill D, Matthews E, Strayer LJ. Randomized, double-blind, placebo-controlled trial of 2 dosages of sustained-release Bupropion for adolescent smoking cessation. <i>Archives of Pediatrics & Adolescent Medicine</i> . 2007;161(11):1068-74.	Wrong population
Murphy JM, Mahoney MC, Cummings KM, Hyland AJ, Lawvere S. A randomized trial to promote pharmacotherapy use and smoking cessation in a Medicaid population (United States). <i>Cancer Causes and Control</i> . 2005;16(4):373-82.	Wrong or missing intervention
Murray RL, Leonardi-Bee J, Marsh J, Jayes L, Li J, Parrott S, et al. Systematic identification and treatment of smokers by hospital based cessation practitioners in a secondary care setting: Cluster randomised controlled trial. <i>BMJ (Online)</i> . 2013;347 (7917) (no pagination)(f4004).	Wrong study design
Murray RL, Szatkowski L, Ussher M. Evaluation of a Refined, Nationally Disseminated Self-Help Intervention for Smoking Cessation ("Quit Kit-2"). <i>Nicotine and Tobacco Research</i> . 2013;15(8):1365-71.	Wrong study design
Nahas GJ, Wilson D, Talbot V, Cartmell KB, Warren GW, Toll BA, et al. Feasibility of implementing a hospital-based "opt-out" tobacco-cessation service. <i>Nicotine & Tobacco Research</i> . 2017;19(8):937-43.	Wrong study design
Nascimento OA, Barros JO, Santos NR, Oliveira NC, Vicente R, Jardim JR. Related factors to smoking cessation rate in individuals with and without COPD after 3 and 12 months in a smoking cessation primary care attention center (prevfumo). <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2018;197(MeetingAbstracts).	Wrong study design
National Institute on Drug A, National Institutes of Health Clinical C. Effect of Varenicline on Tobacco Addiction and on Abstinence-induced Cognitive Impairment. https://ClinicalTrials.gov/show/NCT01309685 ; 2011.	Study withdrawn
National Institute on Drug A. A Preliminary Study of Sustained-Release Bupropion for Smoking Cessation in Bipolar Affective Disorder. https://ClinicalTrials.gov/show/NCT00593099 ; 2004.	Ongoing study
National Institute on Drug A. Bupropion as an Adjunct to the Nicotine Patch Plus CBT. https://ClinicalTrials.gov/show/NCT00142831 ; 1999.	Ongoing study
National Institute on Drug A. Combination Pharmacotherapy for Smoking Cessation Among Methadone Patients. https://ClinicalTrials.gov/show/NCT00060814 ; 2002.	Wrong study design
National Institute on Drug A. Effect of Bupropion on Smoking Behavior in Smokers With Schizophrenia. https://ClinicalTrials.gov/show/NCT00218231 ; 2004.	Ongoing study
National Institute on Drug A. Effects of Transdermal Nicotine on Smoking, Craving and Withdrawal in People With Schizophrenia. https://ClinicalTrials.gov/show/NCT00218218 ; 2002.	Ongoing study
National Institute on Drug A. Impact of Smoking Cessation on Sleep - 5. https://ClinicalTrials.gov/show/NCT00132821 ; 2005.	Ongoing study
National Institute on Drug A. Reduction of Smoking in Cardiac Disease Patients. https://ClinicalTrials.gov/show/NCT00469885 ; 2001.	Ongoing study

National Institute on Drug Dependence C, Bronx VAMC. Association of Functional COMT Val108/Met Polymorphism With Smoking Cessation in Nicotine Replacement Therapy. https://ClinicalTrials.gov/show/NCT01980550 ; 2004.	Ongoing study
National Taiwan University H, Ministry of S, Technology T. The Effectiveness of Smoking Cessation Clinics Combined With Coach-assisted Lifestyle Change in Prediabetic Smokers. https://ClinicalTrials.gov/show/NCT01926041 ; 2013.	Wrong or missing intervention
Nct, Affairs USDoV, Research VAoo, Development. Non-Nicotine Agents for Smoking Cessation. 2003.	Awaiting classification
Nct, Affairs USDoV, Research VAoo, Development. Proactive Tobacco Treatment for Veterans. 2009.	Linked to excluded study
Nct, Affairs USDoV, Research VAoo, Development. Tobacco Cessation for Veterans With Post Traumatic Stress Disorder (PTSD). 2009.	Wrong or missing intervention
Nct, Affairs USDoV, Research VAoo, Development. Treatment to Quit Smoking. 1997.	Wrong or missing intervention
Nct, Agatha C, Patient-Centered Outcomes Research I, University of K. The PATH Home Trial: A Comparative Effectiveness Study of Peripartum Opioid Use Disorder in Rural Kentucky. 2019.	Wrong population
Nct, Albert Einstein College of Medicine I, National Institutes of H, Pfizer, National Institute on Drug A. A Trial of Directly Observed and Long-term Varenicline. 2018.	Ongoing study
Nct, Arizona State U, National Institute on Drug A, Los Angeles Clinical T, University of Nevada R, Pfizer. Varenicline OTC Trial on Efficacy and Safety. 2018.	Ongoing study
Nct, Assistance Publique - Hôpitaux de P. Randomized Trial of Electronic Cigarettes With or Without Nicotine in Smoking Cessation. 2018.	Ongoing study
Nct, Assistance Publique - Hôpitaux de P. Smoking Tobacco Cessation Integrated Program of Patients Treated for the Head and the Neck Cancer. 2018.	Wrong or missing intervention
Nct, Azienda Ospedaliera Universitaria di Bologna Policlinico SOM, Fondazione Salvatore M. Acute Physiological Effects of Electronic Sigarette vs Cigarette Smoking. 2014.	Linked to excluded study
Nct, Birmingham UoAa, Institute NC. Integrated Smoking Cessation Treatment for Low Income Community Corrections. 2009.	Linked to excluded study
Nct, Boston University Charles River C. Engaging Working Memory and Distress Tolerance to Aid Smoking Cessation. 2018.	Wrong or missing intervention
Nct, Brown U, National Cancer I. Quit Smoking With Nicotine Patch and Exercise/Health Education (Females Only). 2012.	Linked to excluded study

Nct, Brown U. Improving Smoking Cessation Outcomes in Heavy Drinkers - 1. 2003.	Linked to excluded study
Nct, Brown U. Targeted Pharmacological and Behavioral Treatments for Smoking in Schizophrenia Study 1. 2018.	Ongoing study
Nct, California State University SM, National Institute of General Medical S. Effects Among Smokers Who Use and Do Not Use E-Cigarettes. 2018.	Ongoing study
Nct, Case Comprehensive Cancer C. Path2Quit Tobacco Intervention For African Americans. 2018.	Ongoing study
Nct, Center HLMC, Research I, James, Esther King Biomedical Research P, University of M. Smoking-cessation: A Spanish-Language Clinical Trial. 2015.	Ongoing study
Nct, Center HLMC, Research I, National Cancer I, Florida Department of H. Spanish-Language Smoking Cessation Trial. 2016.	Wrong or missing intervention
Nct, Center MDAC, National Cancer I. Combined Treatment for Tobacco and Alcohol Use in Puerto Rico. 2007.	Ongoing study
Nct, Center MDAC, National Cancer I. Smoking Cessation for HIV/AIDS Patients. 2004.	Wrong or missing intervention
Nct, Center MDAC, National Cancer I. Tailored Smoking Cessation Intervention in Promoting Sexual and Gender Minority Smokers to Quit Smoking. 2018.	Participants received same drug
Nct, Center MDAC, National Institutes of H, The University of Texas Health Science Center H, National Cancer I. Youth Perceptions of Tobacco. 2013.	Wrong or missing intervention
Nct, Center MDAC. Combined Treatment for Tobacco and Alcohol Use. 2007.	Study withdrawn
Nct, Central South U, Pfizer. Boosting Chinese Healthcare Service Providers' Utilization of Smoking Cessation Interventions by 'WeChat WeQuit' Program. 2019.	Ongoing study
Nct, Centre for A, Mental H. Electronic Cigarettes in Daily Dependent Smokers. 2014.	Wrong study design
Nct, Clinica Universidad de Navarra UdN, University of N. Intervention Study for Smoking Cessation in Spanish College Students. 2013.	Wrong or missing intervention
Nct, Company RAIS, Inflamax Research I, Davita Clinical R. CSD170301: An Unblinded, Parallel, Randomized Study to Assess Nicotine Uptake in Smokers From Four Different Electronic Cigarettes. 2017.	Ongoing study
Nct, Company RJRV, Celerion, Company RAIS. Comparison of Nicotine Plasma Concentrations and Subjective Effects for Three Electronic Cigarettes vs Combustible Cigarettes and Nicotine Gum. 2014.	Linked to excluded study
Nct, Company RJRV, Davita Clinical R, Company RAIS. Assess Biomarkers of Tobacco Exposure and Nicotine Pharmacokinetics in Smokers After a 5-Day In-Clinic Confinement Switch to an Electronic Cigarette or Nicotine Gum. 2014.	Ongoing study
Nct, Duke U, National Institute on Drug A. The Role of Nicotine Dose and Route of Delivery in Affecting Adoption of E-cigarettes and Reducing Exposure to Toxic Combustion Products. 2018.	Ongoing study

Nct, Duke U. Perioperative Smoking Cessation Interventions. 2017.	Study terminated
Nct, Duke U. Project Q Pilot: Smoking Cessation for Light Smokers. 2018.	Wrong or missing intervention
Nct, Duke U. Reactions to Warnings on Hookah Ads. 2018.	Wrong or missing intervention
Nct, East Carolina U. Mating-EFT Smoking Cessation Intervention. 2019.	Wrong or missing intervention
Nct, Edward Ellerbeck MM, National Cancer I, University of Kansas Medical C. Smoking Cessation in Rural Hospitals. 2010.	Awaiting classification
Nct, Elliott Crouser MD, American Thoracic S, Ohio State U. Study of Nicotine Patches in Patients With Sarcoidosis. 2008.	Study terminated
Nct, Emory U, John EFIC. Smoke-free Air Coalitions in Georgia and Armenia. 2018.	Wrong or missing intervention
Nct, European Institute of O. Early Smoking Reduction or Cessation by Means of no Nicotine Electronic Cigarette Added to Standard Counselling. 2011.	Ongoing study
Nct, Fertin Pharma AS. Comparison of Pharmacokinetic Profiles of 3 New Nicotine Gum Formulations With a Reference Gum. 2014.	Wrong study design
Nct, Fordham U, Albert Einstein College of Medicine I, University of Texas at A, William Marsh Rice U. Smoking Cessation Intervention for Smokers With Depression Receiving Outpatient Psychiatric Treatment. 2015.	Wrong or missing intervention
Nct, Fundación Pública Andaluza para la gestión de la Investigación en S, European C, Andaluz Health S. So-Lo-Mo Intervention Applied to the Smoking Cessation Process. 2016.	Wrong study design
Nct, Glasgow NHSG, Clyde. The Cardiovascular Impacts of Electronic Cigarettes in Comparison to the Use of Nicotine Replacement Patches. 2016.	Ongoing study
Nct, GlaxoSmithKline. Nicotine Mouth Film for Craving Relief. 2012.	Wrong comparator
Nct, GlaxoSmithKline. Oral Tolerability Of Two Nicotine Dosage Forms. 2007.	Wrong comparator
Nct, Hackensack Meridian H, Lombardi Comprehensive Cancer C, Children's Mercy Hospital Kansas C, University of K. Mobile Cessation Support for Latino Smokers. 2018.	Wrong study design
Nct, Health UC, National Institute on Alcohol A, Alcoholism. Smoking Cessation in Substance Abuse Treatment Patients: A Feasibility Study. 2004.	Linked to excluded study
Nct, Heidelberg U, Pfizer. Brief Intervention Study for Quitting Smoking. 2011.	Ongoing study

Nct, Heinrich-Heine University D, German Research F. Social Cognition,Attentional Network and Nicotine Drug Dependency - A Pharmacological Clinical Trail. 2008.	Linked to excluded study
Nct, Hospital Authority HK, The Hong Kong Research Grants C. A Randomized Controlled Trial of a Nurse Delivered Stage-matched Smoking Cessation Intervention to Promote Heart Health of Cardiac Patients. 2002.	Linked to excluded study
Nct, Hospital Authority HK, The University of Hong K. A Randomized Controlled Trial on Smoking Cessation and Adherence Intervention on Patients With Erectile Dysfunction. 2004.	Linked to excluded study
Nct, Hospital Authority HK, The University of Hong K. Smoking Reduction Intervention for Smokers Not Willing to Quit Smoking: a Randomised Control Trial. 2004.	Awaiting classification
Nct, Hospital Mutua de T. Training Programme in Physical Activity. 2014.	Wrong population
Nct, Hospital Universitari Vall d'Hebron Research I, Clínica G, Hospital Sant R, Hospital de Sant P. Varenicline for Smoking Cessation in Hospitalized Patients With Psychiatric Disorders. 2019.	Ongoing study
Nct, Indiana U, National Institute on Alcohol A, Alcoholism. Reactions to E-cigs and Alcohol. 2017.	Wrong study design
Nct, Institute of P, Neurology W. Smoking Cessation Interventions in Stroke Patients. 2012.	Ongoing study
Nct, Janssen Pharmaceutical KK. A Study to Assess the Nicotine Pharmacokinetics, Tolerability and Safety With a New Oral Nicotine Replacement Product in Healthy Japanese Smokers. 2017.	Wrong study design
Nct, Jeil Pharmaceutical Co L. Compare the Pharmacokinetic Characteristics and the Safety Between Zeropix and Champix in Healthy Adult Male Subjects. 2017.	Wrong study design
Nct, Johns Hopkins U. Cultural Adaptation and Piloting of a Smoking Cessation Intervention for Smokers With HIV. 2018.	Wrong or missing intervention
Nct, Jordi Gol i Gurina F, Preventive S, Health Promotion Research N. ISTAPS: A Stepped Primary Care Smoking Cessation Intervention. 2003.	Linked to excluded study
Nct, Juul Labs I. Clinical Study Comparing 7 ENDS Products and 1 Combustible Cigarette Using 2 Delivery Methods. 2018.	Wrong study design
Nct, Juul Labs I. United States Pre-Market Tobacco Application Pharmacokinetics. 2018.	Wrong study design
Nct, Kaiser P, Oregon H, Science U, Legacy Health S, National Heart L, et al. Cessation Service Use and Effectiveness for Hospitalized Smokers. 2011.	Wrong study design
Nct, Karolinska I, Stockholm County Council S. Smoking Cessation in Primary Health Care Patients With Obstructive Pulmonary Disease. 2010.	Ongoing study
Nct, Karolinska I, Stockholm County Council S. Smoking Cessation Intervention in Acute Orthopedic Surgery. 2004.	Ongoing study

Nct, Kırıkkale U. Synbiotic as an Adjunct to Mechanical Treatment of Gingivitis. 2014.	Wrong or missing intervention
Nct, London South Bank U, Lambeth NHS, Southwark NHS, Allen Carr E. Study Comparing Two Stop Smoking Intervention Efficacies. 2017.	Wrong or missing intervention
Nct, Lorillard Tobacco C. Characterization of Nicotine Exposure and Urge-to-Smoke Following Use of Electronic and Conventional Cigarettes. 2014.	Linked to excluded study
Nct, Massachusetts General H, Bay Cove Human S, Brigham, Women's H, University of Massachusetts A, et al. Integrated Smoking Cessation Treatment for Smokers With Serious Mental Illness. 2016.	Ongoing study
Nct, Massachusetts General H, Partners HealthCare I. Comparative Effectiveness of Two Tobacco Cessation Interventions for Employees of Partners HealthCare, Inc. 2015.	Ongoing study
Nct, Massachusetts General H, Robert Wood Johnson F. Pilot Study of Bupropion for Smoking Cessation in Postpartum Non-breastfeeding Women. 2005.	Study terminated
Nct, Massachusetts General H. Effects of Electronic Cigarettes in Cigarette Smokers With Mild to Moderate COPD. 2019.	Ongoing study
Nct, Mayo C, National Institute on Alcohol A, Alcoholism. Behavioral Counseling for Alcohol Dependent Smokers (Nicotine Patch). 1999.	Participants received same drug
Nct, Mayo C. Brief Pain and Smoking Cessation Intervention in Adults With Chronic Pain. 2015.	Wrong or missing intervention
Nct, McNeil AB, Johnson, Johnson C, Personal Products W. Efficacy and Safety Following Use of a Novel Nicotine Replacement Therapy. 2009.	Awaiting classification
Nct, McNeil AB, Johnson, Johnson C, Personal Products W. Smoking Reduction or Cessation With Nicotine Replacement Therapy. 1999.	Awaiting classification
Nct, McNeil Consumer Healthcare Division of McNeil-Ppc I, McNeil AB, GlaxoSmithKline, Johnson, Johnson C, et al. Study to Determine if a New Nicotine Replacement Therapy Can Safely Help Smokers to Quit Smoking. 2011.	Study terminated
Nct, Medical University of South C, National Cancer I. Clinical Outcomes of a Nationwide, Naturalistic E-Cig Trial (CONNECT). 2018.	Ongoing study
Nct, Medical University of South C, Virginia Tech Carilion School of M, Research I. QuitFast: Evaluating Transcranial Magnetic Stimulation as a Tool to Reduce Smoking Directly Following a Quit Attempt. 2018.	Wrong or missing intervention
Nct, Medical University of South C. Impact of Exclusive Use of Low Nicotine Cigarettes on Compensatory Smoking. 2018.	Wrong or missing intervention
Nct, Medical University of South C. The Effects of Clavulanic Acid on Tobacco Use Disorder. 2019.	Wrong or missing intervention

Nct, Memorial University of N. Evaluation of an Intensive Individualized Smoking Cessation Program Delivered by Pharmacists. 2018.	Wrong or missing intervention
Nct, Miami VAHS. Adverse Airway Effects From Tobacco and E-Cigarettes. 2016.	Ongoing study
Nct, Milton SHMC, National Institute on Drug A. Effects of Random Nicotine Delivery on Smoking Cessation. 2019.	Ongoing study
Nct, Minneapolis Veterans Affairs Medical C, National Institute on Alcohol A, Alcoholism. Timing of Smoking Intervention in Alcohol Treatment (Nicotine Patch). 1997.	Study terminated
Nct, Montefiore Medical C, Albert Einstein College of Medicine I, Yale U. A Trial to Test the Efficacy of a Tailored Intensive Smoking Cessation Intervention in Persons With HIV (PWHs). 2009.	Linked to excluded study
Nct, Montefiore Medical C. Effect of Electronic Cigarette Use on Cough Reflex Sensitivity. 2014.	Linked to excluded study
Nct, National Institute on Alcohol A, Alcoholism. Bupropion as a Smoking Cessation Aid in Alcoholics. 2002.	Awaiting classification
Nct, National Institute on Drug A, National Institutes of Health Clinical C. State and Trait Mediated Response to TMS in Substance Use Disorder. 2019.	Wrong or missing intervention
Nct, National Institute on Drug A. Effectiveness of the Nicotine Patch in Suppressing Nicotine Withdrawal Symptoms in Women Versus Men - 1.	Linked to excluded study
Nct, New York State Psychiatric I, National Institute on Drug A. Effects of Varenicline on Cigarette Self Administration. 2008.	Wrong study design
Nct, New York State Psychiatric I, National Institute on Drug A. Electronic Cigarettes or Nicotine Inhaler for Smoking Cessation. 2013.	Study withdrawn
Nct, New York State Psychiatric I. Effects of Nicotine Replacement and Repeated Cue Exposure on Cigarette Craving. 2009.	Linked to excluded study
Nct, New York University School of M, National Heart L, Blood I. Effectiveness of Smoking-cessation Interventions for Urban Hospital Patients. 2011.	Linked to excluded study
Nct, Norwegian Institute of Public H, Norwegian Cancer S, Helse Stavanger HF, Oslo University H, Sorlandet Hospital HF, et al. Smoking Cessation in Cancer Treatment. 2017.	Ongoing study
Nct, Odense University H, Danish Heart F, Hjernesagen, Funen C. Smoking Cessation Intervention After Stroke or Transient Ischaemic Attack. 2005.	Linked to excluded study

Nct, Ohio State U, National Cancer I. Tobacco Cessation Interventions With Ohio Appalachian Smokers. 2010.	Linked to excluded study
Nct, Oregon Center for Applied Science I. Pharmacological Aids for Interactive Smoking Cessation. 2001.	Wrong or missing intervention
Nct, Oregon Research Behavioral Intervention Strategies I, National Institute on Drug A. Expanding the Click City Tobacco Prevention Program to Include E-cigarettes and Other Novel Tobacco Products. 2018.	Wrong population
Nct, Ottawa Heart Institute Research C. Integrating the "Ottawa Model" for Smoking Cessation Into Primary Care Practice. 2012.	Wrong or missing intervention
Nct, Ottawa Hospital Research I. E-Cigarette Inner City RCT. 2018.	Ongoing study
Nct, Pacific Institute for R, Evaluation, National Institute on Alcohol A, Alcoholism, The Scripps Research I. Strategies: Motivational Interviewing/Psychoeducation. 2018.	Wrong or missing intervention
Nct, Personal Improvement Computer S, National Cancer I. Computer-Assisted Scheduling of Nicotine Inhaler Use in Participants Who Plan to Stop Smoking. 2001.	Participants received same drug
Nct, Peter G, National Institute on Drug A, University of P. Combining a Smoke Ending Aid With Behavioral Treatment - 1. 1999.	Wrong or missing intervention
Nct, Pfizer. A 6-Week Multicenter Trial Of Varenicline Tartrate For Cognitive Impairment In Subjects With Schizophrenia. 2009.	Study withdrawn
Nct, Pfizer. A Single Dose Study Investigating the Absorption and Elimination as Well as the Tolerability of Varenicline Transdermal Delivery System (e.g., a Patch) as Compared to Oral Varenicline in Adult Smokers. 2008.	Awaiting classification
Nct, Pfizer. Impact Of Smoking Cessation Treatment Reimbursement On The Quit Rates In Smokers Motivated To Quit. 2009.	Linked to excluded study
Nct, Pfizer. Multiple Dose, Dose Escalation Study of Varenicline Controlled Release Formulation in Adult Smokers. 2007.	Awaiting classification
Nct, Pharmaceutical Productions I, National Institute on Drug A. Pilot Pharmacokinetic Study of Investigational Sublingual Nicotine Tablet Versus COMMIT Nicotine Lozenge. 2013.	Wrong study design
Nct, Pierre Fabre M. Nicotine Patch Safety Study. 2012.	Awaiting classification
Nct, Queen Mary University of L. Effect of the Electronic Cigarette on Withdrawal Symptoms. 2012.	Wrong study design

Nct, Rebecca A, Novo Nordisk AS, University of P. Daily Liraglutide for Nicotine Dependence. 2018.	Wrong or missing intervention
Nct, Research VAOo, Development, Fred Hutchinson Cancer Research C. Promoting Smoking Cessation in Lung Cancer Screening Through Proactive Treatment. 2018.	Ongoing study
Nct, Rhode Island H. Smoking Cessation Intervention for Diabetic Patients. 2011.	Participants received same drug
Nct, Roswell Park Cancer I, National Cancer I, National Institute on Drug A. Nicotine Delivery From Novel Non-Tobacco Electronic System in Smokers. 2016.	Ongoing study
Nct, Rutgers TSUoNJ. E-cigarette vs. Nicotine Inhaler Comparisons. 2012.	Wrong study design
Nct, Saint-Joseph U, Hotel Dieu de France H. Efficacy of Smoking Cessation Intervention for Head and Neck Cancer Patients. 2015.	Ongoing study
Nct, Saluda Medical Americas I, Saluda Medical Pty L. Safety and Efficacy Study of the Evoke™ SCS System With Feedback vs. Conventional Stimulation. 2017.	Wrong or missing intervention
Nct, Santa Casa Medical S. TNS for Treating Nicotine Craving. 2014.	Ongoing study
Nct, Shanghai 6th People's H, University of North Carolina CH, Asian Center for Health E, Shanghai Municipal Commission of H, Family P, et al. Peer Support To Enhance The Shanghai Integration Model Of Diabetes Care: Extension & Dissemination. 2017.	Wrong population
Nct, Sir Mortimer BDJGH, Canadian Tobacco Control Research I. The Safety of Transdermal Nicotine Immediately Following an Acute Coronary Syndrome. 2005.	Study terminated
Nct, Stanford U. Transnasal Humidified Rapid-Insufflation Ventilatory Exchange (THRIVE) Use in Pediatric Procedures. 2018.	Wrong or missing intervention
Nct, State University of New York - Upstate Medical U, National Alliance for Research on S, Depression. Varenicline Treatment in Alcohol and Nicotine Dependent Patients With Schizophrenia. 2008.	Study terminated
Nct, State University of New York at B, National Cancer I, Pfizer. EVarQuit: Extended Pre-quit Varenicline to Assist in Quitting Smoking. 2017.	Ongoing study
Nct, Swedish Match AB. Serbian Smoking Reduction/Cessation Trial (2SRT). 2008.	Linked to excluded study
Nct, Tata Memorial H. Evaluation of Efficacy of Different Methods of Tobacco Cessation Interventions Among BEST Employees in Mumbai. 2015.	Ongoing study
Nct, Tata Memorial H. Medium Vs Low Intensity Intervention in Tobacco Cessation and Control in Zari Workers. 2013.	Ongoing study
Nct, The Miriam H, Brown U, University of California B, International RTI, Rhode Island H, et al. Sustaining Smoking Cessation in Smokers With Kids With Asthma. 2005.	Participants received same drug
Nct, The University of Texas Health Science Center H. Medical Nutrition Therapy for Medically Complex Infants in the Pediatric Outpatient Setting. 2018.	Wrong population

Nct, Universita degli Studi di C. Antismoking Effects of Electronic Cigarettes in Subjects With Schizophrenia and Their Potential Influence on Cognitive Functioning. 2018.	Ongoing study
Nct, Universita degli Studi di C. High Cessation Rates in Smokers Using Personal Vaporizers. 2013.	Linked to excluded study
Nct, University Health Network T, Pfizer. Smoking Cessation Program in the Preadmission Clinic. 2008.	Ongoing study
Nct, University Hospital BS. Smoking Cessation in Alcoholics. 2007.	Linked to excluded study
Nct, University Hospital GC, Danish Research Foundation FSSCD. Smoking Cessation With Smokeless Tobacco, Nicotine Tablets and Nicotine Patches. 2006.	Ongoing study
Nct, University Hospital GC, Pfizer. Varenicline for Long-term Nicotine Replacement Therapy (NRT) Users. 2009.	Linked to excluded study
Nct, University Hospital GC. Nurse Smoking Cessation of Patients With Chronic Obstructive Pulmonary Disease (COPD) With Nicotine Replacement Therapy (NRT) and Behavioral Support. 2002.	Study terminated
Nct, University Hospital SF. Nicotine Pharmacokinetic in Blood and Analysis of Exhaled Breath After E-cigarette Smoking. 2018.	Wrong study design
Nct, University Hospitals Cleveland Medical C, Government Calicut Medical C, Paramukhswami Medical C, Science PSGIoM, Research. Medical Student Counseling for Hospitalized Patients Addicted to Tobacco. 2018.	Participants received same drug
Nct, University Medicine G, German Federal Ministry of E, Research, German Research F. Smoking Interventions in General Medical Practices. 2004.	Ongoing study
Nct, University Medicine G, German Federal Ministry of E, Research. Smoking Cessation and Relapse Prevention in Women Postpartum. 2001.	Wrong or missing intervention
Nct, University of Alabama at B. Balanced Placebo Design With Varenicline: Pharmacological and Expectancy Effects on Medication Adherence. 2018.	Ongoing study
Nct, University of Alabama at B. NRT Sampling and Selection to Increase Medication Adherence. 2018.	Participants received same drug
Nct, University of B, University of L, University of Geneva S. The ESTxENDS Trial- Electronic Nicotine Delivery Systems (ENDS/Vaporizer/E-cigarette) as an Aid for Smoking Cessation. 2018.	Ongoing study
Nct, University of C, National Institute on Drug A, Ortho-McNeil Janssen Scientific Affairs LLC. Smoking Cessation for Adults With Attention Deficit Hyperactivity Disorder (ADHD). 2005.	Wrong or missing intervention
Nct, University of California SF, Food, Drug A, National Institutes of H, National Cancer I. Cardiovascular Assessment of the Effects of Tobacco and Nicotine Delivery Products. 2013.	Ongoing study

Nct, University of L, Pharmacia. Nicotine Substitute Prescribed at Hourly Intake or ad Libitum for Heavy Smokers Willing to Quit. 2000.	Linked to excluded study
Nct, University of M, National Institute of Mental H. An Integrated Approach to Smoking Cessation in Severe Mental Illness (SMI). 2009.	Linked to excluded study
Nct, University of M, Prevent Cancer F. Randomized Trial Assessing the Effectiveness of a Pharmacist-Delivered Program for Smoking Cessation. 2005.	Linked to excluded study
Nct, University of M. Smoking Cessation After Acute Coronary Syndrome. 2015.	Ongoing study
Nct, University of M. Smoking Cessation Interventions for People Living With HIV in Nairobi, Kenya. 2019.	Ongoing study
Nct, University of Minnesota C, Translational Science I, National Institute on Drug A. Medicinal Nicotine for Preventing Stress Induced Craving and Withdrawal Symptoms. 2011.	Linked to excluded study
Nct, University of Minnesota C, Translational Science I. Medicinal Nicotine for Preventing Cue Induced Craving. 2013.	Linked to excluded study
Nct, University of North Carolina CH, National Cancer I, Emory U, The University of Texas Health Science Center H, Washington University School of M. Brief Intervention to Create Smoke-Free Homes Policies in Low-Income Households: North Carolina Effectiveness Trial. 2013.	Wrong or missing intervention
Nct, University of O, Lapland Central Hospital Rovaniemi F. Efficacy and Safety of E-cigarettes for Smoking Cessation in Middle-aged Heavy Smokers. 2018.	Ongoing study
Nct, University of O, Pfizer. Effectiveness of Combination Varenicline and Oral Nicotine Replacement Therapy. 2019.	Ongoing study
Nct, University of O. Project Phoenix: Pilot Intervention for Smokers Who Are Not Ready to Quit Smoking. 2018.	Ongoing study
Nct, University of O. Smoking Cessation Treatment for Depressed Smokers. 2015.	Wrong or missing intervention
Nct, University of O. Smoking Cessation Treatment for Substance Use Dependents. 2018.	Wrong or missing intervention
Nct, University of P, AstraZeneca. Effects of Chantix on Relapse Prevention for Smoking Cessation. 2006.	Wrong study design
Nct, University of P, National Institute of Mental H, Johns Hopkins Bloomberg School of Public H. Impact of Representative Payee Services on ART Adherence Among Marginalized People Living With HIV/AIDS. 2017.	Wrong population
Nct, University of Paris 5 - Rene D, Assistance Publique - Hôpitaux de P. Effect of Daily Nicotine Patch Application on Mechanical Ventilation Weaning in Smoking Patients. 2010.	Ongoing study

Nct, University of Roma La S. Simultaneous Coronary Artery Evaluation and Lung Cancer CT Screening. 2018.	Wrong or missing intervention
Nct, University of South C, Academy of N, Dietetics. The Plant-Based and Soul-Full Study (PASS). 2018.	Wrong population
Nct, University of Southern C, Illinois Institute of T. Effects of Nicotine in Tobacco Smokers. 2018.	Ongoing study
Nct, University of Southern C, University of California LA, University of California SD. High Dose Bupropion for Smoking Cessation. 2018.	Ongoing study
Nct, University of Southern D, Danish Cancer S, National Board of Health D. Comparison of Four Different Smoking Cessation Programmes. 2011.	Ongoing study
Nct, University of T, National Heart L, Blood I. Nicotine Patch and Bupropion to Reduce Smoking Rates in Younger, Low-Income, and Minority Individuals. 2004.	Awaiting classification
Nct, University of Texas at A. Smoking Cessation and Sexual Health in Men. 2008.	Linked to excluded study
Nct, University of V, Vanderbilt University Medical C. The Nicotinic Cholinergic System and Cognitive Aging. 2016.	Participants received same drug
Nct, University of W, National Institute on Drug A. Personalized Health Assessment Related to Medications (Project PHARM). 2017.	Wrong or missing intervention
Nct, University of Wisconsin M, Dean Health S, Mercy Health S, Aurora Health C, National Cancer I. Identifying Optimal Smoking Cessation Intervention Components (Cessation). 2010.	Linked to excluded study
Nct, University of Wisconsin M, Institute NC. A Quit Smoking Study Using Smartphones. 2014.	Participants received same drug
Nct, University of Wisconsin M, University of Illinois at C, Dean Health S, Mercy Health S, Aurora Health C, et al. Identifying Treatments to Motivate Smokers to Quit. 2010.	Linked to excluded study
Nct, Vanderbilt U. Smoking Cessation Intervention in the Emergency Department (ED). 2010.	Wrong or missing intervention
Nct, Washington University School of M, National Institutes of H, National Cancer I. Helping Poor Smokers Quit. 2017.	Wrong or missing intervention
Nct, Western University C. Aerobic Exercise and Inhibitory Control. 2018.	Wrong or missing intervention

Nct, Yale U, National Institute on Drug A. Framing Messages for Smoking Cessation With Bupropion - 6. 2000.	Linked to excluded study
Nct, Yale U, National Institutes of H. A Smoking Cessation Treatment for Adult Women. 2011.	Wrong or missing intervention
Nct, Yale U, Research VAOo, Development. Treatment for Nicotine Addiction in Women. 2015.	Ongoing study
Nct, Yale U. A Study to Help People Quit Smoking. 2013.	Wrong or missing intervention
Nct, Yale U. The Effects of Menthol as Delivered by an Electronic Cigarette on the Desirability of Nicotine in Tobacco Users. 2014.	Ongoing study
Nct, Yu-Li H, National Health Research Institutes T, Department of Health EYRO. The Pharmacogenetic Study , Readiness to Change, and Pharmacological Intervention for Smoking Cessation in Schizophrenia. 2006.	Awaiting classification
nct. A Pilot Drug Trial That Evaluates the Whitening Potential of Nicotine Gum. https://clinicaltrials.gov/show/nct00174499 . 2005.	Awaiting classification
Nct. A Smoking Intervention Study Using Scheduled Gradual Reduction With Varenicline to Help With Cessation. https://ClinicalTrials.gov/show/NCT01772641 ; 2013.	Study terminated
Nct. A Study to Evaluate the Safety of Electronic Vapor Products for 2 Years. https://ClinicalTrials.gov/show/NCT02143310 ; 2014.	Participants received same drug
Nct. A Study to Evaluate the Safety Profile of an e-Vapour Product. https://ClinicalTrials.gov/show/NCT02029196 ; 2013.	Ongoing study
Nct. A Trial of E-cigarettes in Current Cigarette Smokers. https://ClinicalTrials.gov/show/NCT02357173 ; 2014.	Ongoing study
Nct. Advancing Tobacco Use Treatment for African American Smokers. https://ClinicalTrials.gov/show/NCT02360631 ; 2015.	Ongoing study
Nct. Changes in Psychological and Somatic Parameters Associated With Changing From Smoking to E-cigarettes. https://ClinicalTrials.gov/show/NCT02635620 ; 2015.	Ongoing study
Nct. Comparison of Varenicline and Placebo for Smoking Cessation in Schizophrenia. https://ClinicalTrials.gov/show/NCT00554840 ; 2007.	Ongoing study
Nct. Contingency Management and Pharmacotherapy for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00683280 ; 2008.	Participants received same drug
Nct. Dissemination of a Tailored Tobacco Quitline for Rural Veteran Smokers. https://ClinicalTrials.gov/show/NCT01892813 ; 2013.	Ongoing study
Nct. Early In-hospital Initiation of Pharmacotherapy for Smoking Cessation, Patients After ACS. https://ClinicalTrials.gov/show/NCT02106637 ; 2014.	Ongoing study
Nct. E-Cigarettes: Dynamic Patterns of Use and Health Effects. https://ClinicalTrials.gov/show/NCT02527980 ; 2015.	Ongoing study
Nct. Effects of Varenicline on Smoking Lapse in Smokers With and Without Schizophrenia. https://ClinicalTrials.gov/show/NCT01850953 ; 2013.	Ongoing study
nct. Efficacy and Safety of an Electronic Nicotine Delivery Device (E-Cigarette) Without Nicotine Cartridges. https://clinicaltrials.gov/show/nct01194583 . 2010.	Wrong study design

Nct. Efficacy and Safety of Dianicline Treatment as an Aid to Smoking Cessation in Cigarette Smokers (AMERIDIAN). https://ClinicalTrials.gov/show/NCT00387946 ; 2006.	Wrong or missing intervention
Nct. Efficacy And Safety Of Smoking Cessation With Varenicline Tartrate In Diabetic Smokers: (DIASMOKE). https://ClinicalTrials.gov/show/NCT01387425 ; 2011.	Ongoing study
Nct. Efficacy of Varenicline on Smoking Cessation at the Acute Phase of an Exacerbation of Chronic Obstructive Pulmonary Disease. https://ClinicalTrials.gov/show/NCT01694732 ; 2012.	Ongoing study
Nct. Electronic Cigarettes (E-cigarettes) as a Harm Reduction Strategy. https://ClinicalTrials.gov/show/NCT02628964 ; 2015.	Ongoing study
nct. Electronic Cigarettes and Reactivity to Smoking Cues. https://clinicaltrials.gov/show/nct01782599 . 2013.	Ongoing study
Nct. Electronic Cigarettes in Smokers With Mental Illness. https://ClinicalTrials.gov/show/NCT02212041 ; 2014.	Wrong comparator
Nct. Electronic Nicotine Delivery Systems as a Smoking Cessation Treatment. https://ClinicalTrials.gov/show/NCT02487953 ; 2015.	Ongoing study
nct. Evaluating the Acute Effects of Electronic Nicotine Delivery Devices Marketed to Smokers. https://clinicaltrials.gov/show/nct00932295 . 2009.	Wrong study design
Nct. Evaluating the Efficacy of E-Cigarette Use for Smoking Cessation (E3) Trial. https://ClinicalTrials.gov/show/NCT02417467 ; 2015.	Ongoing study
Nct. Evaluation of Appeal and Impact of E-Cigarettes Among Chronic Smokers With Smoking-Related Cancers. https://ClinicalTrials.gov/show/NCT02648178 ; 2016.	Ongoing study
nct. Examining the Effectiveness of the Nicotine Patch in Male and Female Smokers - 2. https://clinicaltrials.gov/show/nct00135746 . 2005.	Wrong study design
Nct. Exercise or Relaxation for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00921388 ; 2009.	Wrong comparator
Nct. Extended Treatment for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01330043 ; 2010.	Ongoing study
Nct. Genetically Informed Smoking Cessation Trial. https://ClinicalTrials.gov/show/NCT02351167 ; 2015.	Ongoing study
Nct. Improving Varenicline Adherence and Outcomes in Homeless Smokers. https://ClinicalTrials.gov/show/NCT00786149 ; 2008.	Participants received same drug
Nct. Maintaining Nonsmoking. https://ClinicalTrials.gov/show/NCT01162239 ; 2010.	Participants received same drug
Nct. Methadone Maintenance Treatment and Smoking Cessation. https://ClinicalTrials.gov/show/NCT00906386 ; 2009.	Ongoing study
nct. Nicotine Patch for Nicotine Dependence in Individuals With Schizophrenia or Schizoaffective Disorder - 1. https://clinicaltrials.gov/show/nct00046813 . 2002.	Study terminated
Nct. Penn State TXT2Quit Study. https://ClinicalTrials.gov/show/NCT02367391 ; 2015.	Participants received same drug
Nct. Personal Vaporizer vs Cigalike. https://ClinicalTrials.gov/show/NCT02398487 ; 2014.	Ongoing study

nct. Pharmacologic Intervention for Postcessation Weight Gain. https://clinicaltrials.gov/show/nct00005704 . 2000.	Wrong or missing intervention
Nct. Pilot Study of Nicotine Nasal Spray and Varenicline on Smoking in Methadone-Maintained Patients. https://ClinicalTrials.gov/show/NCT02147132 ; 2014.	Wrong study design
Nct. Reward Sensitivity and Pharmacotherapy for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02162849 ; 2014.	Ongoing study
Nct. Safety and Efficacy of the Novel Selective Nicotinic Receptor Partial Agonist, CHANTIX (Varenicline) in Patients With Acute Coronary Syndrome. https://ClinicalTrials.gov/show/NCT01170338 ; 2010.	Ongoing study
Nct. Short Term Effects of Electronic Cigarettes in Tobacco Dependent Adults. https://ClinicalTrials.gov/show/NCT02498145 ; 2015.	Ongoing study
Nct. Smoking Cessation in Women With Gynecological Conditions. https://ClinicalTrials.gov/show/NCT01989923 ; 2013.	Ongoing study
Nct. Smoking Cessation Program in the Preadmission Clinic: The Use of a Teachable Moment. https://ClinicalTrials.gov/show/NCT01243203 ; 2010.	Ongoing study
Nct. Smoking Cessation Strategies in Community Cancer Programs for Lung and Head and Neck Cancer Patients. https://ClinicalTrials.gov/show/NCT02048917 ; 2014.	Ongoing study
Nct. Smoking Cessation Study In Healthy Adolescent Smokers. https://ClinicalTrials.gov/show/NCT01312909 ; 2011.	Wrong population
Nct. Smoking Cessation Treatment for Head and Neck Cancer Patients. https://ClinicalTrials.gov/show/NCT00931021 ; 2009.	Study terminated
Nct. Smoking Habits and Smoking Cessation in Young Adults. https://ClinicalTrials.gov/show/NCT01531049 ; 2012.	Ongoing study
Nct. Smoking Study With Behavioral Therapy for Hypertensive Patients. https://ClinicalTrials.gov/show/NCT00879177 ; 2009.	Wrong comparator
Nct. Study of Varenicline (Champix) for Smoking Cessation/Reduction in Patients With Bipolar Disorder. https://ClinicalTrials.gov/show/NCT01093937 ; 2010.	Ongoing study
Nct. Tailored Tobacco Quitline for Rural Veterans. https://ClinicalTrials.gov/show/NCT01592695 ; 2012.	Wrong study design
Nct. The MATCH (Medication Aids for Tobacco Cessation and Health) Study. https://ClinicalTrials.gov/show/NCT02146911 ; 2014.	Ongoing study
Nct. The Moment Study: Mixed Method Ecigarette Study. https://ClinicalTrials.gov/show/NCT02261363 ; 2014.	Wrong comparator
Nct. The Role of Nicotine and Non-Nicotine Alkaloids in E-Cigarette Use and Dependence. https://ClinicalTrials.gov/show/NCT02590393 ; 2015.	Ongoing study
Nct. The Use of Nicotine Patches Together With E-cigarettes (With and Without Nicotine) for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02521662 ; 2015.	Ongoing study
Nct. Tobacco Dependence in Breast Cancer Patients Trial of Varenicline (Chantix). https://ClinicalTrials.gov/show/NCT01532232 ; 2012.	Study terminated
Nct. Treatment of Smoking Among Individuals With PTSD. https://ClinicalTrials.gov/show/NCT00937235 ; 2009.	Wrong comparator
Nct. UK-Czech E-cigarette Study. https://ClinicalTrials.gov/show/NCT01842828 ; 2013.	Ongoing study
Nct. Varenicline and Combined Nicotine Replacement Therapy (NRT) for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02271919 ; 2014.	Ongoing study

Nct. Varenicline for Adolescent Smoking Cessation. https://ClinicalTrials.gov/show/NCT01509547 ; 2011.	Wrong population
Nct. Varenicline for Nicotine Dependence Among Those With HIV/AIDS. https://ClinicalTrials.gov/show/NCT01710137 ; 2012.	Ongoing study
Nct. Varenicline in Drug Treatment. https://ClinicalTrials.gov/show/NCT01286584 ; 2011.	Ongoing study
Nct. Varenicline In-Patient Study. https://ClinicalTrials.gov/show/NCT01413516 ; 2011.	Ongoing study
Nct. Varenicline or Nicotine Patch in Promoting Smoking Cessation Among Current Smokers. https://ClinicalTrials.gov/show/NCT01771627 ; 2013.	Ongoing study
Nelson PR, Chen P, Battista DR, Pillitteri JL, Shiffman S. Randomized Trial to Compare Smoking Cessation Rates of Snus, with and without Smokeless Tobacco Health-Related Information, and a Nicotine Lozenge. <i>Nicotine & Tobacco Research</i> . 2018;24:24.	Wrong comparator
Nerin I, Beamonte A, Gargallo P, Jimenez-Muro A, Marqueta A. Weight gain and anxiety levels in recent ex-smokers. [Spanish]. <i>Archivos de Bronconeumologia</i> . 2007;43(1):9-15.	Wrong study design
Neuropsychiatric safety of varenicline and bupropion 2016. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/075/CN-01195075/frame.html .	Wrong study design
New York State Psychiatric I, National Institute on Drug A. Effect of Memantine Versus Bupropion on Smoking Relapse in Nicotine-Dependent Individuals - 3. https://ClinicalTrials.gov/show/NCT00136786 ; 2006.	Wrong study design
New York State Psychiatric I, National Institute on Drug A. The Effects of Memantine and Bupropion on Acute, Reinforcing, and Conditioned Effects of Cigarettes - 1. https://ClinicalTrials.gov/show/NCT00136747 ; 2004.	Wrong study design
New York State Psychiatric I. Effects of Nicotine Replacement Therapy and D-cycloserine on Nicotine Treatment Seekers. https://ClinicalTrials.gov/show/NCT01907594 ; 2013.	Wrong comparator
New York U, National Institute on Drug A. Smoking Cessation Treatment With Transdermal Nicotine Replacement Therapy - 1. https://ClinicalTrials.gov/show/NCT00067158 ; 2003.	Participants received same drug
New York University School of M, Lincoln M, Mental Health C. Smoking Cessation Peds Asthma. https://ClinicalTrials.gov/show/NCT02640521 ; 2015.	Study withdrawn
New York University School of M, Pfizer. Study of Varenicline Effects on Cigarette Smoking Reward and Craving During a Model of Brief Quit Attempt. https://ClinicalTrials.gov/show/NCT00571805 ; 2007.	Study withdrawn
New York University School of M, Robert Wood Johnson F, System VANYHH. Financial Incentives for Smoking Treatment. https://ClinicalTrials.gov/show/NCT02506829 ; 2015.	Participants received same drug
New York University School of M. Effect of Electronic Cigarettes on Platelets, Endothelium and Inflammation. https://ClinicalTrials.gov/show/NCT02662075 ; 2015.	Study withdrawn
Ng P, Oliverio S, Sobota M. Kick the butt!-a pilot smoking cessation program using group visits in an internal medicine resident clinic. <i>Journal of General Internal Medicine</i> . 2016;1):S832.	Wrong study design
Nguyen L, Mullen C, Saxon L, Chastain L. Assessing rates of tobacco cessation within a multimodal group intervention in a veterans affairs medical center. <i>Journal of the American Geriatrics Society</i> . 2015;63:S85.	Cannot locate full text
Nides MA, Leischow SJ, Bhattar M, Simmons M. Nicotine blood levels and short-term smoking reduction with an electronic nicotine delivery system. 2014;38(2):265-74.	Wrong study design

Nieva G, Comin M, Valero S, Bruguera E. Cigarette dependence and depressive symptoms as predictors of smoking status at five-year follow-up after a workplace smoking cessation program. <i>Addictive Behaviors</i> . 2017;73:9-15.	Wrong study design
Nishio C, Tomioka H, Mamesaya N, Nakao S, Sekiya R, Yamashita S, et al. Health-related quality of life in asthmatic patients who attended a smoking cessation program. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2014;44(no pagination).	Participants received same drug
Nishio C, Tomioka H, Sekiya R, Yamashita S, Kaneda T, Kida Y, et al. Characteristics of mentally ill smokers participating in a smoking cessation program. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2012;40(no pagination).	Participants received same drug
Njoy I, Trials LAC, National Institute on Drug A. Nicotine Pharmacokinetics From Research Electronic Nicotine Delivery System S-TA-U001 in Smokers and E-Cigarette Users. https://ClinicalTrials.gov/show/NCT02792426 ; 2016.	Wrong study design
Nohlert E, Ohrvik J, Helgason AR. Effectiveness of proactive and reactive services at the Swedish National Tobacco Quitline in a randomized trial. <i>Tobacco Induced Diseases</i> . 2014;12 (1) (no pagination)(9).	Wrong or missing intervention
Nohlert E, Ohrvik J, Tegelberg A, Tillgren P, Helgason AR. Long-term follow-up of a high- and a low-intensity smoking cessation intervention in a dental setting- a randomized trial. <i>Bmc Public Health</i> . 2013;13.	Wrong or missing intervention
Nollen N, Ahluwalia JS, Mayo MS, Richter K, Choi WS, Okuyemi KS, et al. A randomized trial of targeted educational materials for smoking cessation in African Americans using transdermal nicotine. <i>Health Education & Behavior</i> . 2007;34(6):911-27.	Participants received same drug
Nollen NL, Cox LS, Mayo MS, Ellerbeck EF, Madhusudhana S, Ahluwalia JS. A randomized clinical trial of counseling and nicotine replacement therapy for treatment of African American non-daily smokers: Design, accrual, and baseline characteristics. <i>Contemporary Clinical Trials</i> . 2018;70:72-82.	Ongoing study
North Shore Medical C. Smoking Cessation After Hospitalization for a Cardiopulmonary Illness. https://ClinicalTrials.gov/show/NCT01791803 ; 2006.	Ongoing study
Northwell H. Comparison of Conservative and Aggressive Smoking Cessation Treatment Strategies in a Vascular Surgery Office Practice. https://ClinicalTrials.gov/show/NCT01850589 ; 2011.	Study terminated
Northwestern U, National Cancer I. Behavioral Activation and Varenicline for Smoking Cessation in Depressed Smokers. https://ClinicalTrials.gov/show/NCT02378714 ; 2015.	Ongoing study
Nunes T, Etchevers MJ, Marin I, Gallego S, Merino Ochoa O, Menchen Viso L, et al. Effectiveness of smoking cessation in Crohn's disease patients. <i>Journal of Crohn's and Colitis</i> . 2012;6:S162.	Cannot locate full text
Oertel W, Muller H, Schade-Brittinger C, Kamp C, Balthasar K, Articus K, et al. The NIC-PD-study-A randomized, placebo-controlled, double-blind, multi-centre trial to assess the disease-modifying potential of transdermal nicotine in early Parkinson's disease in Germany and N. America. <i>Movement disorders Conference: 22nd international congress of parkinson's disease and movement disorders, MDS 2018 Hong kong</i> . 2018;33(Supplement 2):S159.	Wrong population
Ojo-Fati O, John F, Thomas J, Joseph AM, Raymond NC, Cooney NL, et al. Integrating smoking cessation and alcohol use treatment in homeless populations: Study protocol for a randomized controlled trial. <i>Trials</i> . 2015;16 (1) (no pagination)(385).	Participants received same drug
Okuyemi KS, Goldade K, Whembolua GL, Thomas JL, Eischen S, Guo H, et al. Smoking characteristics and comorbidities in the power to quit randomized clinical trial for homeless smokers. <i>Nicotine & Tobacco Research</i> . 2013;15(1):22-8.	Participants received same drug

Okuyemi KS, Goldade K, Whembolua GL, Thomas JL, Eischen S, Sewali B, et al. Motivational interviewing to enhance nicotine patch treatment for smoking cessation among homeless smokers: a randomized controlled trial. <i>Addiction</i> . 2013;108(6):1136-44.	Participants received same drug
Okuyemi KS, Thomas JL, Hall S, Nollen NL, Richter KP, Jeffries SK, et al. Smoking cessation in homeless populations: A pilot clinical trial. <i>Nicotine & Tobacco Research</i> . 2006;8(5):689-99.	Participants received same drug
O'Malley SS, Cooney JL, Krishnan-Sarin S, Dubin JA, McKee SA, Cooney NL, et al. A controlled trial of naltrexone augmentation of nicotine replacement therapy for smoking cessation. <i>Archives of Internal Medicine</i> . 2006;166(6):667-74.	Wrong comparator
O'Malley SS, Wu R, Mayne ST, Jatlow PI. Smoking cessation is followed by increases in serum bilirubin, an endogenous antioxidant associated with lower risk of lung cancer and cardiovascular disease. <i>Nicotine and Tobacco Research</i> . 2014;16 (8) (no pagination)(ntu067).	Participants received same drug
O'Malley SS, Zweben A, Fucito L. Varenicline for the treatment of alcohol dependence in smokers: Can one treatment affect drinking and smoking? <i>Alcoholism: Clinical and Experimental Research</i> . 2016;40:251A.	Ongoing study
Oncken C, Arias AJ, Feinn R, Litt M, Covault J, Sofuoglu M, et al. Topiramate for smoking cessation: a randomized, placebo-controlled pilot study. <i>Nicotine & Tobacco Research</i> . 2014;16(3):288-96.	Wrong or missing intervention
Oncken C, Dornelas E, Greene J, Sankey H, Glasmann A, Feinn R, et al. Nicotine gum for pregnant smokers - A randomized controlled trial. <i>Obstetrics and Gynecology</i> . 2008;112(4):859-67.	Wrong population
Oncken CA, Litt MD, McLaughlin LD, Burki NA. Nicotine Concentrations With Electronic Cigarette Use: Effects of Sex and Flavor. <i>Nicotine & Tobacco Research</i> . 2015;17(4):473-8.	Wrong study design
Onen ZP, Sen E, Eris Gulbay B, Ozturk A, Akkoca Yildiz O, Acican T, et al. [Smoking cessation in patients with cardiopulmonary diseases]. <i>Anadolu Kardiyoloji Dergisi</i> . 2011;11(3):244-9.	Wrong study design
Onen ZP, Sen E, Gulbay BE, Kabalak PA, Yildiz OA, Acican T, et al. Effectiveness of smoking cessation in patients with psychiatric disease. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2012;40(no pagination).	Cannot locate full text
Onizawa S, Taniguchi H, Nozu T, Asano A, Katsura H, Saiki K, et al. [Sustainable efficacy of oral varenicline for smoking cessation]. <i>Nihon Kokyuki Gakkai Zasshi</i> . 2010;48(11):791-6.	Participants received same drug
Ormston R, van der Pol M, Ludbrook A, McConville S, Amos A. quit4u: The effectiveness of combining behavioural support, pharmacotherapy and financial incentives to support smoking cessation. <i>Health Education Research</i> . 2015;30(1):121-33.	Participants received same drug
Oslo University H. Transdermal Nicotine Replacement in Smokers With Acute Aneurysmal Subarachnoid Hemorrhage. https://ClinicalTrials.gov/show/NCT02350335 ; 2011.	Wrong comparator
Ostacher M. Collaborative framework for a multisite clinical trial in bipolar disorder and comorbid substance abuse: varenicline in bipolar smokers. <i>Bipolar Disorders</i> . 2013;15:34-.	Awaiting classification
Ostroff JS, Borderud SP, Li Y, Burkhalter J, Sheffer CE. Electronic cigarette use among cancer patients: Characteristics of e-cigarette users and their smoking cessation outcomes. <i>Psycho-Oncology</i> . 2014;23:139-40.	Wrong comparator
Otero UB, Perez CdA, Szklo M, Esteves GA, dePinho MM, Szklo AS, et al. Randomized clinical trial: effectiveness of the cognitive-behavioral approach and the use of nicotine replacement transdermal patches for smoking cessation among adults in Rio de Janeiro, Brazil. <i>Cadernos de saude publica</i> . 2006;22(2):439-49.	Wrong study design

Ottawa Heart Institute Research C, Heart, Stroke Foundation of O. Efficacy and Cost-Effectiveness of Cost-free Pharmacotherapy for Smoking Cessation for High-risk Smokers With Cerebrovascular Disease. https://ClinicalTrials.gov/show/NCT00962988 ; 2009.	Ongoing study
Ottawa Heart Institute Research C, Heart, Stroke Foundation of O. Self-directed Titrated Transdermal Nicotine Patch Versus Standard Treatment for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01622998 ; 2011.	Ongoing study
Ottawa Heart Institute Research C. Cost Free Pharmacotherapy for Smoking Cessation in Stroke Patients. https://ClinicalTrials.gov/show/NCT01404546 ; 2008.	Ongoing study
Ottawa Hospital Research I, Canadian Institutes of Health R, Network CCHT. The Canadian HIV Quit Smoking Trial: Tackling the Co-morbidities of Depression and Cardiovascular Disease in HIV+ Smokers. https://ClinicalTrials.gov/show/NCT01800019 ; 2014.	Ongoing study
Ozer A, Seydaoglu G. The success rates of smoking cessation methods. American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS. 2015;191(no pagination).	Cannot locate full text
Pacifici R, Pichini S, Graziano S, Pellegrini M, Massaro G, Beatrice F. Successful Nicotine Intake in Medical Assisted Use of E-Cigarettes: A Pilot Study. Int J Environ Res Public Health. 2015;12(7):7638-46.	Wrong study design
Pack QR, Priya A, Lagu T, Pekow P, Rigotti N, Lindenauer P. Smoking cessation therapies are persistently and significantly underutilized among hospitalized smokers with coronary heart disease. Journal of the American College of Cardiology. 2017;69 (11 Supplement 1):1755.	Awaiting classification
Pack QR, Priya A, Lagu TC, Pekow PS, Atreya A, Rigotti NA, et al. Short-Term Safety of Nicotine Replacement in Smokers Hospitalized With Coronary Heart Disease. Journal of the American Heart Association. 2018;7(18).	Awaiting classification
Palamidas A, Gennimata SA, Kaltsakas G, Tsikrika S, Vakali S, Gratziou C, et al. Acute effect of an e-cigarette with and without nicotine on lung function. European Respiratory Journal Conference: European Respiratory Society Annual Congress. 2013;42(no pagination).	Wrong study design
Palamidas A, Gennimata SA, Kaltsakas G, Tsikrika S, Vakali S, Gratziou C, et al. Acute effect of an e-cigarette with and without nicotine on lung function. Tobacco Induced Diseases Conference: 11th Annual Conference of the International Society for the Prevention of Tobacco Induced Diseases, ISPTID. 2013;12(no pagination).	Wrong study design
Paone G, Serpilli M, Girardi E, Conti V, Principe R, Puglisi G, et al. The combination of a smoking cessation programme with rehabilitation increases stop-smoking rate. Journal of Rehabilitation Medicine. 2008;40(8):672-7.	Participants received same drug
Pardell H, Salto E, Ciruela J, Tresserras R, Gascon P, Salleras L. [Smoking cessation program using nicotine patch among physicians and pharmacists of Catalonia: results after 3-12 months]. Anales de Medicina Interna. 1996;13(1):16-20.	Participants received same drug
Partin MR, An LC, Nelson DB, Nugent S, Snyder A, Fu SS, et al. Randomized Trial of an Intervention to Facilitate Recycling for Relapsed Smokers. American Journal of Preventive Medicine. 2006;31(4):293-9.	Participants received same drug
Patten CA, Rummans TA, Croghan IT, Hurt RD, Hays JT. Development of depression during placebo-controlled trials of bupropion for smoking cessation: case reports. Journal of Clinical Psychiatry. 1999;60(7):436-41.	Wrong study design
Pearson JL, Smiley SL, Rubin LF, Anesetti-Rothermel A, Elmasry H, Davis M, et al. The Moment Study: protocol for a mixed method observational cohort study of the Alternative Nicotine Delivery Systems (ANDS) initiation process among adult cigarette smokers. BMJ Open. 2016;6(4):e011717.	Wrong comparator
Peckham E, Man MS, Mitchell N, Li J, Becque T, Knowles S, et al. Smoking cessation intervention for severe mental ill health trial (SCIMITAR): A pilot randomised control trial of the clinical effectiveness and cost-effectiveness of a bespoke smoking cessation service. Health Technology Assessment. 2015;19(25):1-148.	Participants received same drug

Pederson LL, Wanklin JM, Lefcoe NM. THE EFFECTS OF COUNSELING ON SMOKING CESSATION AMONG PATIENTS HOSPITALIZED WITH CHRONIC OBSTRUCTIVE PULMONARY-DISEASE - A RANDOMIZED CLINICAL-TRIAL. <i>International Journal of the Addictions</i> . 1991;26(1):107-19.	Wrong or missing intervention
Pedrol-Clotet E, Deig-Comerma E, Ribell-Bachs M, Vidal-Castell I, Garcia-Rodriguez P, Soler A. Bupropion use for smoking cessation in HIV-infected patients receiving antiretroviral therapy. [Spanish]. <i>Enfermedades Infecciosas y Microbiologia Clinica</i> . 2006;24(8):509-11.	Wrong study design
Penn P, Gallagher SM, Schindler E, Layne WK. Comparing smoking cessation interventions for persons with serious mental illness. <i>Drug and Alcohol Dependence</i> . 2015;146:e68.	Cannot locate full text
Perez-Tortosa S, Roig L, Manresa JM, Martin-Cantera C, Puigdomenech E, Roura P, et al. Continued smoking abstinence in diabetic patients in primary care: a cluster randomized controlled multicenter study. <i>Diabetes Research & Clinical Practice</i> . 2015;107(1):94-103.	Wrong or missing intervention
Pfizer. A Seven-Week Dose-Ranging Study of CP-526,555 Compared With Placebo and Zyban for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00150241 ; 2000.	Ongoing study
Pfizer. A Study to Evaluate the Safety and Efficacy of Three Doses of a Controlled Release Formulation of Varenicline for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00741884 ; 2010.	Study withdrawn
Philip D, National Heart L, Blood I, Ohio State U. Smoking Cessation Using Motivational Therapy and Varenicline. https://ClinicalTrials.gov/show/NCT00701896 ; 2008.	Ongoing study
Piper ME, Fiore MC, Smith SS, Fraser D, Bolt DM, Collins LM, et al. Identifying effective intervention components for smoking cessation: A factorial screening experiment. <i>Addiction</i> . 2015;No Pagination Specified.	Wrong study design
Piper ME, Fiore MC, Smith SS, Fraser D, Bolt DM, Collins LM, et al. Identifying effective intervention components for smoking cessation: a factorial screening experiment. <i>Addiction</i> . 2016;111(1):129-41.	Wrong study design
Plebani JG, Lynch KG, Yu Q, Pettinati HM, O'Brien CP, Kampman KM. Results of an initial clinical trial of varenicline for the treatment of cocaine dependence. <i>Drug and Alcohol Dependence</i> . 2012;121(1-2):163-6.	Wrong population
Poh PY, Chyi Ng CC. A retrospective review of a pharmacist-managed smoking cessation program in a Singapore Hospital. <i>Pharmacotherapy</i> . 2015;35 (11):e242-e3.	Cannot locate full text
Pollak KI, Oncken CA, Lipkus IM, Lyna P, Swamy GK, Pletsch PK, et al. Nicotine replacement and Behavioral therapy for smoking cessation in pregnancy. <i>American Journal of Preventive Medicine</i> . 2007;33(4):297-305.	Wrong population
Polonio I, Saruhashi T, Endo C, Kiso K, Migotto M, Sgavioli M, et al. Profile of patients attending the outpatient smoking cessation clinic of Santa Casa De Sao Paulo. Comparison of two populations: Military police and general population. <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2014;189(no pagination).	Cannot locate full text
Polosa R, Caponnetto P, Cibella F, Le-Houezec J. Quit and smoking reduction rates in vape shop consumers: a prospective 12-month survey. <i>Int J Environ Res Public Health</i> . 2015;12(4):3428-38.	Wrong study design
Polosa R, Caponnetto P, Maglia M, Morjaria JB, Russo C. Success rates with nicotine personal vaporizers: a prospective 6-month pilot study of smokers not intending to quit. <i>Bmc Public Health</i> . 2014;14:1159.	Wrong study design
Polosa R, Caponnetto P, Morjaria JB, Papale G, Campagna D, Russo C. Effect of an electronic nicotine delivery device (e-Cigarette) on smoking reduction and cessation: a prospective 6-month pilot study. <i>Bmc Public Health</i> . 2011;11:786.	Wrong study design
Polosa R, Morjaria J, Caponnetto P, Caruso M, Strano S, Battaglia E, et al. Effect of Smoking Abstinence and Reduction in Asthmatic Smokers Switching to Electronic Cigarettes: Evidence for Harm Reversal. <i>Int J Environ Res Public Health</i> . 2014;11(5):4965-77.	Wrong study design

Pomerleau OF, Pomerleau CS, Marks JL, Snedecor SM, Mehringer AM, Brouwer RJN, et al. Prolonged nicotine patch use in quitters with past abstinence-induced depressed mood. <i>Journal of Substance Abuse Treatment</i> . 2003;24(1):13-8.	Participants received same drug
Pourmand G, Alidaee MR, Rasuli S, Maleki A, Mehraei A. Do cigarette smokers with erectile dysfunction benefit from stopping?: a prospective study. <i>BJU International</i> . 2004;94(9):1310-3.	Participants received same drug
Prapavessis H, De Jesus S, Fitzgeorge L, Faulkner G, Maddison R, Batten S. Exercise to Enhance Smoking Cessation: the Getting Physical on Cigarette Randomized Control Trial. <i>Annals of Behavioral Medicine</i> . 2016;50(3):358-69.	Participants received same drug
Pravettoni G, Masiero M, Lucchiari C, Maisenneuve P, Mazzocco K, Veronesi G. The role of electronic cigarettes in smoking cessation among heavy smokers undergoing a lung cancer screening program: Preliminary results of a randomized controlled study. <i>Psycho-Oncology</i> . 2016;25:72.	Ongoing study
Primack BA. Varenicline, combination NRT, and nicotine patch did not differ for smoking cessation at 6 mo: Commentary. <i>Annals of Internal Medicine</i> . 2016;164(10):JC54.	Wrong study design
Prochaska JJ, Epperson A, Skan J, Oppezco M, Barnett P, Delucchi K, et al. The Healing and Empowering Alaskan Lives Toward Healthy-Hearts (HEALTHH) Project: Study protocol for a randomized controlled trial of an intervention for tobacco use and other cardiovascular risk behaviors for Alaska Native People. <i>Contemporary Clinical Trials</i> . 2018;71:40-6.	Ongoing study
Prochaska JJ, Fromont SC, Ramo DE, Young-Wolff KC, Delucchi K, Brown RA, et al. Gender differences in a randomized controlled trial treating tobacco use among adolescents and young adults with mental health concerns. <i>Nicotine and Tobacco Research</i> . 2014;17(4):479-85.	Wrong population
Prochaska JJ, Hall SE, Delucchi K, Hall SM. Efficacy of initiating tobacco dependence treatment in inpatient psychiatry: a randomized controlled trial. <i>American journal of public health</i> . 2014;104(8):1557-65.	Wrong study design
Prochaska JJ, Hall SE, Hall SM. Stage-Tailored Tobacco Cessation Treatment in Inpatient Psychiatry. <i>Psychiatric Services</i> . 2009;60(6):848-.	Linked to excluded study
Prochazka AV, Kick S, Steinbrunn C, Miyoshi T, Fryer GE. A randomized trial of nortriptyline combined with transdermal nicotine for smoking cessation. <i>Archives of Internal Medicine</i> . 2004;164(20):2229-33.	Participants received same drug
Prokhorov AV, Marani S, Vost T, Luca M, Mullen M. Project SUCCESS : students using computerized coaching to end smoking successfully. Society for research on nicotine & tobacco 17th annual meeting, february 16-19, toronto, 2011. 2011:18.	Linked to excluded study
Przulj D, Wehbe L, McRobbie H, Hajek P. Progressive nicotine patch dosing prior to quitting smoking: feasibility, safety and effects during the pre-quit and post-quit periods. <i>Addiction</i> . 2018;29:29.	Wrong study design
Puente D, Cabezas C, Rodriguez-Blanco T, Fernandez-Alonso C, Cebrian T, Torrecilla M, et al. The role of gender in a smoking cessation intervention: a cluster randomized clinical trial. <i>BMC Public Health</i> . 2011;11.	Wrong study design
Puerto A, Marqueta A, Nerin I, Jimenez A, Lanuza J, Saenz MA. Profile of prescription, medication adherence and efficacy of the smoking cessation treatment. <i>European Journal of Clinical Pharmacology</i> . 2010;66:S88.	Cannot locate full text
Puri S, Mirza S, DeRycke EC, Bastian L. Inpatient smoking cessation counseling and pharmacotherapy: A practice gap. Chest Conference: CHEST. 2015;148(4 MEETING ABSTRACT).	Cannot locate full text

Queen Mary University of L, Pfizer. De-nicotinised Cigarettes Study. https://ClinicalTrials.gov/show/NCT01250301 ; 2011.	Wrong or missing intervention
Queen Mary University of L. Combination of Nicotine Replacement Therapy (NRT) and Varenicline to Increase Cessation of Tobacco (CONVICT Study). https://ClinicalTrials.gov/show/NCT01184664 ; 2011.	Ongoing study
Raja M, Saha S, Krishna-Reddy V, Mohd S, Narang R, Sood P. Effectiveness of oral health education versus nicotine replacement therapy for tobacco cessation- a parallel randomized clinical trial. Journal of Clinical & Experimental Dentistry. 2016;8(1):e64-70.	Wrong comparator
Raja M, Saha S, Mohd S, Narang R, Reddy LVK, Kumari M. Cognitive Behavioural Therapy versus Basic Health Education for Tobacco Cessation among Tobacco Users: A Randomized Clinical Trail. Journal of clinical and diagnostic research : JCDR. 2014;8(4):ZC47-9.	Wrong or missing intervention
Ramo DE, Thrul J, Chavez K, Delucchi KL, Prochaska JJ. Feasibility and Quit Rates of the Tobacco Status Project: A Facebook Smoking Cessation Intervention for Young Adults. Journal of medical Internet research. 2015;17(12):e291.	Wrong or missing intervention
Ray LA, MacKillop J, Hutchison KE. Pharmacogenetics of the micro -opioid receptor and the treatment of addictions. Personalized Medicine. 2007;4(2):217-20.	Wrong study design
Regan AK, Promoff G, Dube SR, Arrazola R. Electronic nicotine delivery systems: adult use and awareness of the 'e-cigarette' in the USA. Tobacco Control. 2013;22(1):19-23.	Wrong study design
Reid RD, Armstrong A, McDonnell L, Aitken DA, Robert L, LaRue A, et al. VARENICLINE VERSUS TRANSDERMAL NICOTINE PATCH FOR SMOKING CESSATION IN PATIENTS WITH CORONARY HEART DISEASE: A PILOT RANDOMIZED TRIAL. Canadian Journal of Cardiology. 2010;26:53D-D.	Ongoing study
Reid RD, Pipe A, Dafoe WA. Is telephone counselling a useful addition to physician advice and nicotine replacement therapy in helping patients to stop smoking? A randomized controlled trial. Canadian Medical Association Journal. 1999;160(11):1577-81.	Participants received same drug
Reid ZZ, Regan S, Kelley JHK, Streck JM, Ylioja T, Tindle HA, et al. Comparative effectiveness of post-discharge strategies for hospitalized smokers: study protocol for the Helping HAND 2 randomized controlled trial. BMC Public Health. 2015;15.	Wrong study design
Research VAOo, Development. CPT and Smoking Cessation. https://ClinicalTrials.gov/show/NCT01901848 ; 2013.	Participants received same drug
Research VAOo, Development. Family-supported Smoking Cessation for Chronically Ill Veterans. https://ClinicalTrials.gov/show/NCT00448344 ; 2008.	Participants received same drug
Research VAOo, Development. Mood and Smoking: A Comparison of Smoking Cessation Treatments. https://ClinicalTrials.gov/show/NCT00075972 ; 2003.	Wrong study design
Research VAOo, Development. Nicotine Delivery Systems: Research & Treatment. https://ClinicalTrials.gov/show/NCT00108342 ; 2007.	Study terminated
Research VAOo, Development. Smoking Cessation for Veterans With Severe and Persistent Mental Illness. https://ClinicalTrials.gov/show/NCT00960375 ; 2010.	Wrong or missing intervention

Rettig EM, Fakhry C, Hales RK, Kisuule F, Quon H, Kiess AP, et al. Pilot randomized controlled trial of a comprehensive smoking cessation intervention for patients with upper aerodigestive cancer undergoing radiotherapy. <i>Head and Neck-Journal for the Sciences and Specialties of the Head and Neck</i> . 2018;40(7):1534-47.	Wrong study design
Richmond R, Indig D, Butler T, Wilhelm K, Archer V, Wodak A. A randomized controlled trial of a smoking cessation intervention conducted among prisoners. <i>Addiction (Abingdon, England)</i> . 2013;108(5):966-74.	Wrong comparator
Richmond R, Indig D, Butler T, Wilhelm K, Archer V, Wodak A. A randomized controlled trial of a smoking cessation intervention conducted among prisoners. <i>Addiction</i> . 2013;108(5):966-74.	Participants received same drug
Rigotti NA, Bitton A, Kelley JK, Hoepfner BB, Levy DE, Mort E. Supplementing office-based care with a populationbased direct-to-smoker outreach intervention offering free treatment to smokers in a community health center: A randomized controlled trial. <i>Journal of General Internal Medicine</i> . 2011;26:S33-S4.	Wrong study design
Rigotti NA, Chang Y, Tindle HA, Kalkhoran SM, Levy DE, Regan S, et al. Association of E-Cigarette Use With Smoking Cessation Among Smokers Who Plan to Quit After a Hospitalization: A Prospective Study. <i>Annals of Internal Medicine</i> . 2018;168(9):613-20.	Linked to excluded study
Rigotti NA, Gonzales D, Dale LC, Lawrence D, Chang Y, Cirrus Study G. A randomized controlled trial of adding the nicotine patch to rimonabant for smoking cessation: efficacy, safety and weight gain. <i>Addiction</i> . 2009;104(2):266-76.	Wrong study design
Rigotti NA, Japuntich S, Regan S, Kelley JH, Chang Y, Reyen M, et al. Promoting smoking cessation after hospital discharge: The helping hand randomized controlled comparative effectiveness trial. <i>Journal of General Internal Medicine</i> . 2013;28:S160.	Linked to excluded study
Rigotti NA, Regan S, Levy DE, Japuntich S, Chang Y, Park ER, et al. Sustained care intervention and postdischarge smoking cessation among hospitalized adults a randomized clinical trial. <i>JAMA - Journal of the American Medical Association</i> . 2014;312(7):719-28.	Wrong study design
Robinson JD, Li L, Chen M, Lerman C, Tyndale RF, Schnoll RA, et al. Evaluating the temporal relationships between withdrawal symptoms and smoking relapse. <i>Psychology of Addictive Behaviors</i> . 2019;07:07.	Awaiting classification
Roddy E, Romilly N, Challenger A, Lewis S, Britton J. Use of nicotine replacement therapy in socioeconomically deprived young smokers: a community-based pilot randomised controlled trial. <i>Tobacco Control</i> . 2006;15(5):373-6.	Wrong population
Rodu B, Stegmayr B, Nasic S, Cole P, Asplund K. The influence of smoking and smokeless tobacco use on weight amongst men. <i>Journal of Internal Medicine</i> . 2004;255(1):102-7.	Wrong study design
Roelsgaard I, Thomsen T, Ostergaard M, Christensen R, Hetland M, Jacobsen S, et al. The effect of an intensive smoking cessation intervention on disease activity in patients with rheumatoid arthritis: study protocol for a randomised controlled trial2017; 18(1):[570 p.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/175/CN-01446175/frame.html .	Ongoing study
Rogers E, Fu S, Krebs P, Noorbaloochi S, Nugent S, Gravely A, et al. Proactive tobacco treatment for smokers using VA mental health clinics: A randomized controlled trial. <i>Journal of General Internal Medicine</i> . 2018;33 (2 Supplement 1):313.	Linked to excluded study
Rogers ES, Fu SS, Krebs P, Noorbaloochi S, Nugent SM, Gravely A, et al. Proactive Tobacco Treatment for Smokers Using Veterans Administration Mental Health Clinics. <i>American Journal of Preventive Medicine</i> . 2018;54(5):620-9.	Participants received same drug
Rogers ES, Fu SS, Krebs P, Noorbaloochi S, Nugent SM, Rao R, et al. Proactive outreach for smokers using VHA mental health clinics: protocol for a patient-randomized clinical trial. <i>BMC public health</i> . 2014;14:1294.	Participants received same drug

Rohsenow DJ, Monti PM, Colby SM, Martin RA. Brief interventions for smoking cessation in alcoholic smokers. <i>Alcoholism-Clinical and Experimental Research</i> . 2002;26(12):1950-1.	Linked to excluded study
Roig L, Perez S, Prieto G, Martin C, Advani M, Armengol A, et al. Cluster randomized trial in smoking cessation with intensive advice in diabetic patients in primary care. ITADI Study. <i>BMC Public Health</i> . 2010;10.	Linked to excluded study
Rolnick SJ, Klevan D, Cherney L, Lando HA. Nicotine replacement therapy in a group model HMO. <i>HMO practice</i> . 1997;11(1):34-7.	Linked to excluded study
Rose JE, Behm FM, Westman EC, Kukovich P. Precessation treatment with nicotine skin patch facilitates smoking cessation. <i>Nicotine & Tobacco Research</i> . 2006;8(1):89-101.	Wrong study design
Rose JE, Behm FM, Westman EC, Levin ED, Stein RM, Ripka GV. MECAMYLAMINE COMBINED WITH NICOTINE SKIN PATCH FACILITATES SMOKING CESSATION BEYOND NICOTINE PATCH TREATMENT ALONE. <i>Clinical Pharmacology & Therapeutics</i> . 1994;56(1):86-99.	Wrong study design
Rose Research Center LLC, National Institutes of H, Food, Drug A. Abuse Liability of Reduced Nicotine Content Cigarettes in the Context of Concurrent E-Cigarette Use. https://ClinicalTrials.gov/show/NCT02870218 ; 2017.	Ongoing study
Roswell Park Cancer I, National Cancer I. Varenicline or Nicotine Patch in Promoting Smoking Cessation Among Current Smokers. https://ClinicalTrials.gov/show/NCT01771627 ; 2012.	Ongoing study
Roswell Park Cancer I, Pfizer. An Alternative Dosing Schedule of Varenicline for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00835900 ; 2009.	Ongoing study
Roswell Park Cancer I. Bupropion in Helping Adults Stop Smoking. https://ClinicalTrials.gov/show/NCT00534001 ; 2006.	Ongoing study
Roth MT, Westman EC. Use of bupropion SR in a pharmacist-managed outpatient smoking-cessation program. <i>Pharmacotherapy:The Journal of Human Pharmacology & Drug Therapy</i> . 2001;21(5):636-41.	Wrong comparator
Rowe AS, McMillen JC, Wylie DR. Outcomes associated with nicotine replacement therapy in patients with subarachnoid hemorrhage. <i>Neurocritical Care</i> . 2011;1):S109.	Wrong comparator
Rozario HP. Nicotine dependence assessment using Fagerstrom test and Nicotine Replacement therapy (NRT) recommendation techniques for smoking cessation among Paniya tribes. <i>Annals of Oncology</i> . 2016;27 (Supplement 9):ix187-ix8.	Wrong study design
Rutgers TSUoNJ, National Institute on Drug A. Developing a Smoking Cessation Intervention for Methadone Maintained Smokers. https://ClinicalTrials.gov/show/NCT01393392 ; 2011.	Ongoing study
Ruther T, Ruderer A, Wirth C, Schuler V, Lang V, Linhardt A, et al. Smoking Cessation Program for Inpatients with Substance Use Disorder: A Quasi-Randomized Controlled Trial of Feasibility and Efficacy. <i>European Addiction Research</i> . 2016;22(5):268-76.	Wrong study design
Saiz Martinez PA, Fernandez-Artamendi S, Al-Halabi S, Diaz-Mesa E, Garcia-Alvarez L, Martinez E, et al. Tobacco use and clinical symptoms in a sample of outpatients diagnosed with schizophrenia. <i>European Psychiatry</i> . 2015;30:266.	Cannot locate full text
Sallsten G, Thoren J, Barregard L, Schutz A, Skarping G. Long-term use of nicotine chewing gum and mercury exposure from dental amalgam fillings. <i>Journal of Dental Research</i> . 1996;75(1):594-8.	Wrong population
Sampablo I, Lores L, Coll-Klein F, Jimenez C, Rebasa P. Predictive factors in smoking cessation with combined therapy with bupropion and nicotine patches. <i>Monaldi Archives for Chest Disease</i> . 2003;59(2):171-6.	Participants received same drug
Sampablo Lauro I, Lores L, Coll F, Rebasa P. Combining nicotine patches and bupropion in smoking cessation therapy. [Spanish]. <i>Archivos de Bronconeumologia</i> . 2000;36(7):377-80.	Wrong study design

Sarkar BK, Shahab L, Arora M, Lorencatto F, Reddy KS, West R. A cluster randomized controlled trial of a brief tobacco cessation intervention for low-income communities in India: study protocol. <i>Addiction</i> . 2014;109(3):371-8.	Wrong or missing intervention
Saul JE, Lien R, Schillo B, Kavanaugh A, Wendling A, Luxenberg M, et al. Outcomes and cost-effectiveness of two nicotine replacement treatment delivery models for a tobacco quitline. <i>International Journal of Environmental Research & Public Health [Electronic Resource]</i> . 2011;8(5):1547-59.	Wrong study design
Saules KK, Schuh LM, Arfken CL, Reed K, Kilbey MM, Schuster CR. Double-blind placebo-controlled trial of fluoxetine in smoking cessation treatment including nicotine patch and cognitive-behavioral group therapy. <i>American Journal on Addictions</i> . 2004;13(5):438-46.	Participants received same drug
Schlagintweit HE, Good P, Barrett SP. The impact of nicotine lozenges and stimulus expectancies on cigarette craving. <i>Behavioural Pharmacology</i> . 2013;24:e50.	Cannot locate full text
Schlam TR, Fiore MC, Smith SS, Fraser D, Bolt DM, Collins LM, et al. Comparative effectiveness of intervention components for producing long-term abstinence from smoking: A factorial screening experiment. <i>Addiction</i> . 2015:No Pagination Specified.	Wrong study design
Schmidt F. MEDICAL SUPPORT OF NICOTINE WITHDRAWAL. <i>Munchener Medizinische Wochenschrift</i> . 1974;116(11):557-64.	Wrong or missing intervention
Schmidt N, Raines A, Allan N, Zvolensky M. Anxiety sensitivity risk reduction in smokers: A randomized control trial examining effects on panic2016; 77(R01 MH076629-0 (NIMH) *National Institute of Mental Health*):[138-46 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/060/CN-01128060/frame.html .	Wrong comparator
Schneider D, Meienberg A, Deman A, Burkard T. Setting up a smoking cessation service-what can we expect from the first year? <i>Praxis</i> . 2015;104:123.	Wrong comparator
Schnoll RA, Cappella J, Lerman C, Pinto A, Patterson F, Wileyto EP, et al. A Novel Recruitment Message to Increase Enrollment Into a Smoking Cessation Treatment Program: Preliminary Results From a Randomized Trial. <i>Health Communication</i> . 2011;26(8):735-42.	Participants received same drug
Schoenberg NE, Studts CR, Shelton BJ, Liu M, Clayton R, Bispo JB, et al. A randomized controlled trial of a faith-placed, lay health advisor delivered smoking cessation intervention for rural residents. <i>Preventive Medicine Reports</i> . 2016;3:317-23.	Wrong study design
Schuck K, Bricker JB, Otten R, Kleinjan M, Brandon TH, Engels RC. Effectiveness of proactive quitline counselling for smoking parents recruited through primary schools: results of a randomized controlled trial. <i>Addiction</i> . 2014;109(5):830-41.	Wrong or missing intervention
Secades-Villa R, Alonso-Perez F, Garcia-Rodriguez O, Fernandez-Hermida JR. EFFECTIVENESS OF THREE INTENSITIES OF SMOKING CESSATION TREATMENT IN PRIMARY CARE. <i>Psychological Reports</i> . 2009;105(3):747-58.	Wrong or missing intervention
Selby P, Brosky G, Oh P, Raymond V, Arteaga C, Ranger S. A pragmatic, randomized, controlled study evaluating the impact of access to smoking cessation pharmacotherapy coverage on the proportion of successful quitters in a Canadian population of smokers motivated to quit (ACCESSATION). <i>BMC public health</i> . 2014;14:433.	Participants received same drug
Semmelweis U. <i>Epidemiology and Acute Oral Effects Electronic Cigarette</i> . https://ClinicalTrials.gov/show/NCT03011710 ; 2015.	Ongoing study
Seoul National University H, Pfizer. Intervention Trial of Culturally-Appropriate Decision Aids for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01566097 ; 2012.	Wrong study design

Severson HH, Akers L, Andrews JA, Lichtenstein E, Jerome A. Evaluating two self-help interventions for smokeless tobacco cessation. Addictive Behaviors. 2000;25(3):465-70.	Wrong or missing intervention
Severson HH, Andrews JA, Lichtenstein E, et al. Smokeless tobacco cessation through dental offices: An intervention that works. Journal of Dental Research. 1998;77:1206.	Linked to excluded study
Severson HH, Andrews JA, Lichtenstein E, Gordon JS, Barckley M, Akers L. A self-help cessation program for smokeless tobacco users: comparison of two interventions. Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco. 2000;2(4):363-70.	Wrong or missing intervention
Severson HH, Andrews JA, Lichtenstein E, Gordon JS, Barckley MF. Using the hygiene visit to deliver a tobacco cessation program: Results of a randomized clinical trial. Journal of the American Dental Association. 1998;129(7):993-9.	Wrong or missing intervention
Severson HH, Gordon JS, Danaher BG, Akers L. ChewFree.com: Evaluation of a Web-based cessation program for smokeless tobacco users. Nicotine & Tobacco Research. 2008;10(2):381-91.	Wrong or missing intervention
Severson HH, Peterson AL, Andrews JA, Gordon JS, Cigrang JA, Danaher BG, et al. Smokeless tobacco cessation in military personnel: A randomized controlled trial. Nicotine & Tobacco Research. 2009;11(6):730-8.	Wrong or missing intervention
Shaare Zedek Medical C. Extended (6-Week) Varenicline Preloading: Does it Facilitate Smoking Reduction and Cessation? : https://ClinicalTrials.gov/show/NCT02634281 ; 2016.	Ongoing study
Shaare Zedek Medical C. Varenicline for "Gradual" vs. "Abrupt" Smoking Cessation in Low-motivated COPD Smokers. https://ClinicalTrials.gov/show/NCT02894957 ; 2016.	Ongoing study
Shah PA, Cunningham CO, Brisbane MT, DeLuca JP, Nahvi S. Use of Smoking Cessation Methods Among Patients Receiving Office-based Buprenorphine Maintenance Treatment. Journal of Addiction Medicine. 2017;11(6):494-7.	Wrong study design
Shahab L, Mortimer E, Bauld L, McGowan JA, McNeill A, Tyndale RF. Characterising the nicotine metabolite ratio and its association with treatment choice: A cross sectional analysis of Stop Smoking Services in England. Scientific Reports. 2017;7.	Wrong study design
Shand J, Howe A, Cinnamon N, Hussey S, Menown IB. Real world 12 month evaluation of varenicline therapy. Atherosclerosis Supplements. 2011;12 (1):170-1.	Cannot locate full text
Shaughnessy AF, Davis RE, Reeder CE. Nicotine chewing gum: Effectiveness and the influence of patient education in a family practice. Journal of Family Practice. 1987;25(3):266-9.	Awaiting classification
Sheba Medical C, Hadassah Medical O, Barzilai Medical C. Early In-hospital Initiation of Pharmacotherapy for Smoking Cessation, Patients After ACS. https://ClinicalTrials.gov/show/NCT02106637 ; 2014.	Ongoing study
Sheffield Hallam U, King's College L, Heart Research UK, Help2Change. E-cigarettes and Cardiovascular Function. https://ClinicalTrials.gov/show/NCT03061253 ; 2017.	Ongoing study
Shelley D, Cantrell J, Warn D, Wong S. Smoking cessation among sheltered homeless: a pilot. American Journal of Health Behavior. 2010;34(5):544-52.	Wrong study design
Shelley D, Tseng TY, Gonzalez M, Krebs P, Wong S, Furberg R, et al. Correlates of Adherence to Varenicline Among HIV+ Smokers. Nicotine & Tobacco Research. 2015;17(8):968-74.	Ongoing study

Shelley D, Zraik D, Tseng T, Jannat-Khah D, Pae-Cho C, Moon L. Effect of short acting vs long acting nicotine replacement therapy among a low SES population of light smokers (POS3-77)2012:[113 p.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/596/CN-00814596/frame.html .	Awaiting classification
Sherman S, Link AR, Rogers E, Krebs P, Ladapo JA, Shelley D, et al. Smoking cessation interventions for urban hospital patients: A randomized comparative effectiveness trial. <i>Journal of General Internal Medicine</i> . 2015;30:S253.	Participants received same drug
Shewale AR, Borse MS, Brown JD, Li C. Mental health status of varenicline and bupropion users during a quit attempt compared to current smokers, other quitters, and non-smokers. <i>Drug and Alcohol Dependence</i> . 2015;154:132-8.	Wrong study design
Shiffman S, Paty JA, Rohay JM, Di Marino ME, Gitchell J. The efficacy of computer-tailored smoking cessation material as a supplement to nicotine polacrilex gum therapy. <i>Archives of Internal Medicine</i> . 2000;160(11):1675-81.	Participants received same drug
Shimony A, Grandi SM, Pilote L, Joseph L, O'Loughlin J, Paradis G, et al. Utilization of evidence-based therapy for acute coronary syndrome in high income and middle/low income countries. <i>Canadian Journal of Cardiology</i> . 2013;1):S297-S8.	Cannot locate full text
Short NA, Mathes BM, Gibby B, Oglesby ME, Zvolensky MJ, Schmidt NB. Insomnia symptoms as a risk factor for cessation failure following smoking cessation treatment. <i>Addiction Research & Theory</i> . 2017;25(1):17-23.	Linked to excluded study
Siahpush M, Shaikh RA, McCarthy M, Sikora Kessler A, Tibbits M, Singh GK. Association between duration of use of pharmacotherapy and smoking cessation: Findings from a national survey. <i>BMJ Open</i> . 2015;5 (1) (no pagination)(e006229).	Wrong study design
SianJennings CSC, Kotseva K, De Backer G, De Bacquer D, Gaita D, Keil U, et al. How well supported are high CVD risk smokers to quit their habit? EUROASPIRE III survey of individuals at high risk of developing CVD from 12 European regions. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> . 2010;17:S93.	Wrong study design
Signes-Costa J, Sancho JN, Chiner E, Senent C, Gomez-Merino E, Mediero G, et al. Efficacy of varenicline in smokers with chronic obstructive pulmonary disease. <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2011;183(1 MeetingAbstracts).	Participants received same drug
Sigurgeirsdottir J, Petursdottir G, Gudmundsdottir T, Jonsdottir A, Ingimarsdottir IE. The results of a nurse-led nicotine dependence therapy as a part of interdisciplinary rehabilitation in Iceland. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2014;44(no pagination).	Wrong study design
Silver B, Ramey SJ, Diaz DA, Abreu Z, Vega M, Yechieli R, et al. Smoking cessation at a safety-net hospital: Results from a resident-led prospective quality improvement initiative. <i>International Journal of Radiation Oncology Biology Physics</i> . 2018;101 (2 Supplement 1):E41.	Ongoing study
Simbec R, creative CN. A Study of Blood Levels of Nicotine Following an Electronic Cigarette. https://ClinicalTrials.gov/show/NCT02794220 ; 2012.	Ongoing study
Simon JA, Duncan C, Carmody TP, Hudes ES. Bupropion plus nicotine replacement no better than replacement alone. <i>Journal of Family Practice</i> . 2004;53(12):953-4.	Awaiting classification
Sippel JM, Osborne ML, Bjornson W, Goldberg B, Buist AS. Smoking cessation in primary care clinics. <i>Journal of General Internal Medicine</i> . 1999;14(11):670-6.	Wrong or missing intervention
Smit ES, Evers S, de Vries H, Hoving C. Cost-Effectiveness and Cost-Utility of Internet-Based Computer Tailoring for Smoking Cessation. <i>Journal of Medical Internet Research</i> . 2013;15(3):55-70.	Wrong or missing intervention
Smith M, Fu S, Nugent S, Nelson D, Joseph A, An L. A COST-EFFECTIVE TELEHEALTH INTERVENTION FOR SMOKING CESSATION. <i>Journal of General Internal Medicine</i> . 2010;25:208-.	Wrong study design

Smith SS, Jorenby DE, Fiore MC, Anderson JE, Mielke MM, Beach KE, et al. Strike while the iron is hot: Can stepped-care treatments resurrect relapsing smokers? Journal of Consulting and Clinical Psychology. 2001;69(3):429-39.	Participants received same drug
Smith SS, Keller PA, Kobinsky KH, Baker TB, Fraser DL, Bush T, et al. Enhancing tobacco quitline effectiveness: Identifying a superior pharmacotherapy adjuvant. Nicotine and Tobacco Research. 2013;15(3):718-28.	Wrong study design
Smoke-Break I. Clinical Study of Smoke-Break Liquid Cigarettes. https://ClinicalTrials.gov/show/NCT00715871 ; 2008.	Wrong or missing intervention
Solano Rein S, Vaquero Lozano P, Solano Garcia-Tenorio R, Lopez Ruiz T, Jimenez Ruiz CA, de Granda Orive JI. Treatment of smoking habit in chronic obstructive pulmonary disease. [Spanish]. Revista de Patologia Respiratoria. 2012;15(4):123-8.	Awaiting classification
Soroka University Medical C, Novartis. In-patient Smoking Cessation Intervention Using Counseling, Spirometry and Nicotine Replacement Therapy. https://ClinicalTrials.gov/show/NCT02470923 ; 2015.	Ongoing study
Sousa AB, Lotufo JPB, Fukuoka MH, Sakai MC, Galvao GF, Ribeiro E, et al. Smoking cessation pharmacotherapy improves patients adherence and increases the ability to achieve abstinence from cigarette smoking in a Brazilian teaching hospital. European Journal of Hospital Pharmacy: Science and Practice. 2012;19 (2):220-1.	Awaiting classification
Southwest Oncology G, National Cancer I, Cancer, Leukemia Group B, Eastern Cooperative Oncology G. S0002 - A Program to Quit Smoking With or Without Bupropion in Treating Patients With Stage I or II Non-Small Cell Lung Cancer Who Have Undergone Surgery. https://ClinicalTrials.gov/show/NCT00032084 ; 2002.	Study terminated
Spangler EL, Brooke BS, Beck AW, Hoel A, Farber A, Goodney PP, et al. Design and initial enrollment in the Vascular Physicians Offer and Report (VAPOR) trial. Journal of Vascular Surgery. 2016;63(4):1121-5.e2.	Wrong study design
SS AL, BinShabaib M, Akram Z, Rahman I, Romanos GE, Javed F. Impact of cigarette smoking and vaping on the outcome of full-mouth ultrasonic scaling among patients with gingival inflammation: a prospective study. Clinical Oral Investigations. 2018;26:26.	Wrong study design
Stanford U, International SRI, Johns Hopkins U, University of B, National Institute on Drug A. Behaviorally Enhanced Counseling on Nicotine Dependence (BEACON) Trial. https://ClinicalTrials.gov/show/NCT00991081 ; 2009.	Linked to excluded study
Stanford U, National Institute of Mental H. Evaluation of Tobacco Treatment Strategies for Inpatient Psychiatry. https://ClinicalTrials.gov/show/NCT00968513 ; 2009.	Ongoing study
Stanford U, National Institute on Drug A, University of California SF. Treating Tobacco Dependence in Inpatient Psychiatry - 1. https://ClinicalTrials.gov/show/NCT00136812 ; 2006.	Participants received same drug
Stanford U, National Institute on Drug A. Behavioral Maintenance Treatment for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01615770 ; 2004.	Participants received same drug
Stanford U, National Institute on Drug A. Extended Treatment for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01330043 ; 2010.	Ongoing study
Stanford U, San Francisco Employment Development D, Marin Employment C, JobTrain. Tobacco Treatment for Employable Californians, Total IMPACT Study. https://ClinicalTrials.gov/show/NCT02478697 ; 2015.	Ongoing study
Stanford U, University of California SF, Alaska Native Tribal Health C, University of Alaska A, Norton Sound Health C. Healing and Empowering Alaskan Lives Towards Healthy-Hearts Study. https://ClinicalTrials.gov/show/NCT02137902 ; 2015.	Ongoing study

Stevens VJ, Severson H, Lichtenstein E, Little SJ, Leben J. MAKING THE MOST OF A TEACHABLE MOMENT - A SMOKELESS-TOBACCO CESSATION INTERVENTION IN THE DENTAL OFFICE. American Journal of Public Health. 1995;85(2):231-5.	Wrong or missing intervention
Stoleru M, Mihaicuta S. Smoking cessation: A problem for older adults? European Respiratory Journal Conference: European Respiratory Society Annual Congress. 2012;40(no pagination).	Cannot locate full text
Streck JM, Regan S, Chang Y, Kelley JHK, Singer DE, Rigotti NA. Examining the effects of illicit drug use on tobacco cessation outcomes in the Helping HAND 2 randomized controlled trial. Drug and Alcohol Dependence. 2017;178:586-92.	Linked to excluded study
Tadrous M, Martins D, Yao Z, Mamdani MM, Juurlink DN, Gomes T, et al. Varenicline and Risk of Self-Harm: A Nested Case-Control Study. 2016;11(9):e0163681.	Participants received same drug
Taggar JS, Lewis S, Docherty G, Bauld L, McEwen A, Coleman T. Do cravings predict smoking cessation in smokers calling a national quit line: secondary analyses from a randomised trial for the utility of 'urges to smoke' measures. Substance Abuse Treatment Prevention and Policy. 2015;10.	Linked to excluded study
Tan AS, Bigman CA, Sanders-Jackson A. Sociodemographic correlates of self-reported exposure to ecigarette communications and its association with public support for smoke-free and vape-free policies: results from a national survey of US adults. Tobacco Control. 2015;24:574-81.	Wrong study design
Tan ASL, Bigman CA. E-Cigarette Awareness and Perceived Harmfulness Prevalence and Associations with Smoking-Cessation Outcomes. American Journal of Preventive Medicine. 2014;47(2):141-9.	Wrong study design
Ternullo SR, Abdolahi A, Williams GC. Study of monotherapy versus combination therapy for tobacco dependence among heavily addicted smokers. Journal of the American Pharmacists Association. 2017;57(1):77-+.	Awaiting classification
The Mind Research N, National Institute on Drug A. TDCS and Cognitive Retraining to Augment Pharmacotherapy for the Treatment of Nicotine Dependence. https://ClinicalTrials.gov/show/NCT02534454 ; 2015.	Wrong or missing intervention
The Queen Elizabeth H, Basil Hetzel Institute for Translational R, University of A, University of South A, Clinical Practice U, Therapeutics Research C. Evaluation of Transdermal Nicotine Replacement Therapy (NRT) Activity Through Metabolic Induction. https://ClinicalTrials.gov/show/NCT01438944 ; 2011.	Wrong comparator
The University of Hong K, Food, Health Bureau HK, Queen Mary Hospital HK, Tuen Mun H. Smoking Cessation in Cancer Patients. https://ClinicalTrials.gov/show/NCT01685723 ; 2012.	Ongoing study
The University of Hong K. Motivating Smokers at Outdoor Public Smoking Hotspots to Have Quit Attempt With Nicotine Replacement Therapy Sampling. https://ClinicalTrials.gov/show/NCT02491086 ; 2015.	Wrong comparator
The University of Texas Health Science Center H, Michael Deakey Veterans Affairs Medical C, Baylor College of M. Exenatide Once Weekly for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02975297 ; 2016.	Wrong or missing intervention
The University of Texas Health Science Center H. E-cigarettes to Promote Smoking Reduction Among Individuals With Schizophrenia. https://ClinicalTrials.gov/show/NCT02918630 ; 2016.	Ongoing study
Thomas D, Abramso nMJ, Bonevski B, Taylor S, Poole S, Weeks GR, et al. A pharmacist-led system-change smoking cessation intervention for smokers admitted to Australian public hospitals (GIVE UP FOR GOOD): study protocol for a randomised controlled trial. Trials. 2013;14:148.	Participants received same drug

Thomas D, Leckbee G, Pope N, Montemayor D, Ashley J, Cervantes J. Assessment of community pharmacist intervention to improve outcomes in smoking cessation using a behavioral and pharmacotherapy approach. <i>Journal of the American Pharmacists Association</i> . 2012;52 (2):247.	Ongoing study
Tidey JW, Cassidy RN, Miller ME. Smoking Topography Characteristics of Very Low Nicotine Content Cigarettes, With and Without Nicotine Replacement, in Smokers With Schizophrenia and Controls. <i>Nicotine & Tobacco Research</i> . 2016;18(9):1807-12.	Wrong study design
Tidey JW, Rohsenow DJ, Kaplan GB, Swift RM, Ahnallen CG. Separate and combined effects of very low nicotine cigarettes and nicotine replacement in smokers with schizophrenia and controls. <i>Nicotine and Tobacco Research</i> . 2013;15(1):121-9.	Wrong study design
Tidey JW, Rohsenow DJ. Intention to quit moderates the effect of bupropion on smoking urge. <i>Nicotine & Tobacco Research</i> . 2009;11(3):308-12.	Wrong study design
Toll BA, Leary V, Wu R, Salovey P, Meandzija B, O'Malley SS. A preliminary investigation of naltrexone augmentation of bupropion to stop smoking with less weight gain. <i>Addictive Behaviors</i> . 2008;33(1):173-9.	Wrong comparator
Toll BA, White M, Wu R, Meandzija B, Jatlow P, Makuch R, et al. Low-dose naltrexone augmentation of nicotine replacement for smoking cessation with reduced weight gain: a randomized trial. <i>Drug & Alcohol Dependence</i> . 2010;111(3):200-6.	Participants received same drug
Tomasulo MMV, Scanlin MC, Patterson SM. The Effects of Nicotine and Nicotine Abstinence on Stress-Induced Cardiovascular Reactivity Do Habitual and Light/Intermittent Cigarette Smokers Differ? <i>Journal of Psychophysiology</i> . 2017;31(3):116-33.	Awaiting classification
Tomioka H, Sekiya R, Yamashita S, Kaneda T, Kida Y, Nishio C, et al. Health-related quality of life in smokers who attended a smoking cessation clinic-cross-sectional and longitudinal study. <i>American Journal of Respiratory and Critical Care Medicine Conference: American Thoracic Society International Conference, ATS</i> . 2012;185(no pagination).	Cannot locate full text
Tonnesen P, Mikkelsen K. A placebo-controlled trial with varenicline for long term nicotine replacement product users. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2011;38(no pagination).	Wrong population
Tonnesen P, Mikkelsen K. Varenicline to Stop Long-term Nicotine Replacement Use: A Double-Blind, Randomized, Placebo-Controlled Trial. <i>Nicotine & Tobacco Research</i> . 2013;15(2):419-27.	Wrong population
Tonstad S, Gustavsson G, Kruse E, Walmsley JM, Westin A. Symptoms of nicotine toxicity in subjects achieving high cotinine levels during nicotine replacement therapy. <i>Nicotine and Tobacco Research</i> . 2014;16 (9) (no pagination)(ntu076).	Wrong study design
Tonstad S, Heggen E, Giljam H, Lagerback PA, Tonnesen P, Wikingsson LD, et al. Niccine (R), a Nicotine Vaccine, for Relapse Prevention: A Phase II, Randomized, Placebo-Controlled, Multicenter Clinical Trial. <i>Nicotine & Tobacco Research</i> . 2013;15(9):1492-501.	Wrong or missing intervention
Tonstad S, Heggen E, Giljam H, Lagerback PA, Tonnesen P, Wikingsson LD, et al. Niccine, a nicotine vaccine, for relapse prevention: A phase ii, randomized, placebo-controlled, multicenter clinical trial. <i>Nicotine and Tobacco Research</i> . 2013;15(9):1492-501.	Wrong or missing intervention
Tran-Van D, Herve Y, Labadie P, Deroudilhe G, Avargue's P. Restlessness in intensive care unit: think to the nicotinic withdrawal syndrome [French]. <i>Annales françaises d'anesthésie et de réanimation</i> . 2004;23(6):604-6.	Wrong study design
Trofor L, Albu AM, Podgornii A, Barnea E, Trofor AC. Long term evaluation of the effectiveness of the first reimbursed smoking cessation program conducted in the Clinic of pulmonary diseases of Iasi, Romania. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2013;42(no pagination).	Cannot locate full text
Trumbo CW, Harper R. Use and Perception of Electronic Cigarettes Among College Students. <i>Journal of American College Health</i> . 2013;61(3):149-55.	Wrong study design
Umeå U, Karolinska I. Acute Cardiorespiratory Effects of E-cigarette Inhalation. https://ClinicalTrials.gov/show/NCT02899234 ; 2016.	Ongoing study

Universita degli Studi di C. Efficacy And Safety Of Smoking Cessation With Varenicline Tartrate In Diabetic Smokers: (DIASMOKE). https://ClinicalTrials.gov/show/NCT01387425 ; 2011.	Ongoing study
Université Libre de B. Comparison of Electronic Cigarettes and Tobacco Cigarettes on Cardiovascular Function and Oxidative Stress. https://ClinicalTrials.gov/show/NCT03036644 ; 2017.	Ongoing study
University Health Network T, Pfizer. Smoking Cessation Program in the Preadmission Clinic. https://ClinicalTrials.gov/show/NCT01320462 ; 2010.	Ongoing study
University Health Network T, Pfizer. Smoking Cessation Program in the Preadmission Clinic: The Use of a Teachable Moment. https://ClinicalTrials.gov/show/NCT01243203 ; 2007.	Ongoing study
University Hospital B. Efficacy of Varenicline on Smoking Cessation at the Acute Phase of an Exacerbation of Chronic Obstructive Pulmonary Disease. https://ClinicalTrials.gov/show/NCT01694732 ; 2012.	Ongoing study
University of A. The Effects of Electronic Cigarette Smoking on the Arterial Wall and Endothelial Glycocalyx Properties of Smokers. https://ClinicalTrials.gov/show/NCT03039920 ; 2015.	Wrong comparator
University of Alabama at B. In Vivo Experience With NRT to Increase Adherence and Smoking Abstinence. https://ClinicalTrials.gov/show/NCT02938403 ; 2017.	Participants received same drug
University of Auckland NZ, Health New Zealand Ltd CNZ, Auckland District Health B. The Use of Nicotine Patches Together With E-cigarettes (With and Without Nicotine) for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02521662 ; 2016.	Ongoing study
University of Auckland NZ, Lakes District Health B, Brunel U. Cytisine Versus Varenicline for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02957786 ; 2017.	Wrong comparator
University of British C, Providence H, Services, St. Paul's Hospital C. Nicotine Replacement Provided at a Tertiary Care Hospital. https://ClinicalTrials.gov/show/NCT01614054 ; 2012.	Participants received same drug
University of C, Pfizer. Varenicline Augmentation of Patch Outcomes in Heavy Drinkers' Smoking Cessation. https://ClinicalTrials.gov/show/NCT02859142 ; 2016.	Ongoing study
University of C. Impact of Smoking Cessation and γ -Tocopherol to Restore Vascular Endothelial Function. https://ClinicalTrials.gov/show/NCT01314443 ; 2011.	Wrong comparator
University of C. Pilot Study of Nicotine Nasal Spray and Varenicline on Smoking in Methadone-Maintained Patients. https://ClinicalTrials.gov/show/NCT02147132 ; 2016.	Ongoing study
University of California LA, National Institute on Drug A. Combining Varenicline and Naltrexone for Smoking Cessation. https://ClinicalTrials.gov/show/NCT02698215 ; 2016.	Wrong comparator
University of California LA, Tarzana Treatment C. Dopamine D2/D3 Receptor Upregulation by Varenicline in Methamphetamine Users. https://ClinicalTrials.gov/show/NCT03059563 ; 2017.	Ongoing study
University of California LA. Arterial Inflammation and E-Cigarettes. https://ClinicalTrials.gov/show/NCT02734888 ; 2015.	Wrong comparator
University of California LA. E-cigarettes and Blood Vessel Function. https://ClinicalTrials.gov/show/NCT02740595 ; 2015.	Ongoing study
University of California LA. E-Cigarettes and SNA. https://ClinicalTrials.gov/show/NCT02724241 ; 2015.	Ongoing study
University of California LA. E-cigarettes, Nicotine Inhaler, and Blood Vessel Function. https://ClinicalTrials.gov/show/NCT03072628 ; 2017.	Wrong study design

University of California LA. Effects of Varenicline in Heavy Drinking Smokers. https://ClinicalTrials.gov/show/NCT02488889 ; 2015.	Wrong study design
University of California SD, California Department of Health S. Telephone Counseling and the Distribution of Nicotine Patches to Smokers. https://ClinicalTrials.gov/show/NCT00851357 ; 2009.	Ongoing study
University of California SD. Smoking Cessation With the Nicotine Patch: A Pilot Study of Patch Messaging. https://ClinicalTrials.gov/show/NCT01287377 ; 2010.	Participants received same drug
University of California SF, Kaiser P, National Institute on Drug A. Interventions for Smoking Among Persons in Recovery. https://ClinicalTrials.gov/show/NCT00714896 ; 2007.	Wrong study design
University of California SF, National Heart L, Blood I. Chinese Community Smoking Cessation Project. https://ClinicalTrials.gov/show/NCT00139516 ; 2001.	Ongoing study
University of California SF, National Institute on Drug A. Tobacco Intervention in Buprenorphine Treatment. https://ClinicalTrials.gov/show/NCT01350011 ; 2011.	Ongoing study
University of California SF, National Institutes of H. E-cigarette THC Study. https://ClinicalTrials.gov/show/NCT02955329 ; 2017.	Wrong comparator
University of California SF, University of C. Smoking Reduction and Cessation Interventions for Chinese. https://ClinicalTrials.gov/show/NCT00714532 ; 2007.	Participants received same drug
University of California SF. Low-Cost Contingency Management for Smoking Cessation. https://ClinicalTrials.gov/show/NCT01040260 ; 2009.	Participants received same drug
University of California SF. The END Perioperative Smoking Pilot Study. https://ClinicalTrials.gov/show/NCT02482233 ; 2015.	Ongoing study
University of D, Tayside NHS, British Heart F. Vascular EffectS of regUlar Cigarettes Versus electronic Cigarette USe. https://ClinicalTrials.gov/show/NCT02878421 ; 2016.	Ongoing study
University of F, National Institute on Drug A. Smoking Cessation Intervention. https://ClinicalTrials.gov/show/NCT03072511 ; 2016.	Ongoing study
University of I, National Heart L, Blood I. Preventing Weight Gain and Controlling Blood Pressure During Smoking Cessation in Hypertensive Smokers. https://ClinicalTrials.gov/show/NCT00113074 ; 2004.	Participants received same drug
University of Illinois at C, National Institute on Drug A, National Cancer I, Howard Brown Health C. Culturally Targeted & Individually Tailored Smoking Cessation Study: LGBT Smokers. https://ClinicalTrials.gov/show/NCT01633567 ; 2012.	Participants received same drug
University of K, Kentucky Lung Cancer Research P. Optimization of Smoking Cessation Strategies Concurrent With Treatment of Tobacco Related Malignancies. https://ClinicalTrials.gov/show/NCT02048917 ; 2014.	Ongoing study
University of Kansas Medical C, National Cancer I, GlaxoSmithKline. Disease Management for Smoking Cessation. https://ClinicalTrials.gov/show/NCT00440115 ; 2009.	Participants received same drug
University of Kansas Medical C, National Heart L, Blood I. Changing the Default for Tobacco Treatment. https://ClinicalTrials.gov/show/NCT02721082 ; 2016.	Participants received same drug

University of Kansas Medical C, Patient-Centered Outcomes Research I. Smoking Cessation Versus Long-term Nicotine Replacement Among High-risk Smokers. https://ClinicalTrials.gov/show/NCT02148445 ; 2014.	Ongoing study
University of L. The Acute Cardiovascular Effects of Nicotine in E-cigarettes (ACE) Study. https://ClinicalTrials.gov/show/NCT03010995 ; 2017.	Ongoing study
University of M, National Cancer I. Smoking Cessation in Hispanic Construction Workers. https://ClinicalTrials.gov/show/NCT02873377 ; 2017.	Ongoing study
University of M. Optimizing Smoking Cessation for People With HIV/AIDS Who Smoke. https://ClinicalTrials.gov/show/NCT02460900 ; 2015.	Ongoing study
University of Massachusetts B. Smoking Cessation Intervention for Women Living With HIV. https://ClinicalTrials.gov/show/NCT02898597 ; 2016.	Participants received same drug
University of Massachusetts W. Tobacco Dependence Treatment for Asian Americans. https://ClinicalTrials.gov/show/NCT01091363 ; 2009.	Participants received same drug
University of Minnesota C, Translational Science I, National Cancer I. Offering Proactive Tobacco Treatment. https://ClinicalTrials.gov/show/NCT01123967 ; 2011.	Linked to excluded study
University of Minnesota C, Translational Science I, National Heart L, Blood I. Enhancing Smoking Cessation in the Homeless Population. https://ClinicalTrials.gov/show/NCT01932996 ; 2015.	Study withdrawn
University of Minnesota C, Translational Science I, National Institute on Drug A, National Cancer I. Efficacy of Oral Tobacco Products Compared to a Medicinal Nicotine. https://ClinicalTrials.gov/show/NCT00710034 ; 2010.	Wrong comparator
University of Minnesota C, Translational Science I, National Institute on Drug A, Pfizer. Human Laboratory Study of Varenicline and Bupropion for Nicotine Dependence. https://ClinicalTrials.gov/show/NCT00749658 ; 2008.	Wrong study design
University of Minnesota C, Translational Science I, National Institute on Drug A, Pfizer. Human Laboratory Study Of Varenicline in Smokers. https://ClinicalTrials.gov/show/NCT00709696 ; 2008.	Ongoing study
University of Minnesota C, Translational Science I, National Institute on Drug A. Tobacco Cessation in Postmenopausal Women (Part II) - 2. https://ClinicalTrials.gov/show/NCT00061074 ; 1993.	Wrong study design
University of Minnesota C, Translational Science I. SMART for Smoking Cessation in Lung Cancer Screening. https://ClinicalTrials.gov/show/NCT02597491 ; 2015.	Participants received same drug
University of N, Department of Health UK. Trial Comparing Different Methods of Support With Stopping Smoking (PORTSSS/Stop Together Trial). https://ClinicalTrials.gov/show/NCT00775944 ; 2009.	Wrong study design
University of O. Smoking Cessation in Women With Gynecological Conditions. https://ClinicalTrials.gov/show/NCT01989923 ; 2013.	Ongoing study
University of P, National Institute on Drug A, National Institutes of H. Varenicline and Smoking Behavior. https://ClinicalTrials.gov/show/NCT01524627 ; 2011.	Ongoing study
University of P, National Institute on Drug A. Cessation in Non-Daily Smokers. https://ClinicalTrials.gov/show/NCT02168855 ; 2014.	Ongoing study
University of P, National Institute on Drug A. Treatment of Smoking Among Individuals With PTSD. https://ClinicalTrials.gov/show/NCT00937235 ; 2009.	Participants received same drug
University of P, National Institute on Drug A. Varenicline for Nicotine Dependence Among Those With HIV/AIDS. https://ClinicalTrials.gov/show/NCT01710137 ; 2012.	Ongoing study

University of P, Pfizer. Measuring Smoking Behaviors While Using Varenicline. https://ClinicalTrials.gov/show/NCT00948155 ; 2009.	Wrong study design
University of P, Vitality I. Randomized Clinical Trial to Reduce Harm From Tobacco. https://ClinicalTrials.gov/show/NCT02328794 ; 2015.	Ongoing study
University of P. Extended Varenicline Treatment for Smoking Among Cancer Patients. https://ClinicalTrials.gov/show/NCT01756885 ; 2013.	Ongoing study
University of P. Nicotine Receptor Levels and Smoking Cessation. https://ClinicalTrials.gov/show/NCT01788943 ; 2013.	Ongoing study
University of P. Project 1, Study 2: The Combined Impact of Nicotine Replacement and Spectrum Cigarettes. https://ClinicalTrials.gov/show/NCT02301325 ; 2014.	Wrong comparator
University of Sao P. Effect of Smoking Cessation on Non-surgical Periodontal Therapy. https://ClinicalTrials.gov/show/NCT02493270 ; 2007.	Wrong study design
University of T, National Cancer I. Strategies to Promote Cessation in Smokers Who Are Not Ready To Quit. https://ClinicalTrials.gov/show/NCT02905656 ; 2016.	Ongoing study
University of Texas at A, National Institute of Mental H. Smoking Cessation Following Psychiatric Hospitalization. https://ClinicalTrials.gov/show/NCT02204956 ; 2015.	Ongoing study
University of Texas at A. Enhancing Panic and Smoking Reduction Treatment With D-Cycloserine. https://ClinicalTrials.gov/show/NCT01944423 ; 2013.	Participants received same drug
University of V, National Cancer I. Does Abstinence From E-cigarettes Produce Withdrawal Symptoms? : https://ClinicalTrials.gov/show/NCT02825459 ; 2016.	Wrong study design
University of Wisconsin M, Penn State U, University of Illinois at C. A Comparative Effectiveness RCT of Optimized Cessation Treatments. https://ClinicalTrials.gov/show/NCT02301403 ; 2015.	Ongoing study
University of Wisconsin M, Penn State U, University of Illinois at C. The Long-Term Quitting (Smoking Cessation) Study. https://ClinicalTrials.gov/show/NCT02564315 ; 2015.	Participants received same drug
University of Wisconsin M. UW-CTRI Smoking Cessation Study. https://ClinicalTrials.gov/show/NCT02681510 ; 2016.	Wrong study design
Unrod M, Simmons VN, Sutton SK, Cummings KM, Celestino P, Craig BM, et al. Relapse-prevention booklets as an adjunct to a tobacco quitline: A Randomized Controlled Effectiveness Trial. <i>Nicotine and Tobacco Research</i> . 2016;18(3):298-305.	Participants received same drug
Ura S, Wada H, Satoh-Asahara N, Akao M, Abe M, Ono K, et al. Self-rating depression scale score as an inverse and independent predictor of successful smoking cessation with a pharmacological treatment for nicotine addiction. <i>Circulation Conference: American Heart Association's</i> . 2014;130(no pagination).	Cannot locate full text
van Rossem C, Spigt M, Smit ES, Viechtbauer W, Mijnheer KK, van Schayck CP, et al. Combining intensive practice nurse counselling or brief general practitioner advice with varenicline for smoking cessation in primary care: Study protocol of a pragmatic randomized controlled trial. <i>Contemporary Clinical Trials</i> . 2015;41:298-312.	Ongoing study
van Staden SR, Groenewald M, Engelbrecht R, Becker PJ, Hazelhurst LT. Carboxyhaemoglobin levels, health and lifestyle perceptions in smokers converting from tobacco cigarettes to electronic cigarettes. <i>S Afr Med J</i> . 2013;103(11):865-8.	Wrong study design
Vansickel AR, Cobb CO, Weaver MF, Eissenberg TE. A Clinical Laboratory Model for Evaluating the Acute Effects of Electronic "Cigarettes": Nicotine Delivery Profile and Cardiovascular and Subjective Effects. <i>Cancer Epidemiology Biomarkers & Prevention</i> . 2010;19(8):1945-53.	Wrong study design

Veldheer S, Yingst J, Midya V, Hummer B, Lester C, Krebs N, et al. Pulmonary and other health effects of electronic cigarette use among adult smokers participating in a randomized controlled smoking reduction trial. <i>Addictive Behaviors</i> . 2018;29:29.	Ongoing study
Vickerman KA, Keller PA, Deprey M, Lachter RB, Jenssen J, Dreher M. Never Quit Trying: Reengaging Tobacco Users in Statewide Cessation Services. <i>Journal of Public Health Management and Practice</i> . 2018;24(3):E25-E33.	Wrong study design
Vidrine DJ, Arduino RC, Gritz ER. Impact of a cell phone intervention on mediating mechanisms of smoking cessation in individuals living with HIV/AIDS. <i>Nicotine & Tobacco Research</i> . 2006;8(Suppl 1):S103-S8.	Participants received same drug
Vidrine JI, Businelle MS, Reitzel LR, Cao Y, Cinciripini PM, Marcus MT, et al. Coping mediates the association of mindfulness with psychological stress, affect, and depression among smokers preparing to quit. <i>Mindfulness</i> . 2014:[ePub ahead of print].	Participants received same drug
Vijay Ramchandani PD, National Institute on Alcohol A, Alcoholism, National Institutes of Health Clinical C. Varenicline to Reduce Alcohol Consumption in Heavy Drinkers. https://ClinicalTrials.gov/show/NCT00695500 ; 2008.	Wrong population
Villardaga R, Heffner JL, Mercer LD, Bricker JB. Do counselor techniques predict quitting during smoking cessation treatment? A component analysis of telephone-delivered Acceptance and Commitment Therapy. <i>Behaviour Research and Therapy</i> . 2014;61:89-95.	Linked to excluded study
Virginia Commonwealth U, National Institute on Drug A. Examining the Effect of the Nicotine Patch in Male and Female Smokers - 3. https://ClinicalTrials.gov/show/NCT00390559 ; 2005.	Wrong study design
Virtanen SE, Zeebari Z, Rohyo I, Galanti MR. Evaluation of a brief counseling for tobacco cessation in dental clinics among Swedish smokers and snus users. A cluster randomized controlled trial (the FRITT study). <i>Preventive Medicine</i> . 2015;70:26-32.	Wrong or missing intervention
Wake Forest University Health S, National Cancer I. Bupropion Hydrochloride or Patient's Choice for Smoking Cessation in Patients With Squamous Cell Head and Neck Cancer Undergoing Radiation Therapy With or Without Chemotherapy. https://ClinicalTrials.gov/show/NCT02582008 ; 2016.	Ongoing study
Wakefield M, Olver I, Whitford H, Rosenfeld E. Motivational interviewing as a smoking cessation intervention for patients with cancer: Randomized controlled trial. <i>Nursing Research</i> . 2004;53(6):396-405.	Wrong comparator
Walker N, Howe C, Bullen C, Grigg M, Glover M, McRobbie H, et al. Does improved access and greater choice of nicotine replacement therapy affect smoking cessation success? Findings from a randomized controlled trial. <i>Addiction</i> . 2011;106(6):1176-85.	Wrong study design
Walker N, Howe C, Glover M, McRobbie H, Barnes J, Nosa V, et al. Cytisine versus nicotine for smoking cessation. <i>New England Journal of Medicine</i> . 2014;371(25):2353-62.	Wrong comparator
Walker N, Verbiest M, Kurdziel T, Laking G, Laugesen M, Parag V, et al. Effectiveness and safety of nicotine patches combined with e-cigarettes (with and without nicotine) for smoking cessation: study protocol for a randomised controlled trial. <i>BMJ Open</i> . 2019;9(2):e023659.	Ongoing study
Walsh MM, Hilton JF, Ellison JA, Gee L, Chesney MA, Tomar SL, et al. Spit (smokeless) tobacco intervention for high school athletes: Results after 1 year. <i>Addictive Behaviors</i> . 2003;28(6):1095-113.	Wrong population
Walsh MM, Langer TJ, Kavanagh N, Mansell C, MacDougal W, Kavanagh C, et al. Smokeless tobacco cessation cluster randomized trial with rural high school males: Intervention interaction with baseline smoking. <i>Nicotine & Tobacco Research</i> . 2010;12(6):543-50.	Wrong population
Wang X, Wu I, Chen H, Bordnick P, Essien EJ, Johnson ML, et al. Comparison of cardiovascular risks following smoking cessation treatments using varenicline versus NRT among schizophrenic smokers. <i>Value in Health</i> . 2016;19 (3):A182-A3.	Cannot locate full text
Wang X, Wu I, Chen H, Bordnick P, Johnson ML, Essien EJ, et al. Comparison of suicide attempts/behaviors following smoking cessation treatments among schizophrenic smokers. <i>Value in Health</i> . 2016;19 (3):A111.	Cannot locate full text

Wang X, Yang M, Chen H, Essien EJ, Abughosh S. Predeictors of a successful cessation outcome among adult smokers using smoking cessation medication. Value in Health. 2014;17 (3):A180.	Cannot locate full text
Warner C, Sewali B, Olayinka A, Eischen S, Wang Q, Guo H, et al. Smoking cessation in homeless populations: Who participates and who does not. Nicotine and Tobacco Research. 2014;16(3):369-72.	Linked to excluded study
Washington University School of M, National Institutes of H, National Institute on Drug A. Genetically Informed Smoking Cessation Trial. https://ClinicalTrials.gov/show/NCT02351167 ; 2015.	Ongoing study
Wayne State U, National Institute on Drug A. Smoking Cessation for People Living With HIV/AIDS. https://ClinicalTrials.gov/show/NCT01965405 ; 2013.	Ongoing study
Weaver KE, Kaplan S, Griffin L, Urbanic J, Zbikowski S, Danhauer SC. Satisfaction with a Quitline-based Smoking Cessation Intervention among Cancer Survivors. Cancer Epidemiology, Biomarkers & Prevention. 2015;24(4):759.	Participants received same drug
Weaver KE, Urbanic JJ, Case D, Kaplan SG, Lesser GJ, Zbikowski S, et al. Preliminary efficacy of an enhanced quitline smoking cessation intervention for cancer patients. Journal of Clinical Oncology Conference. 2015;33(15 SUPPL. 1).	Participants received same drug
Webb A. Safety of nicotine replacement therapy in critically ill patients. Anaesthesia and Intensive Care. 2017;45(3):423-4.	Awaiting classification
Wells QS, Freiberg MS, Greevy RA, Jr., Tyndale RF, Kundu S, Duncan MS, et al. Nicotine Metabolism-informed Care for Smoking Cessation: A Pilot Precision RCT. Nicotine & Tobacco Research. 2017;14:14.	Wrong study design
West R, May S, McEwen A, McRobbie H, Hajek P, Vangeli E. A randomised trial of glucose tablets to aid smoking cessation. Psychopharmacology. 2010;207(4):631-5.	Wrong study design
West R, Zhou X. Is nicotine replacement therapy for smoking cessation effective in the "real world"? Findings from a prospective multinational cohort study. Thorax. 2007;62(11):998-1002.	Wrong study design
West Virginia U, Pfizer, National Institute on Drug A. Smoking Cessation & Opioid Dependence Treatment Integration. https://ClinicalTrials.gov/show/NCT02854800 ; 2016.	Wrong study design
Westman JG, Gowin JL, Vatsalya V, Schwandt ML, Momenan R, Bartlett SE, et al. Varenicline diminishes amygdalar bold response to negative facial expression cues in heavy drinkers. Alcoholism: Clinical and Experimental Research. 2015;39:149A.	Cannot locate full text
Wewers ME, Ferketich AK, Harness J, Paskett ED. Effectiveness of a nurse-managed, lay-led tobacco cessation intervention among ohio appalachian women. Cancer Epidemiology, Biomarkers & Prevention. 2009;18(12):3451-8.	Participants received same drug
Wiggers LCW, Smets EMA, Oort FJ, Storm-Versloot MN, Vermeulen H, van Loenen LBM, et al. Adherence to nicotine replacement patch therapy in cardiovascular patients. International Journal of Behavioral Medicine. 2006;13(1):79-88.	Wrong study design
Williams D, Ussher M, Dunsiger S, Miranda R, Gwaltney C, Monti P, et al. Overcoming limitations in previous research on exercise as a smoking cessation treatment: rationale and design of the "Quit for Health" trial2014; 37(1):[33-42 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/642/CN-01015642/frame.html .	Participants received same drug
Williams NJ, Arheart KL, Klesges R. A SMOKELESS TOBACCO CESSATION PROGRAM FOR POSTSECONDARY STUDENTS. Health Values. 1995;19(3):33-42.	Wrong or missing intervention

Williams NJ. A smokeless tobacco cessation program for postsecondary students [thesis]. Dissertation Abstracts International. 1992;53(6-A):1821.	Wrong or missing intervention
Wilson JS, Elborn JS, Fitzsimons D. 'It's not worth stopping now': why do smokers with chronic obstructive pulmonary disease continue to smoke? A qualitative study. <i>Journal of Clinical Nursing</i> . 2011;20(5-6):819-27.	Wrong study design
Wing VC, Wass CE, Bacher I, Rabin RA, George TP. Varenicline modulates spatial working memory deficits in smokers with schizophrenia. <i>Schizophrenia Research</i> . 2013;149(1-3):190-1.	Wrong study design
Winickoff JP, Nabi-Burza E, Chang Y, Finch S, Regan S, Wasserman R, et al. Implementation of a Parental Tobacco Control Intervention in Pediatric Practice. <i>Pediatrics</i> . 2013;132(1):109.	Wrong study design
Wisborg K, Henriksen TB, Jespersen LB, Secher NJ. Nicotine patches for pregnant smokers: A randomized controlled study. <i>Obstetrics and Gynecology</i> . 2000;96(6):967-71.	Wrong population
Wittebole X, Hahm S, Coyle SM, Kumar A, Calvano SE, Lowry SF. Nicotine exposure alters in vivo human responses to endotoxin. <i>Clinical & Experimental Immunology</i> . 2007;147(1):28-34.	Wrong population
Won Choi PMPH, National Cancer I, University of Kansas Medical C. Smoking Cessation for American Indians. https://ClinicalTrials.gov/show/NCT01106456 ; 2010.	Wrong study design
Wong R, Tuckett G. Kick the habit: A pharmacist-managed tobacco cessation clinic. <i>Journal of the American Pharmacists Association</i> . 2015;55 (2):e231.	Cannot locate full text
Wu BJ, Chen HK, Lee SM. Do atypical antipsychotics really enhance smoking reduction more than typical ones?: the effects of antipsychotics on smoking reduction in patients with schizophrenia. <i>Journal of Clinical Psychopharmacology</i> . 2013;33(3):319-28.	Awaiting classification
Wu D, Ma GX, Zhou K, Zhou D, Liu A, Poon AN. The effect of a culturally tailored smoking cessation for Chinese American smokers. <i>Nicotine & Tobacco Research</i> . 2009;11(12):1448-57.	Participants received same drug
Wu I, Chen H, Bordnick P, Essien EJ, Johnson ML, Peters RJ, et al. Comparative effectiveness of smoking cessation medications among schizophrenic smokers. <i>Value in Health</i> . 2016;19 (3):A119-A20.	Cannot locate full text
Wyman J, Allen A, Hertsgaard L, Overson E, Allen S, Hatsukami D. Effect of smoking cessation on overactive bladder symptoms in adults: A pilot study 2014; 33(6):[866-7 pp.]. Available from: http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/235/CN-01059235/frame.html .	Wrong comparator
Yalcin BM, Unal M. The mid-term effect of grandchildren and children motivation to quit rates of smokers. <i>Journal of Experimental and Clinical Medicine (Turkey)</i> . 2017;34(1):73-7.	Wrong study design
Yale U, Columbia U. Varenicline Treatment of Alcohol Dependence in Smokers. https://ClinicalTrials.gov/show/NCT01553136 ; 2012.	Ongoing study
Yale U, National Cancer I. Progesterone Augmentation of Nicotine Replacement Therapy Study. https://ClinicalTrials.gov/show/NCT02685072 ; 2016.	Wrong comparator
Yale U, National Institute on Alcohol A, Alcoholism. Do Treatments for Smoking Cessation Affect Alcohol Drinking? Study 1: Nicotine Replacement Therapy. https://ClinicalTrials.gov/show/NCT00699556 ; 2006.	Wrong comparator
Yale U, National Institute on Alcohol A, Alcoholism. Do Treatments for Smoking Cessation Affect Alcohol Drinking? Study 2: Do Varenicline (Chantix) and Bupropion (Zyban) Change Alcohol Drinking? : https://ClinicalTrials.gov/show/NCT00580645 ; 2007.	Ongoing study
Yale U, National Institute on Drug A. Smoking Cessation for Smokers With Sleep Problems. https://ClinicalTrials.gov/show/NCT01277887 ; 2010.	Participants received same drug

Yale U, National Institutes of H, National Institute on Drug A. Enhancing a High School Based Smoking Cessation Program. https://ClinicalTrials.gov/show/NCT01145001 ; 2009.	Wrong population
Yale U, Research VAOo, Development, National Institute on Drug A. Varenicline and Nicotine Interactions in Humans (VA). https://ClinicalTrials.gov/show/NCT00606892 ; 2007.	Wrong study design
Yale U. A Smoking Cessation Intervention for Yale Dining Employees. https://ClinicalTrials.gov/show/NCT02562521 ; 2015.	Wrong study design
Yale U. Effect of Varenicline on Fixed-Dose Alcohol Administration in Participants With Alcohol Use Disorders. https://ClinicalTrials.gov/show/NCT01000987 ; 2009.	Ongoing study
Yale U. New Approaches to Smoking Cessation in Heavy Drinkers. https://ClinicalTrials.gov/show/NCT02151591 ; 2014.	Participants received same drug
Yale U. Novel Treatments to Enhance Smoking Cessation Before Cancer Surgery: Effects on Smoking and Surgical Outcomes. https://ClinicalTrials.gov/show/NCT01479998 ; 2011.	Study terminated
Yale U. Personalized Intervention Program: Tobacco Treatment for Patients at Risk for Lung Cancer (PIP). https://ClinicalTrials.gov/show/NCT02658032 ; 2016.	Participants received same drug
Yale U. Smoking Cessation Treatment for Head and Neck Cancer Patients. https://ClinicalTrials.gov/show/NCT00931021 ; 2009.	Study terminated
Yang M, Chen H, Johnson ML, Essien EJ, Peters RJ, Abughosh S. Comparative effectiveness of smoking cessation medications among obese smokers. <i>Value in Health</i> . 2013;16 (3):A111.	Cannot locate full text
Yaqubi A, Chang Y, Berkowitz SA, McGlave C, Rigotti NA, Baggett TP. Subsistence difficulties predict worse abstinence outcomes among homeless smokers: Secondary analysis of a randomized smoking cessation trial. <i>Journal of General Internal Medicine</i> . 2017;32 (2 Supplement 1):S321.	Linked to excluded study
Yayla BA, Yildiz AI, Cosgun G, Ugurlu E, Evyapan F. Factors affecting to quit smoking in an outpatient smoking cessation clinic. <i>European Respiratory Journal Conference: European Respiratory Society Annual Congress</i> . 2011;38(no pagination).	Cannot locate full text
Yoshida N, Nakamura K, Kuroda D, Baba Y, Miyamoto Y, Iwatsuki M, et al. Preoperative Smoking Cessation is Integral to the Prevention of Postoperative Morbidities in Minimally Invasive Esophagectomy. <i>World J Surg</i> . 2018;42(9):2902-9.	Wrong or missing intervention
Yoshii C, Kunitomo F, Kano M, Nishida C, Kawanami Y, Mukae H. Evaluation of one-year outcome of smoking cessation among patients who successfully finished a 12-week smoking cessation program. <i>Respirology</i> . 2012;17:147.	Cannot locate full text
Young-Wolff KC, Kline-Simon AH, Das S, Mordecai DJ, Miller-Rosales C, Weisner C. Smoking trends among adults with behavioral health conditions in integrated health care: A retrospective cohort study. <i>Psychiatric Services</i> . 2016;67(9):996-1003.	Wrong study design
Yu H, Zang Y, Lin J. The effect of the abstinence from smoking with nicotine replacement therapy combining with psychological and behavior intervention. <i>Chinese Journal of Rehabilitation Medicine</i> . 2006;21(12):1104-6.	Participants received same drug
Yujie W, Huliang L. GW25-e5131 Efficacy and Safety of Varenicline for Smoking Cessation in Patients with CAD undergoing PCI. <i>Journal of the American College of Cardiology</i> . 2014;64(16_S).	Cannot locate full text

Zahid R, Dogar O, Mansoor S, Khan A, Kanaan M, Jawad M, et al. The efficacy of varenicline in achieving abstinence among waterpipe tobacco smokers - study protocol for a randomized controlled trial 2017; 18(1):[14 p.]. Available from: http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/947/CN-01454947/frame.html .	Ongoing study
Zawertailo L, Mansoursadeghi-Gilan T, Zhang H, Hussain S, Le Foll B, Selby P. Varenicline and Bupropion for Long-Term Smoking Cessation (the MATCH Study): Protocol for a Real-World, Pragmatic, Randomized Controlled Trial. <i>JMIR Research Protocols</i> . 2018;7(10):e10826.	Ongoing study
Zesiewicz TA, Greenstein PE, Sullivan KL, Wecker L, Miller A, Jahan I, et al. A randomized trial of varenicline (Chantix) for the treatment of spinocerebellar ataxia type 3. <i>Neurology</i> . 2012;78(8):545-50.	Wrong population
Zhang Y, Yang Y, Yue Y, Wang DX. Impact of prior smoking cessation on postoperative pulmonary complications in the elderly: secondary analysis of a prospective cohort study. <i>Eur J Anaesthesiol</i> . 2017;34(12):853-4.	Linked to excluded study
Zhao Q, Schwam E, Fullerton T, O'Gorman M, Burstein AH. Pharmacokinetics, Safety, and Tolerability Following Multiple Oral Doses of Varenicline Under Various Titration Schedules in Elderly Nonsmokers. <i>Journal of Clinical Pharmacology</i> . 2011;51(4):492-501.	Wrong population
Zhu S-H, Gamst A, Lee M, Cummins S, Yin L, Zoref L. The Use and Perception of Electronic Cigarettes and Snus among the US Population. <i>Plos One</i> . 2013;8(10).	Wrong study design
Zwar N, Richmond R, Halcomb E, Furler J, Smith J, Hermiz O, et al. Quit in General Practice: a cluster randomised trial of enhanced in-practice support for smoking cessation. <i>Bmc Family Practice</i> . 2010;11.	Participants received same drug