
Information on p1 obtained from clinical assessor & medical notes Interviewer (initials) Sex Male 1 Female 2 Age Date of interview Date of overdose (if staggered, record latest date) NK 9 Staggered overdose No 0 Yes 1 Time of overdose Overdose included self injury No 0 Yes 1 If YES, Details of self injury Substances taken in overdose Number of tablets Paracetamol ..... ..... ..... Paracetamol level @ 4 hours mmols/L Delayed presentation No 0 Yes 1 NK 9 Alcohol taken as part of the act No 0 Yes 1 NK 9 Alcohol in 6 hours before NK 9 No 0 Yes 1

Number of units (1 unit =  $\frac{1}{2}$  pt beer, 1 small glass wine, 1 measure spirits)

Medical treatment (from casenotes/ward staff)

None 0, Activated charcoal 1, Parvalax 2, Activated charcoal + parvalax 3, Other 4 (specify)





Interview Schedule

1. If patient took paracetamol compound as well as paracetamol::

Did yo	u know that x	contained para	acetamol?
	No 0	Yes 1	NK 9
If NO 1.1	Would you have	e taken them as v	vell as the other paracetamol if you had known?
	No 0	Yes 1	NK 9
If 1.1 = 1.2	= 0 or 1 <b>Why?</b>		

2. **How long was there between first seriously thinking about this overdose and actually doing it?** (*i.e. when patient actually planned the overdose*)

1	0 - < 15 minutes		5	12 - < 24 hours
2	15 - < 60 minutes	6	1 - <	7 days
3	1 - < 3 hours		7	7 – 28 days
4	3 - < 12 hours		8	> 1  month
			9	NK

# 3. **Over what period of time did you take the tablets?**

1	< 5 minutes		4	30 - < 60 minutes
2	5 - < 15 minutes		5	1 - < 3 hours
3	15 - < 30 minutes	6	3 hou	urs +

If longer than 3 hours, RECORD DETAILS.

### 4. Did you take all the paracetamol tablets that were available?

1	All available	3	Minority $(<^{1}/_{2})$
2	Majority $(> \frac{1}{2})$	4	About half

#### How many were available?

RECORD DETAILS: Number of packs, pack size

# 5. Where did you get the tablets?

No 0		Yes 1	NK	9		RECO	RD ALL (ca	n be more th	an one)	
Alread Bought the act	t for * <u>-</u>	chemist supermarke local shop .	7 below t (e.g. garage)							
Friend										
*RECO	ORD AL	L DETAIL	S							
Specifi	c Outlet		Quantity	When bo	ought	Who	bought	> recomm amour		
If > ma	aximum	pack size (3	2 tablets pharr	nacy, 16 tab	lets other	outlet):				
6,	Cher	nists have b	een advised n	ot to sell m	ore than .	32 table	ts /			
	Supe	Supermarkets /garages /shops have been advised not to sell more than 16 tablets								
	Did y	ou try to b	uy any more t	han that at	any one o	outlet?				
	No	)	Yes 1	NK 9						
	If YE 6.1		appened? (i.e.	was there an	ny interve	ntion by	salesperson/	/ pharmacist	?)	
7.	If alr	eady in the l	nome:							
	Whe	re were the	tablets at the	time of the	overdose	?				
	1 2 3	Bathroor Kitchen Bedroom	n cabinet			Other (s 5 9	pecify) More than o NK	one place		
8.		-	fically for the c e <b>you took the</b>		-		,			
9.	Why	were the ta	blets obtained	1?						
	1 2	For the O Other rea	DD ason (specify)			9	NK			
10.	Why	did you ch	oose paraceta	mol for the	o/d?					
	DEC	ים מר								

RECORD:

1. People take overdoses for all sorts of reasons. I'm going to show you some cards with different reasons on them. Please could you look through them and choose any which might explain why you took your overdose. You can choose as many as you like.

#### **YELLOW CARDS**

No 0 Yes 1

I wanted to show how desperate I was feeling I wanted to die I wanted to escape from an unbearable situation I wanted to influence someone I wanted to make someone feel sorry I wanted to get relief from a terrible state of mind I wanted to find out whether someone really loved me I wanted to get help

#### 11.1 Were there any other reasons?

Yes 1 No 0

IF YES, RECORD:

12. Before you took the tablets, what did you think the physical effects of taking an overdose of paracetamol might be?

SPECIFY:

13. I'm going to show you some cards which show possible outcomes from taking an overdose - please could you choose the ONE which best describes what you thought the effects of the paracetamol overdose might have been if you had not received treatment

#### **BLUE CARDS**

- Completely safe 1
- 2 Only mild effects which are short lived
- 3 May be harmful but the effects are not lasting
- 4 May cause permanent damage or harm
- 5 Could kill you if not treated in time
- 6 Don't know

9999

If 3, 4 or 5

What are the dangers? 13.1

#### 13.2 How many tablets of paracetamol do you think could cause death?

NK

**RECORD NUMBER** 

NA (i.e. thought not to be harmful) 8888

### 14. IF NECESSARY, ASK Do you think paracetamol could

## 1. Harm a part of your body?

No 0 Yes 1 NK 9 IF YES, SPECIFY:

# 2. Cause you to lose consciousness?

No 0 Yes 1 NK 9

# 3. Have other effects?

No	0	Yes	1	NK	9
IF YES	S, SPECIFY:				

# 15. What have you been told since you came into hospital about the dangers of an overdose of paracetamol? Who told you?

SPECIFY:

#### 16. IF PATIENT HAS INDICATED THAT THEY KNOW THERE ARE HARMFUL EFFECTS: How quickly do the harmful effects show themselves?

1	At once	3 > 6 up to 24 hours after the o/d
2	1-6 hours after the o/d	4 > 24 hours after the o/d
		9 DK

#### 17. IF UNAWARE OF DANGERS: Sometimes despite treatment paracetamol overdose can cause serious harm, even death

Knowing this, would you still have taken paracetamol?

No 0 Yes 1 NK 9

18. In fact when a paracetamol overdose causes harm, this is usually after several days, not at once, and during this period the person remains conscious for most of the time.

Knowing this, would you still have taken paracetamol?

No 0 Yes 1 NK 9

re	ecomn		se, and to seek		lling you not to t te medical advic		
D	)id yo	u notice th	e warning on t	the pack?			
N	lo	0	Yes	1	NK	9	
	f YES: 9.1		nave any effect	on your c	lecision to take	the tablets?	
N	lo	0	Yes	1	NK	9	
D	DETAI	LS:					
	NO: 0.2	If you be	d seen it woul	d vou hav	e taken the table	ats?	
	V.2	0	Yes	•	NK		
1	i U	U	1 05	1	INK	2	
If	the p	aracetamo	l pack contain	ed fewer t	tablets, would ye	ou still have ta	ken an overdo

 No
 0
 Yes
 1
 NK
 9

 If NO:
 20.1
 Why not?

# 21. PREVIOUS OVERDOSES

# Have you taken any overdoses before, or tried to harm yourself in any other way?

(Note incidences of self harm not involving overdose, but record overdoses in detail)

RECORD DETAILS OF OVERDOSES:

Total number:

When

Drugs taken

# 22. IF PREVIOUSLY TOOK PARACETAMOL: What was the outcome?

If over 5, record information on 1<sup>st</sup> and last 4 episodes

No	0	Yes	1	NA	8		NK	9
				1 <sup>st</sup> ever	Most recent	next	next	next
No mee	lical contact							
Non-ho	ospital (e.g. GP)							
Genera	l hospital, not adn	nitted						
Genera	l hospital, admitte	d						
Special	ist treatment, e.g.	liver unit						

23. For MOST RECENT *previous* paracetamol overdose: (if not known already) **How many tablets did you take?** 

# RECORD NUMBER

24. Did your previous experience affect your decision to use paracetamol this time?

No	0	Yes	1	NA	8	NK	9
If YES 24.1	In what way?						
RECO	RD:						

# 25. If you ever felt like taking an overdose again, would you take paracetamol?

No 0 Yes 1 NK 9

Why?

RECORD DETAILS:

# 26. Do you know anyone who has taken a paracetamol overdose?

No 0 Yes 1 NK 9

If YES 26.1 **Who?** 

RECORD DETAILS: Who, when, outcome

- 1 Friend
- 2 Relative
- 3 Other (specify)
- 27. Have you seen or heard about anything in the media (TV, films, news) about people using paracetamol for overdoses?

No 0 Yes 1 NK 9

**RECORD DETAILS:** 

If YES

27.1 Did this influence you in any way?

28.Do you use the internet?<br/>Have you come across anything on the internet about paracetamol?No0Yes1NA (don't use internet)89

RECORD DETAILS If YES 28.1 **Did this influence you in any way?** 

Before we finish, do you have any further comments you would like to make which might help us understand a bit more about why you chose this method, or about anything else which has been raised?

NK

Thank you very much